TOPCARE CHILDRENS COLD COUGH- brompheniramine maleate, dextromethorphan hydrobromide solution Topco Associates LLC

Topco Associates LLC. Children's Cold & Cough Drug Facts

Active ingredients (in each 10 mL)

Brompheniramine maleate 2 mg

Dextromethorphan HBr 10 mg

Purposes

Antihistamine

Cough suppressant

Uses

- temporarily relieves:
- cough
- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes due to hay fever (or other upper respiratory allergies)
- temporarily alleviates the intensity of coughing

Warnings

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleepy

Ask a doctor before use if you have

- trouble urinating due to an enlarged prostate gland
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- cough that lasts or is chronic such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers

When using this product

- do not use more than directed
- may cause marked drowsiness
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosage cup provided
- keep dosage cup with product
- mL = milliliter

age	dose
adults and children 12 years over	20 mL every 4 hours
children 6 to under 12 years	10 mL every 4 hours
children under 6 years	do not use

Other information

- each 10 mL contains: sodium 4 mg
- store at 20°-25°C (68°-77°F)

Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol solution

Questions or comments?

1-888-423-0139

Package/Label Principal Display Panel

TopCare_® health

COMPARE TO CHILDREN'S DIMETAPP® COLD & COUGH ACTIVE INGREDIENTS

Safe and Effective Relief without Phenylephrine

children's

Cold & Cough

ANTIHISTAMINE - BROMPHENIRAMINE MALEATE

COUGH SUPPRESSANT - DEXTROMETHORPHAN HRr

Relieves Nasal Symptoms

- Runny Nose
- Sneezing

Plus Other Symptoms


- Itchy, Watery Eyes
- Cough

ALCOHOL FREE

Ages 6 Years and over

GRAPE FLAVOR

4 FL OZ (118 mL)





+TopCare® children's **Cold & Cough**

PARENTS: Learn about teen medicine abuse

www.StopMedicineAbuse.org

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING

(in each 10 mL)

Brompheniramine maleate 2 mg......Antihistamine Dextromethorphan HBr 10 mg......Cough suppressan .Antihistamine

Uses

- temporarily relieves:
- cough

- cough

 runny nose
 sneezing
 itching of the nose or throat
 itchy, watery eyes due to hay fever (or other upper respiratory allergies)
 temporarily alleviates the intensity of coughing

Warnings

Do not use

Do not use

if you are now taking a prescription monoamine
oxidase inhibitor (MAOI) (certain drugs for
depression, psychiatric, or emotional conditions, or
Parkinson's disease), or for 2 weeks after stopping
the MAOI drug. If you do not know if your
prescription drug contains an MAOI, ask a doctor or
pharmacist before taking this product.

■ to make a child sleepy

Ask a doctor before use if you have

- trouble urinating due to an enlarged prostate gland
- gradienta
 cough that occurs with too much phlegm (mucus)
 a breathing problem such as emphysema or chronic bronchitis
- cough that lasts or is chronic such as occurs with ng, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

- When using this product

 do not use more than directed

 may cause marked drowsiness

 a void alcoholic beverages

 alcohol, sedatives, and tranquitizers may increase
- be careful when driving a motor vehicle or operating
- excitability may occur, especially in children

Stop use and ask a doctor if cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health onal before use.

Reep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than 6 doses in any 24-hour period
 measure only with dosage cup provided
 keep dosage cup with product
 ml = millitier

age	dose
adults and children 12 years and over	20 mL every 4 hours
children 6 to under 12 years	10 mL every 4 hours
children under 6 years	do not use

Other information

■ each 10 mL contains: sodium 4 mg ■ store at 20°-25°C (68°-77°F)

Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol solution

Questions or comments? 1-888-423-0139

DISTRIBUTED BY TOPCO ASSOCIATES LLC ITASCA, IL 60143 ©TOPCO PERA0924 QUESTIONS? 1-888-423-0139



Scan here for more information or call 1-888-423-0139 **GLUTEN FREE** QUALITY GUARANTEED

*This product is not manufactured or distributed by Foundation Consumer Healthcare, LLC, distributor of Children's Dimetapp® Cold & Cough.



COMPARE TO CHILDREN'S DIMETAPP® COLD & COUGH ACTIVE INGREDIENTS*

Safe and Effective Relief without Phenylephrine

children's Cold &

ANTIHISTAMINE-BROMPHENIRAMINE MALEATE COUGH SUPPRESSANT - DEXTROMETHORPHAN HBr

Relieves Nasal Symptoms

- Runny Nose
- Sneezing

Plus Other Symptoms

· Itchy, Watery Eve







4 FL 0Z (118 mL)



26K26 88 C1

+TopCare_{*}

Safe and Effective Relief without Phenylephrine

children's Cold &

ANTIHISTAMINE - BROMPHENIRAMINE MALEATE COUGH SUPPRESSANT - DEXTROMETHORPHAN HBr

Relieves Nasal Symptoms

- Runny Nose
- Sneezing

Plus Other Symptoms

- · Itchy, Watery Eve
- Cough

ALCOHOL FREE





4FL 0Z (118mL)



TOPCARE CHILDRENS COLD COUGH

brompheniramine maleate, dextromethorphan hydrobromide solution

Product Information

Product Type HUMAN OTC DRUG **Item Code (Source)** NDC:76162-099

Route of Administration ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
BROMPHENIRAMINE MALEATE (UNII: IXA7C9ZN03) (BROMPHENIRAMINE - UNII: H57G17P2FN)	BROMPHENIRAMINE MALEATE	2 mg in 10 mL	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 10 mL	

Inactive Ingredients			
Ingredient Name	Strength		
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)			
EDETATE DISODIUM (UNII: 7FLD91C86K)			
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)			
FD&C RED NO. 40 (UNII: WZB9127XOA)			
GLYCERIN (UNII: PDC6A3C0OX)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
WATER (UNII: 059QF0KO0R)			
SACCHARIN SODIUM (UNII: SB8ZUX40TY)			
SODIUM BENZOATE (UNII: OJ245FE5EU)			
SORBITOL SOLUTION (UNII: 8KW3E207O2)			

Product Characteristics			
Color		Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

ı	Packaging				
7	# Item Code	Package Description	Marketing Start Date	Marketing End Date	
	NDC:76162-099- 26	1 in 1 CARTON	06/12/2025		
	L	118 mL in 1 BOTTLE; Type 0: Not a Combination Product			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/12/2025	

Labeler - Topco Associates LLC (006935977)

Revised: 6/2025 Topco Associates LLC