

ARTISTRY IDEAL RADIANCE ILLUMINATING CC SPF 50 BROAD SPECTRUM SUNSCREEN MEDIUM- ensulizole, octinoxate, oxybenzone, titanium dioxide, zinc oxide cream
Amway Corp

Artistry Ideal Radiance Illuminating CC SPF 50 Broad Spectrum Sunscreen Medium

Drug Facts

Active Ingredients

Ensulizole 2.0% (W/W P/P) Octinoxate 5.5% (W/W P/P) Oxybenzone 0.5% (W/W P/P)
Titanium Dioxide 0.84% (W/W P/P) Zinc Oxide 9.8% (W/W P/P)

Purpose

Sunscreen

Uses

- Helps prevent sunburn

Warnings

For external use only

Do not use

on broken or damaged skin

When using this product

keep out of eyes. Rinse with water to remove.

Keep out of reach of children.

If product is swallowed, get medical help or contact a Poison Control Center right away.

Other Information

- Protect this product from excessive heat and direct sun

Inactive Ingredients

WATER/EAU/AQUA, DIMETHICONE, CYCLOPENTASILOXANE, METHYL TRIMETHICONE, BUTYLENE GLYCOL, GLYCERIN, LAURYL PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE, TALC, DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER, TRIMETHYLOLPROPANE TRICAPRYLATE/TRICAPRATE, TRIETHANOLAMINE, HDI/TRIMETHYLOL HEXYLLACTONE CROSSPOLYMER, POLYGLYCERYL-6 POLYRICINOLEATE, DIMETHICONE/PEG-10/15 CROSSPOLYMER, PHENOXYETHANOL, SODIUM CHLORIDE, SODIUM CITRATE, PEG/PPG-18/18 DIMETHICONE, ALUMINA, POLYGLYCERYL-3, POLYDIMETHYLSILOXYETHYL DIMETHICONE, TRISILOXANE, HYDROGEN DIMETHICONE, ALUMINUM HYDROXIDE, STEARIC ACID, TRIETHOXYCAPRYLYLSILANE, METHYLPARABEN, DISODIUM EDTA, FRAGRANCE/PARFUM, SILICA, PROPYLPARABEN, BELLIS PERENNIS (DAISY) FLOWER EXTRACT, MALTODEXTRIN, ETHYLPARABEN,, IODOPROPYNYL BUTYLCARBAMATE, JASMINUM OFFICINALE (JASMINE) FLOWER EXTRACT, LECITHIN, SOPHORA JAPONICA FLOWER POWDER, GERANIUM ROBERTIANUM EXTRACT, OENOTHERA BIENNIS (EVENING PRIMROSE) SEED EXTRACT, LACTOBACILLUS/ACEROLA CHERRY FERMENT, GLYCYRRHIZA GLABRA (LICORICE) ROOT EXTRACT, HYDROLYZED CONCHIOLIN PROTEIN, TOCOPHERYL ACETATE, CERAMIDE-3, BETA-SITOSTEROL, TETRAHEXYLDECYL ASCORBATE, MAY CONTAIN/PEUT CONTEINIR: IRON OXIDES (CI 77491, CI 77492, CI 77499), MICA (CI 77019), TITANIUM DIOXIDE (CI 77891).

Directions

For sunscreen use

■ Apply liberally 15 minutes before sun exposure ■ Use a water resistant sunscreen if swimming or sweating or washing ■ Reapply at least every 2 hours ■ Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including: ■ limit time in the sun, especially from 10 a.m. – 2 p.m. ■ wear long-sleeve shirts, pants, hats, and sunglasses

■ Children under 6 months: ask a doctor

Package Labeling:

Exclusively from **AMWAY**

**Drug Facts (continued)/
Faits sur le produit
(continué)/Datos del
Producto (continuación)**

To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including: ■ limit time in the sun, especially from 10 a.m. – 2 p.m. ■ wear long-sleeve shirts, pants, hats, and sunglasses/Mesures de protection solaire. Passer du temps au soleil augmente votre risque de cancer de la peau et de vieillissement prématuré de la peau. Pour diminuer ce risque, utiliser régulièrement un écran solaire avec un FPS de 15 ou plus à large spectre et d'autres mesures de protection solaire, notamment: • limiter le temps passé au soleil, en particulier entre 10 h 00 et 14 h 00 • porter une chemise à manches longues, un pantalon, un chapeau et des lunettes de soleil/Medidas de Protección Solar. Exponerse por períodos largos al sol aumenta su riesgo de cáncer en la piel y envejecimiento prematuro de la piel. Para reducir el riesgo, debe usar filtro solar de amplia gama con FPS 15 o más regularmente y otras medidas de protección contra el sol, incluyendo: ■ limite el tiempo expuesto al sol, especialmente entre las 10 a.m. – 2 p.m. ■ use camisas de manga larga, pantalones, sombreros y lentes de sol

■ Children under 6 months: ask a doctor/ enfants de moins de 6 mois: consulter un médecin/ Menores de 6 meses: consultar con un médico



**ARTISTRY
IDEAL RADIANCE**

Illuminating CC Cream
SPF 50 Broad Spectrum
Sunscreen Cream
CC Crème Luminance
Crème solaire à large
spectre SPF 50

DIN 02434873

30 ml / 1.0 FL OZ

ARTISTRY IDEAL RADIANCE™

Crema Illuminadora CC con
FPS 50 de Amplio Espectro

USE INSTRUCTIONS: APPLY AS A FOUNDATION TO FACE WITH FINGERTIPS AND BLEND. ALWAYS WORN ALONE OR UNDER FOUNDATION FOR FLAWLESS SKIN PERFECTION. / APPLIQUEZ UNIFORMEMENT SUR LE VISAGE OU BOUT DES DOIGTS, TEL UN FOND DE TENUE. PEUT ÊTRE PORTÉE SEULE OU EN DESSOUS D'UN FOND DE TENUE POUR UNE PEAU LISSE ET PARFAITE. / ÚTILICE COMO UNA BASE AL ROSTRO CON LAS PUNTAS DE LOS DEDOS. PUEDE SER USADA SOLA O DEBAJO DE UNA BASE PARA UNA PIEL IMPECABLEMENTE PERFECTA.

**Drug Facts/Faits sur le produit/
Datos sobre el producto**

Active Ingredients / Ingrédients Actifs / Ingredientes Activos

Ensilizole 2.0% (W/W P/P)
Octinoxate 5.5% (W/W P/P)
Oxybenzone 0.5% (W/W P/P)
Titanium Dioxide 0.84% (W/W P/P)
Zinc Oxide 9.8% (W/W P/P)..... Sunscreen/
Ecran solaire/Protector Solar

Uses/Utilisations/Usos

■ Helps prevent sunburn/Aide à prévenir les coups de soleil/Ayuda a prevenir las quemaduras de sol

**Warnings/Avertissements/
Precauciones**

For external use only/Usage externe seulement/
Para uso externo solamente

Do not use on broken or damaged skin. Ne pas utiliser sur une peau éraflée ou abîmée. No aplicar sobre piel lacerada o con daños

When using this product keep out of eyes. Rinse with water to remove. En utilisant ce produit, le tenir à l'écart des yeux. Rincer avec de l'eau pour l'enlever. Al usar este producto, evite el contacto con los ojos. Enjuague con agua para remover.

Stop use and ask a doctor if rash occurs/Cesser l'utilisation et consulter un médecin en cas d'irritation. Deje de usar y consulte con un médico si ocurre irritación

**Drug Facts (continued)/Faits sur le produit
(continué)/Datos del Producto
(continuación)**

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away. Tenir hors de la portée des enfants. En cas d'ingestion du produit, obtenir immédiatement un avis médical ou contacter un centre anti-poison / Mantener fuera del alcance de los niños. Si se ingiere el producto, debe obtener ayuda médica o comunicarse con un Centro de Control de Envenenamientos de inmediato.

**Other Information/Autre information/
Otra Información**

■ Protect this product from excessive heat and direct sun/ Protéger ce produit de la chaleur excessive et d'une exposition directe au soleil/Proteja este producto de calor excesivo y exposición directa al sol

**Inactive ingredients/Ingrédients inactifs/
Ingredientes inactivos :**

WATER/EAU/AQUA, DIMETHICONE, CYCLOPENTASILOXANE, METHYL TRIMETHICONE, BUTYLENE GLYCOL, GLYCERIN, LAURYL PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE, TALC, DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER, TRIMETHYLOLPROPANE TRICAPRYLATE/TRICAPRATE, TRIETHANOLAMINE, HDI/TRIMETHYLOL HEXYL LACTONE CROSSPOLYMER, POLYGLYCERYL-6 POLYRICINOLEATE, DIMETHICONE/PEG-10/15 CROSSPOLYMER, PHENOXYETHANOL, SODIUM CHLORIDE, SODIUM CITRATE, PEG/PPG-18/18 DIMETHICONE, ALUMINA, POLYGLYCERYL-3, POLYDIMETHYLSILOXYETHYL DIMETHICONE, TRISILOXANE, HYDROGEN DIMETHICONE, ALUMINUM HYDROXIDE, STEARIC ACID, TRIETHOXYCAPRYLYLSILANE, METHYL PARABEN, DISODIUM EDTA, FRAGRANCE/PARFUM, SILICA, PROPYL PARABEN, BELLIS PERENNIS (DAISY) FLOWER EXTRACT, MALTODEXTRIN, ETHYL PARABEN, IODO-PROPENYL BUTYL CARBAMATE, JASMINUM OFFICINALE (JASMINE) FLOWER EXTRACT, LECITHIN, SOPHORA JAPONICA FLOWER POWDER, GERANIUM ROBERTIANUM EXTRACT, OENOTHERA BIENNIS (EVENING PRIMROSE) SEED EXTRACT, LACTOBACILLUS/ACEROLIA CHERRY FERMENT, GLYCERYL RHA GLABRA (LICORICE) ROOT EXTRACT, HYDROLYZED CONCHOLIN PROTEIN, TOCOPHERYL ACETATE, CERAMIDE-3, BETA-SITOSTEROL, TETRAHEXYLDECYL ASCORBATE, MAY CONTAIN/ PEUT CONTENIR: IRON OXIDES (CI 77491, CI 77492, CI 77499), MICA (CI 77019), TITANIUM DIOXIDE (CI 77891).

Directions/Mode d'emploi/Instrucciones

For sunscreen use/Pour une utilisation comme écran solaire/
Usa del protector solar

- Apply liberally 15 minutes before sun exposure/Appliquer généreusement 15 minutes avant de s'exposer au soleil/Aplicar abundantemente 15 minutos antes de exponerse al sol
- Use a water resistant sunscreen if swimming or sweating or washing/Utiliser un écran solaire résistant à l'eau pour nager ou si vous transpirez ou vous lavez/Usa un filtro solar resistente al agua si va a nadar o sudar
- Reapply at least every 2 hours/Réappliquer au moins toutes les 2 heures/Vuelva a aplicar cada dos horas.
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging.



11-9336 MEDIUM

no print area



1043014b

™TRADEMARK LICENSED BY ALTICOR INC./MARQUE DE COMMERCE UTILISÉE SOUS LICENCE PAR ALTICOR INC. MADE IN U.S.A./HECHO EN E.U.A.
DIST. BY/PAR AMWAY CORP., 7575 FULTON AVE., MI 48355 USA/E-U • AMWAY CANADA CORPORATION, LONDON, ON N6Y 5V6 IMP. Y/O DIST. POR: AMWAY DOMINICAN REPUBLIC LLC FABRICADO EN E.U.A. POR ACCESS BUSINESS GROUP LLC, 7575 FULTON STREET EAST, ADA, MI 48355 1043014b

PRINT SIDE UP
BCF8942, DATED 03-07-2014
1 3/8 x 1 1/16 x 5

ARTISTRY IDEAL RADIANCE ILLUMINATING CC SPF 50 BROAD SPECTRUM SUNSCREEN MEDIUM

ensilizole, octinoxate, oxybenzone, titanium dioxide, zinc oxide cream

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:50390-002 |
| Route of Administration | TOPICAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|--------------------------|-----------------|
| ENSULIZOLE (UNII: 9YQ9DI1W42) (ENSULIZOLE - UNII:9YQ9DI1W42) | ENSULIZOLE | 20 mg in 1 mL |
| OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51) | OCTINOXATE | 55 mg in 1 mL |
| OXYBENZONE (UNII: 95OOS7VE0Y) (OXYBENZONE - UNII:95OOS7VE0Y) | OXYBENZONE | 5 mg in 1 mL |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP) | TITANIUM DIOXIDE | 8.8 mg in 1 mL |
| ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC CATION - UNII:13S1S8SF37) | ZINC CATION | 98 mg in 1 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|--|-----------------|
| WATER (UNII: 059QF0KO0R) | |
| DIMETHICONE (UNII: 92RU3N3Y1O) | |
| CYCLOMETHICONE 5 (UNII: 0THT5PCI0R) | |
| METHYL TRIMETHICONE (UNII: S73ZQI0GXM) | |
| BUTYLENE GLYCOL (UNII: 3XUS85K0RA) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| LAURYL PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE (UNII: 25G622K2RA) | |
| TALC (UNII: 7SEV7J4R1U) | |
| TRIMETHYLOLPROPANE TRICAPRYLATE/TRICAPRATE (UNII: N1TRQ06KVJ) | |
| TROLAMINE (UNII: 9O3K93S3TK) | |
| HEXAMETHYLENE DIISOCYANATE/TRIMETHYLOL HEXYLLACTONE CROSSPOLYMER (UNII: WB5K9Y35Y9) | |
| PHENOXYETHANOL (UNII: HIE492ZZ3T) | |
| SODIUM CHLORIDE (UNII: 451W47IQ8X) | |
| SODIUM CITRATE (UNII: 1Q73Q2JULR) | |
| PEG/PPG-18/18 DIMETHICONE (UNII: 9H0AO7T794) | |
| ALUMINUM OXIDE (UNII: LMI26O6933) | |
| TRISILOXANE (UNII: 9G1ZW13R0G) | |
| ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0) | |
| STEARIC ACID (UNII: 4ELV7Z65AP) | |
| TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E) | |
| METHYLPARABEN (UNII: A2I8C7HI9T) | |
| EDETATE DISODIUM (UNII: 7FLD91C86K) | |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4) | |
| PROPYLPARABEN (UNII: Z8IX2SC1OH) | |
| BELLIS PERENNIS FLOWER (UNII: 26I94X9A1K) | |
| MALTODEXTRIN (UNII: 7CVR7L4A2D) | |
| ETHYLPARABEN (UNII: 14255EXE39) | |
| IODOPROPYNYL BUTYLCARBAMATE (UNII: 603P14DHEB) | |
| JASMINUM OFFICINALE FLOWER (UNII: 0Q8K841432) | |
| GERANIUM ROBERTIANUM (UNII: R5I1HKOUBL) | |
| EVENING PRIMROSE OIL (UNII: 3Q9L08K71N) | |

GLICYRRHIZA GLABRA (UNII: 2788Z9758H)
.ALPHA.-TOCOPHEROL ACETATE (UNII: 9E8X80D2L0)
.BETA.-SITOSTEROL (UNII: S347WMO6M4)
TETRAHEXYLDECYL ASCORBATE (UNII: 9LBV3F07AZ)
FERRIC OXIDE RED (UNII: 1K09F3G675)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
FERROSO FERRIC OXIDE (UNII: XM0M87F357)

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:50390-002-30 | 1 in 1 CARTON | 05/01/2015 | |
| 1 | | 30 mL in 1 TUBE; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M020 | 05/01/2015 | |

Labeler - Amway Corp (083416854)

Establishment

| Name | Address | ID/FEI | Business Operations |
|---------------------------|---------|-----------|------------------------|
| Access Business Group LLC | | 839830713 | manufacture(50390-002) |

Revised: 10/2023

Amway Corp