

GPS TOPICAL ANESTHETIC- benzocaine gel
Group Practice Solutions, Inc.

Active Ingredient

Benzocaine 200mg (in each g)

Purpose

Oral Anesthetic

Use

For oral mucosa use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.

Warnings

Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy Alert: Do not use on patients with a history of allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

Do not use

- for more than 7 days unless directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly.
- for teething
- in children under 2 years of age

When using this product Avoid contact with eyes. If contact occurs, flush with water.

Do not exceed recommended dosage. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

If pregnant or breast feeding, ask a physician before use.

Keep out of reach of children.

Directions

- Apply only amount needed to the oral mucosa to prevent or relieve pain.
- children under 2 years of age: do not use

Other Information

Store at 59° - 86°F (15°-30°C). Protect from freezing.

Inactive Ingredients

flavoring, PEG 3350, PEG 400, sodium saccharin. May contain blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, (tartrazine), yellow #6, as a color additive.

Questions or comments?

631-532-2720

PEEL BACK to review drug facts

Manufactured for:



#200 26 Railroad Avenue
Babylon, NY 11702 USA
Tel: 631-532-2720

Quality Dental Products at the Right Price!



TOPICAL
ANESTHETIC GEL
(BENZOCAINE 20%)

Net Content: 1 oz (30 g)

Directions • Apply only amount needed to the oral mucosa to prevent or relieve pain. • children under 2 years of age: do not use	Other information Store at 59° - 86° F (15° - 30° C). Protect from freezing.	Inactive ingredients flavoring, PEG 3350, PEG 400, sodium saccharin, water. May contain blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, (tartrazine), yellow #6, as a color additive.	Questions or comments? 631-532-2720
---	--	---	---

98291751 Rev 11/2018

Drug Facts	Active ingredients Benzocaine 200 mg (in each g)	Purpose Oral Anesthetic	Use For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.	Warnings Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops: • pale, gray or blue colored skin (cyanosis) • headache • rapid heart rate • shortness of breath • dizziness or lightheadedness • fatigue or lack of energy Allergy Alert: Do not use on patients with a history of allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.	Do not use • for more than 7 days unless
-------------------	--	---------------------------------------	---	--	---

GPS TOPICAL ANESTHETIC

benzocaine gel

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69004-004	
Route of Administration	DENTAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)		BENZOCAINE	200 mg in 1 g	
Inactive Ingredients				
Ingredient Name			Strength	
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)				
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)				
SACCHARIN SODIUM (UNII: SB8ZUX40TY)				
FD&C YELLOW NO. 5 (UNII: I753WB2F1M)				
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)				
FD&C RED NO. 3 (UNII: PN2ZH5LOQY)				
FD&C RED NO. 40 (UNII: WZB9127XOA)				
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)				
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)				
D&C RED NO. 28 (UNII: 767IP0Y5NH)				
D&C GREEN NO. 5 (UNII: 8J6RDU8L9X)				
Product Characteristics				
Color	purple	Score		
Shape		Size		
Flavor	GRAPE	Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69004-004-30	30 g in 1 JAR; Type 0: Not a Combination Product	11/01/2018	
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC Monograph Drug	M022		07/01/2013	

Labeler - Group Practice Solutions, Inc. (968188644)