

**NIGHTTIME COLD AND FLU- acetaminophen, dextromethorphan hbr,  
doxylamine succinate liquid  
Walgreens**

-----  
**Drug Facts**

**Active ingredients (in each 30 mL)**

**Acetaminophen 650 mg**

Dextromethorphan HBr 30 mg

Doxylamine succinate 12.5 mg

**Purposes**

**Pain reliever/fever reducer**

Cough suppressant

Antihistamine

**Uses**

- temporarily relieves these common cold/flu symptoms:
  - minor aches and pains
  - headache
  - sore throat
  - fever
  - runny nose and sneezing
  - cough due to minor throat and bronchial irritation

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is

accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

### **Ask a doctor before use if you have**

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to an enlarged prostate gland

### **Ask a doctor or pharmacist before use if you are taking**

- sedatives or tranquilizers
- the blood thinning drug warfarin

### **When using this product**

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives and tranquilizers may increase drowsiness

### **Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## Directions

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in any 24-hours period
- measure only with dosing cup provided. Do not use any other dosing device.
- mL= milliliter
- keep dosing cup with product
- adults and children 12 years and over: 30 mL every 6 hours
- children under 12 years of age: do not use
- **When using other Day Time or Night Time products, carefully read each label to ensure correct dosing**

## Other information

- **each 30 mL contains:** potassium 5 mg
- **each 30 mL contains:** Sodium 19 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

## Inactive ingredients

acesulfame potassium, alcohol, citric acid, D&C yellow #10, FD&C green #3, FD&C yellow #6,, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

## Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

## Principal Display Panel

Compare to the active ingredients in Vicks® NyQuil® Cold & Flu††

### NIGHTTIME

#### Cold & Flu

ACETAMINOPHEN / 650 mg PAIN RELIEVER / FEVER REDUCER

DEXTROMETHORPHAN HBr 30 mg / COUGH SUPPRESSANT

DOXYLAMINE SUCCINATE 12.5 mg / ANTIHISTAMINE

Multi-Symptom

- Relieves headache, fever, sore throat, minor aches & pains, sneezing, runny nose & cough
- For ages 12 years & over
- ALCOHOL 10%

FL OZ (mL)

††This product is not manufactured or distributed by The Procter & Gamble Company.

Vicks® and NyQuil® are registered trademarks of The Procter & Gamble Company.

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.**

DISTRIBUTED BY: WALGREEN CO.

200 WILMOT RD., DEERFIELD, IL 60015

**Product Label**

**Drug Facts**

Active ingredients (in each 30 mL)	Purposes
Dextromethorphan HBr 30 mg.....	Cough suppressant
Doxylamine succinate 12.5 mg.....	Antihistamine

**Uses** ■ temporarily relieves these common cold/flu symptoms  
 ■ minor aches and pains ■ headache ■ sore throat  
 ■ runny nose and sneezing ■ fever  
 ■ cough due to minor throat and bronchial irritation

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:  
 ■ more than 4,000 mg of acetaminophen in 24 hours  
 ■ with other drugs containing acetaminophen  
 ■ 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash  
 If a skin reaction occurs, stop use and seek medical help right away.  
**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**  
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks

◀ PEEL HERE **Drug Facts** (continued under label)

NDC 0363-9976-08



Compare to the active ingredients in Vicks® NyQuil® Cold & Flu™

# NIGHTTIME Cold & Flu

**ACETAMINOPHEN 650 mg / PAIN RELIEVER / FEVER REDUCER**  
**DEXTROMETHORPHAN HBr 30 mg / COUGH SUPPRESSANT**  
**DOXYLAMINE SUCCINATE 12.5 mg / ANTIHISTAMINE**

**Multi-Symptom**

- Relieves headache, fever, sore throat, minor aches & pains, sneezing, runny nose & cough
- For ages 12 years & over
- ALCOHOL 10%

8 FL OZ (237 mL)

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING**

**PARENTS:**  
 Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.  
 ††This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and NyQuil® are registered trademarks of The Procter & Gamble Company.

DISTRIBUTED BY:  
**WALGREEN CO.**  
 200 WILMOT RD.,  
 DEERFIELD, IL 60015  
**100% SATISFACTION GUARANTEED**  
[walgreens.com](http://walgreens.com)  
 ©2022 Walgreen Co. W3ORG1022-F  
 PLD-B407B LB009017 REV-0323



ITEM 808541 W11042-0423-F



3 11917 12394 3

<p><b>Drug Facts (continued)</b></p> <p>after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p><b>Ask a doctor before use if you have</b>          ■ liver disease ■ glaucoma          ■ cough that occurs with too much phlegm (mucus)          ■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema          ■ trouble urinating due to enlarged prostate gland</p> <p><b>Ask a doctor or pharmacist before use if you are</b>          ■ taking sedatives or tranquilizers          ■ taking the blood thinning drug warfarin</p> <p><b>When using this product</b>          ■ excitability may occur, especially in children          ■ marked drowsiness may occur          ■ avoid alcoholic drinks          ■ be careful when driving a motor vehicle or operating machinery          ■ alcohol, sedatives, and tranquilizers may increase drowsiness</p> <p><b>Stop use and ask a doctor if</b>          ■ pain or cough gets worse or lasts more than 7 days          ■ fever gets worse or lasts more than 3 days          ■ redness or swelling is present          ■ new symptoms occur          ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.</p> <p><b>If pregnant or breast-feeding, ask a health professional before use.</b></p>	<p><b>Drug Facts (continued)</b></p> <p><b>Keep out of reach of children.</b>  <b>Overdose warning:</b> Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p><b>Directions</b>          ■ do not take more than directed (see Overdose warning)          ■ do not take more than 4 doses in any 24-hour period          ■ measure only with dosing cup provided. Do not use any other dosing device. ■ mL = milliliter          ■ keep dosing cup with product          ■ adults and children 12 years and over: 30 mL every 6 hours          ■ children under 12 years of age: do not use          ■ when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing</p> <p><b>Other information</b>          ■ each 30 mL contains: potassium 5 mg          ■ each 30 mL contains: sodium 19 mg          ■ store between 20-25°C (68-77°F). Do not refrigerate.</p> <p><b>Inactive ingredients</b> acetylsalicylic acid, alcohol, citric acid, D&amp;C yellow #10, FD&amp;C green #3, FD&amp;C yellow #6, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate</p> <p><b>Questions or comments?</b>          Call 1-877-753-3535 Monday-Friday 9AM-5PM EST</p>
--	---

**WALGREENS Nighttime Cold & Flu**

# NIGHTTIME COLD AND FLU

acetaminophen, dextromethorphan hbr, doxylamine succinate liquid

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0363-9976
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>ACESULFAME POTASSIUM</b> (UNII: 23OV73Q5G9)	
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>HIGH FRUCTOSE CORN SYRUP</b> (UNII: XY6UN3QB6S)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>TRISODIUM CITRATE DIHYDRATE</b> (UNII: B22547B95K)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C GREEN NO. 3</b> (UNII: 3P3ONR6O1S)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-9976-08	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	09/30/2022	01/30/2027

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/30/2022	01/30/2027

**Labeler** - Walgreens (008965063)

