

**NIGHTTIME COLD AND FLU- acetaminophen, dextromethorphan hbr,
doxylamine succinate liquid
Walgreens**

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 30 mg

Doxylamine succinate 12.5 mg

Purposes

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

- temporarily relieves these common cold/flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - fever
 - runny nose and sneezing
 - cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is

accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking

- sedatives or tranquilizers
- the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in any 24-hours period
- measure only with dosing cup provided. Do not use any other dosing device.
- mL= milliliter
- keep dosing cup with product
- adults and children 12 years and over: 30 mL every 6 hours
- children under 12 years of age: do not use
- **When using other Day Time or Night Time products, carefully read each label to ensure correct dosing**

Other information

- **each 30 mL contains:** potassium 5 mg
- **each 30 mL contains:** Sodium 19 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

acesulfame potassium, alcohol, citric acid, D&C yellow #10, FD&C green #3, FD&C yellow #6,, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to the active ingredients in Vicks® NyQuil® Cold & Flu††

NIGHTTIME

Cold & Flu

ACETAMINOPHEN / 650 mg PAIN RELIEVER / FEVER REDUCER

DEXTROMETHORPHAN HBr 30 mg / COUGH SUPPRESSANT

DOXYLAMINE SUCCINATE 12.5 mg / ANTIHISTAMINE

Multi-Symptom

- Relieves headache, fever, sore throat, minor aches & pains, sneezing, runny nose & cough
- For ages 12 years & over
- ALCOHOL 10%

FL OZ (mL)

††This product is not manufactured or distributed by The Procter & Gamble Company.

Vicks® and NyQuil® are registered trademarks of The Procter & Gamble Company.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY: WALGREEN CO.

200 WILMOT RD., DEERFIELD, IL 60015

Product Label

Drug Facts

Active ingredients (in each 30 mL)	Purposes
Dextromethorphan HBr 30 mg.....	Cough suppressant
Doxylamine succinate 12.5 mg.....	Antihistamine

Uses ■ temporarily relieves these common cold/flu symptoms
 ■ minor aches and pains ■ headache ■ sore throat
 ■ runny nose and sneezing ■ fever
 ■ cough due to minor throat and bronchial irritation

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
 ■ more than 4,000 mg of acetaminophen in 24 hours
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
 If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks

◀ PEEL HERE **Drug Facts** (continued under label)

NDC 0363-9976-08



Compare to the active ingredients in Vicks® NyQuil® Cold & Flu™



NIGHTTIME Cold & Flu

ACETAMINOPHEN 650 mg / PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr 30 mg / COUGH SUPPRESSANT
DOXYLAMINE SUCCINATE 12.5 mg / ANTIHISTAMINE

Multi-Symptom

- Relieves headache, fever, sore throat, minor aches & pains, sneezing, runny nose & cough
- For ages 12 years & over
- **ALCOHOL 10%**

8 FL OZ (237 mL)

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING

PARENTS:
 Learn about teen medicine abuse
www.StopMedicineAbuse.org

†Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.
 ††This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and NyQuil® are registered trademarks of The Procter & Gamble Company.

DISTRIBUTED BY:
WALGREEN CO.
 200 WILMOT RD.,
 DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com
 ©2022 Walgreen Co. W3ORG1022-F
 PLD-B407B LB009017 REV-0323



Discard Seals, Empty & Replace Cap

PLASTIC BOTTLE PLASTIC CUP

ITEM 808541 W11042-0423-F



3 11917 12394 3

<p>Drug Facts (continued)</p> <p>after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> ■ liver disease ■ glaucoma ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ trouble urinating due to enlarged prostate gland <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> ■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin <p>When using this product</p> <ul style="list-style-type: none"> ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives, and tranquilizers may increase drowsiness <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ pain or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. <p>If pregnant or breast-feeding, ask a health professional before use.</p>	<p>Drug Facts (continued)</p> <p>Keep out of reach of children.</p> <p>Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions</p> <ul style="list-style-type: none"> ■ do not take more than directed (see Overdose warning) ■ do not take more than 4 doses in any 24-hour period ■ measure only with dosing cup provided. Do not use any other dosing device. ■ mL = milliliter ■ keep dosing cup with product ■ adults and children 12 years and over: 30 mL every 6 hours ■ children under 12 years of age: do not use ■ when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing <p>Other information</p> <ul style="list-style-type: none"> ■ each 30 mL contains: potassium 5 mg ■ each 30 mL contains: sodium 19 mg ■ store between 20-25°C (68-77°F). Do not refrigerate. <p>Inactive ingredients acetylsalicylic acid, alcohol, citric acid, D&C yellow #10, FD&C green #3, FD&C yellow #6, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate</p> <p>Questions or comments? Call 1-877-753-3535 Monday-Friday 9AM-5PM EST</p>
--	---

WALGREENS Nighttime Cold & Flu

NIGHTTIME COLD AND FLU

acetaminophen, dextromethorphan hbr, doxylamine succinate liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-9976
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ALCOHOL (UNII: 3K9958V90M)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-9976-08	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	09/30/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/30/2022	

Labeler - Walgreens (008965063)

