

HEAD CONGESTION FLU SEVERE PE- acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, coated
CHAIN DRUG MARKETING ASSOCIATION INC

1174-QCH-2022-0803

Drug Facts

<i>Active ingredients (in each caplet)</i>	<i>Purposes</i>
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 100 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves these symptoms due to the common cold:
 - nasal congestion
 - headache
 - minor aches and pains
 - cough
 - sore throat
 - sinus congestion and pressure
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make coughs more productive
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none">▪ take 2 caplets every 4 hours▪ do not take more than 10 caplets in 24 hours
children under 12 years	<ul style="list-style-type: none">▪ ask a doctor

Other information

- store between 20-25°C (68-77°F) in a dry place
- retain carton for complete product information and warnings

Inactive ingredients

colloidal silicon dioxide, corn starch, croscarmellose sodium, crospovidone, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide

PRINCIPAL DISPLAY PANEL

QUALITY CHOICE®

NDC 63868-102-24

†Compare to Active Ingredients in SUDAFED PE® Head Congestion + Flu Severe
Head Congestion + Flu Severe

Acetaminophen, Dextromethorphan HBr, Guaifenesin, Phenylephrine HCl

Pain Reliever | Fever Reducer, Cough Suppressant, Expectorant, Nasal Decongestant

For Relief of:

Sinus Pressure

Headache

Sore Throat

Cough

Chest Congestion

Actual Size

24 CAPLETS



FT174020CH_R1

Drug Facts (continued)
Other information
■ store between 20-25°C (68-77°F) in a dry place
■ retain carton for complete product information and warnings
Inactive ingredients
corn starch, croscarmellose sodium, crospovidone, colloidal silicon dioxide

Drug Facts (continued)
FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide

Drug Facts (continued)
Warnings
Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours.
■ with other drugs containing acetaminophen
■ or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ you are now taking a prescription antidepressant or emotional conditions, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
■ If you have ever had an allergic reaction to this product or any of its ingredients
■ If you have ever had an allergic reaction to this product.
Ask a doctor before use if you have
■ high blood pressure
■ liver disease
■ heart disease
■ diabetes
■ thyroid disease

Drug Facts (continued)
Directions
adults and children 12 years and over ■ take 2 caplets every 4 hours ■ do not take more than 10 caplets in 24 hours
children under 12 years ■ ask a doctor
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Keep out of reach of children.
If pregnant or breast-feeding, ask a health professional before use. These could be signs of a serious condition.
■ cough comes back or occurs with rash or headache that lasts
■ redness or swelling is present
■ new symptoms occur
■ fever gets worse or lasts more than 3 days
■ pain, nasal congestion, or cough gets worse or lasts more than 7 days
■ nervousness, dizziness, or sleeplessness occur
Stop use and ask a doctor if
When using this product do not exceed recommended dosage
thinning drug warfarin
Ask a doctor or pharmacist before use if you are taking the blood
■ trouble urinating due to an enlarged prostate gland
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
■ cough that occurs with too much phlegm (mucus)
Drug Facts (continued)

Drug Facts
Active ingredients (in each caplet) Purpose
Acetaminophen 325 mg: Pain reliever/fever reducer
Dextromethorphan HBr 10 mg: Cough suppressant
Guaifenesin 100 mg: Expectorant
Phenylephrine HCl 5 mg: Nasal decongestant

Drug Facts (continued)
Uses
temporarily relieves these symptoms due to the common cold: ■ nasal congestion ■ headache ■ minor aches and pains ■ cough ■ sore throat
■ temporarily reduces fever
■ temporarily reduces fever

†This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Sudafed PE® Head Congestion + Flu Severe.

DO NOT USE IF BLISTER UNITS ARE TORN OR BROKEN

100% SATISFACTION GUARANTEED
Distributed by CDMA, Inc.
Novi, MI 48375
www.qualitychoice.com
Questions: 800-935-2362

NDC 63868-102-24

***Compare to Active Ingredients in SUDAFED PE® Head Congestion + Flu Severe**



Head Congestion + Flu Severe PE

Acetaminophen, Dextromethorphan HBr, Guaifenesin, Phenylephrine HCl

Pain Reliever | Fever Reducer, Cough Suppressant, Expectorant, Nasal Decongestant

For Relief of:

- Sinus Pressure
- Headache
- Sore Throat
- Cough
- Chest Congestion

Actual Size



24 CAPLETS

HEAD CONGESTION FLU SEVERE PE

acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63868-102
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
CROSPVIDONE (UNII: 2S7830E561)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	orange	Score	no score
Shape	OVAL	Size	17mm
Flavor		Imprint Code	AAA;1134
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
---	-----------	---------------------	----------------------	--------------------

1	NDC:63868-102-24	2 in 1 CARTON	08/03/2022	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M012	08/03/2022	

Labeler - CHAIN DRUG MARKETING ASSOCIATION INC (011920774)