

**ACETAMINOPHEN, ASPIRIN, AND CAFFEINE- acetaminophen, aspirin, and caffeine tablet**  
**Bi-Mart**

-----

***Drug Facts***

***Active ingredients (in each geltaab)***

Acetaminophen 250 mg

Aspirin 250 mg (NSAID)\*\*

Caffeine 65 mg

\*\*nonsteroidal anti-inflammatory drug

***Purposes***

Pain reliever

Pain reliever

Pain reliever aid

***Uses***

Temporarily relieves minor aches and pains due to:

- headache
- a cold
- arthritis
- muscular aches
- toothache
- premenstrual & menstrual cramps

***Warnings***

**Reye's syndrome:**Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:**Acetaminophen may cause skin reactions. Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

**Allergy alert:**Aspirin may cause a severe allergic reaction which may include: • hives •

facial swelling • asthma (wheezing) • shock

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 caplets in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDS (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasional rapid heartbeat.

**Do not use** • if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer

- with any other drug containing acetaminophen (prescription or nonprescription). if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if** • you have liver disease • stomach bleeding warning applies to you • you have a history of stomach problems such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you are taking a diuretic • you have asthma

**Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug, or are under a doctor's care for any serious condition

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better
- ringing in the ears or loss of hearing occurs
- painful area is red or swollen
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- any new symptoms occur

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center **(1-800-222-1222)** right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

### ***Directions***

- **do not use more than directed**
- drink a full glass of water with each dose
- **adults and children 12 years and over:**take 2 caplets every 6 hours; not more than 8 caplets in 24 hours
- **children under 12 years:**ask a doctor

### ***Other information***

- store at 20°-25°C(68°-77°F)
- close cap tightly after use
- DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED

### ***Inactive ingredients***

colloidal silicon dioxide, croscarmellose sodium, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, PVPK-30, sodium starch glycolate, stearic acid, talc, titanium dioxide.

### ***Questions?***

**call toll-free 1-844-912-4012**

**PRINCIPAL DISPLAY PANEL-100 Count**

**NDC 37835-531-01**

Compare to the active ingredients in Excedrin® Extra Strength\*

**Extra Strength**

**Acetaminophen, Aspirin (NSAID)\*\*, and Caffeine**

Pain Reliever/ Pain Reliever Aid

100 Caplets

\*This product is not manufactured or distributed by the owner of the registered trademark Excedrin® Extra Strength.

Distributed by:

**BI-MART**

Eugene, OR 97402

0 71357 00184 2

**BI-MART** NDC 37835-531-01

Compare to the active ingredients in EXCEDRIN® EXTRA STRENGTH\*

**EXTRA STRENGTH**  
**Acetaminophen,**  
**Aspirin (NSAID)\*\***  
**and Caffeine**  
**Pain Reliever / Pain Reliever Aid**



**100 CAPLETS**

DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED

**Drug Facts**

**Active ingredients (in each caplet)** Purpose  
Acetaminophen 250 mg ..... Pain reliever  
Aspirin 250 mg (NSAID)\*\* ..... Pain reliever  
Caffeine 65 mg ..... Pain reliever aid  
\*\*nonsteroidal anti-inflammatory drug

**Uses** ■ temporarily relieves minor aches and pains due to:  
■ headache ■ a cold ■ arthritis ■ muscular aches ■ toothache  
■ premenstrual & menstrual cramps

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.  
**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 8 caplets in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or

PEEL FOR DIRECTIONS 531-01-BM 02/25

LIFT HERE

**Drug Facts (continued)**

nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

**Do not use** ■ if you have ever had an allergic reaction to acetaminophen, aspirin or other pain reliever/fever reducer ■ with any other product containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if** ■ you have liver disease ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma

**Ask a doctor or pharmacist before use if you are** ■ taking a prescription drug for diabetes, gout, or arthritis ■ taking any other drug, or are under a doctor's care for any serious condition

**Stop use and ask a doctor if** ■ an allergic reaction occurs. Seek medical help right away, ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ ringing in the ears or loss of hearing occurs ■ painful area is red or swollen ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ any new symptoms occur. These could be signs of a serious condition

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

**Directions** ■ do not use more than directed

■ drink a full glass of water with each dose  
■ adults and children 12 years and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours  
■ children under 12 years: ask a doctor

**Other information** ■ store at 20°-25°C (68°-77°F)

■ close cap tightly after use

**Inactive ingredients** Colloidal silicon dioxide, croscarmellose sodium, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, PVP-K30, sodium starch glycolate, stearic acid, talc, titanium dioxide.

**Questions?** call toll-free 1-844-912-4012

\*This product is not manufactured or distributed by the owner of the registered trademark Excedrin® Extra Strength

Distributed by:  
**BI-MART**  
Eugene, OR 97402

BI-MART UNCONDITIONAL MONEY BACK GUARANTEE  
If you are not satisfied with the quality of this product, return the unused portion and your entire purchase price will be cheerfully refunded.

**PRINCIPAL DISPLAY PANEL-250 Count**

**NDC 37835-531-25**

Compare to the active ingredients in Excedrin® Extra Strength\*

**Extra Strength**

**Acetaminophen, Aspirin (NSAID)\*\* , and Caffeine**

Pain Reliever/ Pain Reliever Aid

250 Caplets

\*This product is not manufactured or distributed by the owner of the registered trademark Excedrin® Extra Strength.

Distributed by:

**BI-MART**

Eugene, OR 97402

5 713571006751 0

**VP VALUE PACK**

**BI-MART** NDC 37835-531-25

Compare to the active ingredients in EXCEDRIN® EXTRA STRENGTH\*

**EXTRA STRENGTH**  
**Acetaminophen,**  
**Aspirin (NSAID)\*\***  
**and Caffeine**  
**Pain Reliever / Pain Reliever Aid**



**250 CAPLETS**

DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED

**Drug Facts**

Active Ingredients (in each caplet)	Purpose
Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID)**	Pain reliever
Caffeine 65 mg	Pain reliever aid

**Uses** ■ temporarily relieves minor aches and pains due to:  
 ■ headache ■ a cold ■ arthritis ■ muscular aches ■ toothache  
 ■ premenstrual & menstrual cramps

**Warnings**

**Fever's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 8 caplets in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinner (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

PEEL FOR DIRECTIONS

531-25-BM 02/25

LIFT HERE

**Drug Facts (continued)**

**Do not use** ■ if you have ever had an allergic reaction to acetaminophen, aspirin or other pain reliever/fever reducer ■ with any other product containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
**Ask a doctor before use if** ■ you have liver disease ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma  
**Ask a doctor or pharmacist before use if you are** ■ taking a prescription drug for diabetes, gout, or arthritis ■ taking any other drug, or are under a doctor's care for any serious condition

**Stop use and ask a doctor if** ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ ringing in the ears or loss of hearing occurs ■ painful area is red or swollen ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ any new symptoms occur. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

**Directions** ■ do not use more than directed  
 ■ drink a full glass of water with each dose  
 ■ adults and children 12 years and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours  
 ■ children under 12 years: ask a doctor

**Other information** ■ store at 20°-25°C (68°-77°F)  
 ■ close cap tightly after use

**Inactive ingredients** Colloidal silicon dioxide, croscarmellose sodium, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, PVP-K30, sodium starch glycolate, stearic acid, talc, titanium dioxide.

**Questions?** call toll-free 1-844-912-4012

\*This product is not manufactured or distributed by the owner of the registered trademark Excedrin® Extra Strength

Distributed by:  
**BI-MART**  
 Eugene, OR 97402

BI-MART UNCONDITIONAL MONEY BACK GUARANTEE  
 If you are not completely satisfied with your BI-MART QUALITY PRODUCT, regardless of the reason, return the unused portion and your entire purchase price will be cheerfully refunded.

**ACETAMINOPHEN, ASPIRIN, AND CAFFEINE**

acetaminophen, aspirin, and caffeine tablet

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:37835-531
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
<b>CAFFEINE</b> (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>SODIUM STARCH GLYCOLATE TYPE A</b> (UNII: H8AV0SQX4D)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	HH6
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37835-531-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	02/01/2025	
2	NDC:37835-531-25	250 in 1 BOTTLE; Type 0: Not a Combination Product	02/01/2025	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/01/2025	

**Labeler -** Bi-Mart (027630078)

## Establishment

Name	Address	ID/FEI	Business Operations
Health Pharma USA LLC		080804485	manufacture(37835-531)

Revised: 2/2025

Bi-Mart