

ACETAMINOPHEN, ASPIRIN, AND CAFFEINE- acetaminophen, aspirin, and caffeine tablet

Bi-Mart

Drug Facts

Active ingredients (in each gelta)

Acetaminophen 250 mg

Aspirin 250 mg (NSAID)**

Caffeine 65 mg

**nonsteroidal anti-inflammatory drug

Purposes

Pain reliever

Pain reliever

Pain reliever aid

Uses

Temporarily relieves minor aches and pains due to:

- headache
- a cold
- arthritis
- muscular aches
- toothache
- premenstrual & menstrual cramps

Warnings

Reye's syndrome:Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert:Acetaminophen may cause skin reactions. Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert:Aspirin may cause a severe allergic reaction which may include: • hives •

facial swelling • asthma (wheezing) • shock

Liver warning:This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 caplets in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning:This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDS (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Caffeine warning:The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasional rapid heartbeat.

Do not use • if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer

- with any other drug containing acetaminophen (prescription or nonprescription). if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if • you have liver disease • stomach bleeding warning applies to you • you have a history of stomach problems such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you are taking a diuretic • you have asthma

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug, or are under a doctor's care for any serious condition

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better
- ringing in the ears or loss of hearing occurs
- painful area is red or swollen
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- any new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center **(1-800-222-1222)** right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed**
- drink a full glass of water with each dose
- **adults and children 12 years and over:**take 2 caplets every 6 hours; not more than 8 caplets in 24 hours
- **children under 12 years:**ask a doctor

Other information

- store at 20°-25°C(68°-77°F)
- close cap tightly after use
- DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED

Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, PVPK-30, sodium starch glycolate, stearic acid, talc, titanium dioxide.

Questions?

call toll-free 1-844-912-4012

PRINCIPAL DISPLAY PANEL-100 Count

NDC 37835-531-01

Compare to the active ingredients in Excedrin® Extra Strength*

Extra Strength

Acetaminophen, Aspirin (NSAID), and Caffeine**

Pain Reliever/ Pain Reliever Aid

100 Caplets

*This product is not manufactured or distributed by the owner of the registered trademark Excedrin® Extra Strength.

Distributed by:

BI-MART

Eugene, OR 97402

0 71357 00184 2

BI-MART NDC 37835-531-01

Compare to the active ingredients in EXCEDRIN® EXTRA STRENGTH*

EXTRA STRENGTH
Acetaminophen,
Aspirin (NSAID)**
and Caffeine
Pain Reliever / Pain Reliever Aid



100 CAPLETS

DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED

Drug Facts

Active ingredients (in each caplet) Purpose
 Acetaminophen 250 mg Pain reliever
 Aspirin 250 mg (NSAID)** Pain reliever
 Caffeine 65 mg Pain reliever aid
 **nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to:
 ■ headache ■ a cold ■ arthritis ■ muscular aches ■ toothache
 ■ premenstrual & menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 8 caplets in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or

PEEL FOR DIRECTIONS 531-01-BM 02/25

LIFT HERE

Drug Facts (continued)

nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use ■ if you have ever had an allergic reaction to acetaminophen, aspirin or other pain reliever/fever reducer ■ with any other product containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if ■ you have liver disease ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma

Ask a doctor or pharmacist before use if you are ■ taking a prescription drug for diabetes, gout, or arthritis ■ taking any other drug, or are under a doctor's care for any serious condition

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away, ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ ringing in the ears or loss of hearing occurs ■ painful area is red or swollen ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ any new symptoms occur. These could be signs of a serious condition

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions ■ do not use more than directed

■ drink a full glass of water with each dose
 ■ adults and children 12 years and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours
 ■ children under 12 years: ask a doctor

Other information ■ store at 20°-25°C (68°-77°F)
 ■ close cap tightly after use

Inactive ingredients Colloidal silicon dioxide, croscarmellose sodium, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, PVP-K30, sodium starch glycolate, stearic acid, talc, titanium dioxide.

Questions? call toll-free 1-844-912-4012

*This product is not manufactured or distributed by the owner of the registered trademark Excedrin® Extra Strength

Distributed by:
BI-MART
 Eugene, OR 97402

BI-MART UNCONDITIONAL MONEY BACK GUARANTEE
 If you are not satisfied with your purchase for any reason, return the unused portion and your entire purchase price will be cheerfully refunded.

PRINCIPAL DISPLAY PANEL-250 Count

NDC 37835-531-25

Compare to the active ingredients in Excedrin® Extra Strength*

Extra Strength

Acetaminophen, Aspirin (NSAID), and Caffeine**

Pain Reliever/ Pain Reliever Aid

250 Caplets

*This product is not manufactured or distributed by the owner of the registered trademark Excedrin® Extra Strength.

Distributed by:

BI-MART

Eugene, OR 97402

5 713571006751

VP VALUE PACK

BI-MART

NDC 37835-531-25

Compare to the active ingredients in EXCEDRIN® EXTRA STRENGTH*

EXTRA STRENGTH

Acetaminophen, Aspirin (NSAID) and Caffeine**

Pain Reliever / Pain Reliever Aid
250 CAPLETS



DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED

Drug Facts

Active Ingredients (in each caplet)	Purpose
Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID)**	Pain reliever
Caffeine 65 mg	Pain reliever aid

Uses ■ temporarily relieves minor aches and pains due to:
 ■ headache ■ a cold ■ arthritis ■ muscular aches ■ toothache
 ■ premenstrual & menstrual cramps

Warnings

Fever's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 8 caplets in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

PEEL FOR DIRECTIONS

531-25-BM 02/25

LIFT HERE

Drug Facts (continued)

Do not use ■ if you have ever had an allergic reaction to acetaminophen, aspirin or other pain reliever/fever reducer ■ with any other product containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
Ask a doctor before use if ■ you have liver disease ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma
Ask a doctor or pharmacist before use if you are ■ taking a prescription drug for diabetes, gout, or arthritis ■ taking any other drug, or are under a doctor's care for any serious condition

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ ringing in the ears or loss of hearing occurs ■ painful area is red or swollen ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ any new symptoms occur. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions ■ do not use more than directed
 ■ drink a full glass of water with each dose
 ■ adults and children 12 years and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours
 ■ children under 12 years: ask a doctor

Other information ■ store at 20°-25°C (68°-77°F)
 ■ close cap tightly after use

Inactive ingredients Colloidal silicon dioxide, croscarmellose sodium, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, PVP-K30, sodium starch glycolate, stearic acid, talc, titanium dioxide.

Questions? call toll-free 1-844-912-4012

*This product is not manufactured or distributed by the owner of the registered trademark Excedrin® Extra Strength

Distributed by:
BI-MART
 Eugene, OR 97402

BI-MART UNCONDITIONAL MONEY BACK GUARANTEE
 If you are not completely satisfied with your BI-MART QUALITY PRODUCT, regardless of the reason, return the unused portion and your entire purchase price will be cheerfully refunded.

ACETAMINOPHEN, ASPIRIN, AND CAFFEINE

acetaminophen, aspirin, and caffeine tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37835-531
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
STARCH, CORN (UNII: O8232NY3SJ)	
POVIDONE K30 (UNII: U725QWY32X)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	white	Score	no score
Shape	OVAL	Size	17mm
Flavor		Imprint Code	HH6
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37835-531-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	02/01/2025	
2	NDC:37835-531-25	250 in 1 BOTTLE; Type 0: Not a Combination Product	02/01/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/01/2025	

Labeler - Bi-Mart (027630078)

Establishment

Name	Address	ID/FEI	Business Operations
ELYSIUM PHARMACEUTICALS LIMITED		915664486	manufacture(37835-531)

Revised: 7/2025

Bi-Mart