

PERIOD PILLS BY BIIKAY- acetaminophen, ibuprofen tablet

Kai Squared

Active Ingredients

Active Ingredients (in each caplet) Purpose

Acetaminophen 250 mg Pain reliever

Ibuprofen 125 mg (NSAID**) Pain reliever

**nonsteroidal anti-inflammatory drug

Directions:

- Do not take more than directed
- Adults and children 12 years and over
- Take 2 caplets every 8 hours while symptoms persist
- Children under 12 years ask a doctor
- Do not take more than 6 caplets in 24 hours, unless directed by a doctor

Warnings

Acetaminophen liver damage warning: This product contains acetaminophen. Severe liver damage may occur if you take: with other drugs containing acetaminophen, more than 6 caplets in 24 hours, which is the maximum daily amount for this product, 3 or more alcoholic drinks every day while using this product.

Acetaminophen allergy alert: may cause severe skin reactions. Symptoms include, skin reddening, blisters, rash.

If skin reactions occurs, stop use and seek medical help right away.

NSAID allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters

If an allergic reaction occurs, stop use and seek medical help right away.

NSAID stomach bleeding warnings: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older, have had stomach ulcers or bleeding problems, take a blood thinning (anticoagulant) or steroid drug, take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others], have 3 or more alcoholic drinks every day while using this product, take more or for a longer time than directed.

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use, with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist, if you have ever had an allergic reaction to acetaminophen or any other pain reliever, right before or after heart surgery.

Ask a doctor before use if:

You have liver disease, stomach bleeding warning applies to you, you have problems or serious side effects from taking pain relievers, you have a history of stomach problems such as heartburn, you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke, you are taking a diuretic.

Ask a doctor or pharmacist before use if you are:

Under a doctor's care for any serious condition, taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin, taking any other drug.

When using this product, take with food or milk if stomach upset occurs.

Stop use and ask a doctor if, you experience any of the following signs of stomach bleeding:

feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better, you have symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling, pain gets worse or lasts more than 10 days, redness or swelling is present in the painful area, any new symptoms appear.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Uses:

Temporarily relieves minor aches and pains due to: headache, toothache, backache, menstrual cramps, muscular aches, minor pain of arthritis

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Uses:

Temporarily relieves minor aches and pains due to:

headache, toothache, backache, menstrual cramps, muscular aches, minor pain of arthritis

Inactive ingredients:

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, crospovidone, ferric oxide red, ferric oxide yellow, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, povidone, pregelatinized starch, sodium lauryl sulfate, stearic acid and titanium dioxide.

Period Pills by Biikay

Pain Reliever



PERIOD PILLS

by Biikay

Pain Reliever
Acetaminophen 250 mg
Ibuprofen (NSAID) 125 mg
36 Caplets

DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 250 mg... Pain reliever
Ibuprofen 125 mg (NSAID)**... Pain reliever
**nonsteroidal anti-inflammatory drug

Uses

temporarily relieves minor aches and pains due to:
■ headache ■ toothache ■ backache ■ menstrual cramps
■ muscular aches ■ minor pain of arthritis

Warnings

Acetaminophen liver damage warning: This product contains acetaminophen. Severe liver damage may occur if you take:
■ with other drugs containing acetaminophen ■ more than 6 caplets in 24 hours, which is the maximum daily amount for this product ■ 3 or more alcohol drinks every day while using this product
Acetaminophen allergy alert: may cause severe skin reactions. Symptoms include ■ skin reddening ■ blisters ■ rash
If skin reactions occur, stop use and seek medical help right away.
NSAID allergy alert: Ibuprofen may cause a severe allergic reaction.

Drug Facts (continued on inside)



L82269
R0125
Distributed by: Kai Squared New York, NY 10001

PEEL
HERE

Lot:
Exp.:

UNVARNISHED

Drug Facts (continued)

especially in people allergic to aspirin. Symptoms may include:

■ hives ■ facial swelling ■ asthma (wheezing) ■ shock
■ skin reddening ■ rash ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

NSAID stomach bleeding warnings: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you have ever had an allergic reaction to acetaminophen or any other pain reliever ■ right before or after heart surgery

Ask a doctor before use if ■ you have liver disease ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers ■ you have a history of stomach problems such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug

When using this product ■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if ■ you experience any of the following signs or symptoms: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling

Drug Facts (continued)

■ pain gets worse or lasts more than 10 days ■ redness or swelling is present in the painful area ■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not take more than directed adults and children ■ take 2 caplets every 8 hours while 12 years and over ■ ask a doctor

■ do not take more than 6 caplets in 24 hours, unless directed by a doctor

Other information ■ read all warnings and directions before use. ■ store at 20 to 25°C (68 to 77°F) ■ avoid excessive heat above 40°C (104°F)

Inactive ingredients carnauba wax, colloidal silicon dioxide, croscarmellose sodium, crospovidone, ferric oxide red, ferric oxide yellow, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, povidone, pregelatinized starch, sodium lauryl sulfate, stearic acid and titanium dioxide.

Questions or comments?

Call 1-800-997-5561 Monday through Friday 9AM - 5PM EST.

PERIOD PILLS BY BIIKAY

acetaminophen, ibuprofen tablet

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:85576-101

Route of Administration

ORAL

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|-------------------|----------|
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN | 250 mg |
| IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM) | IBUPROFEN | 125 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U) | |
| CROSCARMELLOSE SODIUM (UNII: M28OL1HH48) | |
| CROSPVIDONE (UNII: 2S7830E561) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |
| FERRIC OXIDE YELLOW (UNII: EX438O2MRT) | |
| CARNAUBA WAX (UNII: R12CBM0EIZ) | |
| FERRIC OXIDE RED (UNII: 1K09F3G675) | |
| MAGNESIUM STEARATE (UNII: 70097M6I3O) | |
| POVIDONE (UNII: FZ989GH94E) | |
| SODIUM LAURYL SULFATE (UNII: 368GB5141J) | |
| POLYDEXTROSE (UNII: VH2XOU12IE) | |

Product Characteristics

| | | | |
|-----------------|--------|---------------------|----------|
| Color | yellow | Score | no score |
| Shape | OVAL | Size | 14mm |
| Flavor | | Imprint Code | |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:85576-101-36 | 36 in 1 BOTTLE; Type 0: Not a Combination Product | 10/02/2025 | |



Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| ANDA | ANDA216592 | 10/02/2025 | |

Labeler - Kai Squared (130976767)

Establishment

| Name | Address | ID/FEI | Business Operations |
|------------------|---------|-----------|------------------------|
| Allegiant Health | | 079501930 | manufacture(85576-101) |

Revised: 12/2025

Kai Squared