

**TUSSIN DM NIGHTTIME COUGH MAXIMUM STRENGTH- dextromethorphan hbr,
doxylamine succinate liquid
WinCo Foods, LLC**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredients (in each 20 mL)

Dextromethorphan HBr 30 mg

Doxylamine succinate 12.5 mg

Purposes

Cough suppressant

Antihistamine

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - runny nose
 - sneezing
 - itchy, watery eyes
 - itching of the nose or throat
- controls the impulse to cough to help you sleep

Warnings

Do not use

- to sedate a child or to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- glaucoma
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma,

chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers.

When using this product

- do not use more than directed
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur especially in children

Stop use and ask a doctor if

cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- **this adult product is not intended for use in children under 12 years of age**
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- adults and children 12 years and over: 20 mL every 6 hours
- children under 12 years: do not use

Other information

- **each 20 mL contains:** sodium 9 mL
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

anhydrous citric acid, carboxymethylcellulose sodium, FD&C blue #1, FD&C red #40, flavors glycerin, high fructose corn syrup, menthol, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sucralose, xanthan gum

Questions or comments?

Call **1-800-824-1706** Monday-Friday 9AM-5PM MST

Principal Display Panel

*Compare to the active ingredients in Robitussin® Maximum Strength Nighttime Cough DM

MAXIMUM STRENGTH

Tussin DM

Nighttime

DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT

DOXYLAMINE SUCCINATE / ANTIHISTAMINE

Relieves: Cough, Itchy Throat & Runny Nose

Adult Formula, For Ages 12 Years & Over

Alcohol Free

Dosage cup provided

FL OZ (mL)

*This product is not manufactured or distributed by PFIZER CONSUMER HEALTHCARE, distributors of RONITUSSIN® Maximum Strength Nighttime Cough DM.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

DISTRIBUTED BY:

WINCO FOODS, LLC, BOISE, ID 83704

Package Label

Drug Facts (continued)

Directions

- this adult product is not intended for use in children under 12 years of age
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- adults and children 12 years and over: 20 mL every 6 hours
- children under 12 years: do not use

Other information

- each 20 mL contains: sodium 9 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients citric acid, carboxymethylcellulose sodium, FD&C blue #1, FD&C red #40, flavor, glycerin, high fructose corn syrup, menthol, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sucralose, xanthan gum

Questions or comments?
Call 800-824-1706 Monday - Friday 9am - 4pm MST

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

DISTRIBUTED BY:
WINCO FOODS, LLC, BOISE, ID 83704

*This product is not manufactured or distributed by PFIZER CONSUMER HEALTHCARE, distributor of ROBITUSSIN® Maximum Strength Nighttime Cough DM.



The Supermarket Low Price Leader®

100% SATISFACTION GUARANTEE

Comments & Questions?
800-824-1706 - www.wincofoods.com

*Compare to the active ingredients in ROBITUSSIN® Maximum Strength Nighttime Cough DM



DEXTROMETHORPHAN HBr/
COUGH SUPPRESSANT
DOXYLAMINE SUCCINATE/
ANTIHISTAMINE

Relieves: Cough, Itchy Throat & Runny Nose

Adult Formula, For Ages 12 Years & Over

Alcohol Free



4 FL OZ (118 mL)

*Compare to the active ingredients in ROBITUSSIN® Maximum Strength Nighttime Cough DM



DEXTROMETHORPHAN HBr/
COUGH SUPPRESSANT
DOXYLAMINE SUCCINATE/
ANTIHISTAMINE

Relieves: Cough, Itchy Throat & Runny Nose

Adult Formula, For Ages 12 Years & Over

Alcohol Free



4 FL OZ (118 mL)

Drug Facts

Active ingredients (in each 20 mL)	Purposes
Dextromethorphan HBr 30 mg	Cough suppressant
Doxylamine succinate 12.5 mg	Antihistamine

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat
- controls the impulse to cough to help you sleep

Warnings

Do not use

- to sedate a child or to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- trouble urinating due to an enlarged prostate gland
- glaucoma
- a cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

When using this product

- do not use more than directed
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.



PLD-A481B
FC005213

Lot No.:
Exp. Date:

WINCO FOODS Maximum Strength Nighttime Cough

TUSSIN DM NIGHTTIME COUGH MAXIMUM STRENGTH

dextromethorphan hbr, doxylamine succinate liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:67091-333
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 20 mL
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 20 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
MENTHOL (UNII: L7T10EIP3A)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	

Product Characteristics

Color	red	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67091-333-18	1 in 1 BOX	02/01/2019	05/31/2024
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	02/01/2019	05/31/2024

Labeler - WinCo Foods, LLC (056098817)