

WOFANLI QUIT SMOKING NICOTINE PATCHES- nicotine transdermal patches patch
Shenzhen Paxi Trading Co., Ltd.

85835-004

Active Ingredient

Nicotine 21mg

Purpose

Stopsmoking aid

Use

reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking.

Apply one new patch every 24 hours on skin that is dry, clean and hairless. Remove backing from patch and immediately press onto skin. Hold for 10 seconds. Wash hands after applying or removing patch. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch. The used patch should be removed and a new one applied to a different skin site at the same time each day.

If you have vivid dreams, you may remove the patch at bedtime and apply a new one in the morning.

Do not wear more than one patch at a time.

Do not cut patch in half or into smaller pieces.

Do not leave patch on for more than 24 hours because it may irritate your skin and loses strength after 24 hours.

To avoid possible burns, remove patch before undergoing any MRI (magnetic resonance imaging) procedures.

It is important to complete treatment. If you feel you need to use the patch for a longer period to keep from smoking, talk to your healthcare provider.

Warnings

If you are pregnant or breast feeding, only use this medicine on the advice of your health care provider.

Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Do not use

Ask a doctor before use if you have

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- an allergy to adhesive tape or have skin problems, because you are more likely to get rashes.
- Stomach ulcer or diabetes.
- History of seizures.

Ask Doctor

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug.
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

When Using

When using this product

- if you have vivid dreams or other sleep disturbances remove this patch at bedtime.

Stop Use

Stop use and ask a doctor if

- skin redness caused by the patch does not go away after four days, or if your skin swells, or you get a rash.
- irregular heartbeat or palpitations occur.
- you get symptoms of nicotine overdose, such as persistent nausea, vomiting, dizziness, weakness and rapid heartbeat.
- You have symptoms of an allergic reaction (such as difficulty breathing or rash).

Keep Out Of Reach Of Children

Keep out of reach of children and pets

Used patches contain enough nicotine to poison children and pets. If swallowed, seek medical help immediately or contact a Poison Control Center 1-800-222-1222 right away. Save the pouch for patch disposal. Dispose of used patches by folding the sticky ends together and placing them in the pouch.

Directions

If you are under 18 years of age, ask a doctor before use.

Before using this product, read the enclosed self-help guide for complete directions and other information.

Begin using the patch on your quit day.

If you smoke more than 20 cigarettes per day, use the following schedule below:

Weeks 1 and 2

STEP 1+
use 1½
patch/day

Weeks 3 and 6
STEP 1
use 1
patch/day

Weeks 7 and 8
STEP 2
use 1/2
patch/day

Weeks 9 and 10
STEP 3
use 1/4
patch/day

If you smoke more than 10 but not more than 20 cigarettes, start with Step 1, if you smoke 10 or less cigarettes per day, start with Step 2 for 4 weeks (Start with 1 patch or 3/4 patch for the first week), then on to Step 3 for 2 weeks. (Read the enclosed self-help guide for additional information.)

Apply one new patch every 24 hours on skin that is dry, clean and hairless.

Wash hands after applying or removing patch. Save pouch to use for patch disposal.

Dispose of the used patches by folding sticky ends together and putting in pouch.

The used patch should be removed and a new one applied to a different skin site at the same time each day.

If you have vivid dreams, you may remove the patch at bedtime and apply a new one in the morning.

Do not wear more than one patch at a time.

Do not cut patch in half or into smaller pieces.

Do not leave patch on for more than 24 hours because it may irritate your skin and loses strength after 24 hours.

To avoid possible burns, remove patch before undergoing any MRI (magnetic resonance imaging) procedures.

It is important to complete treatment. If you feel you need to use the patch for a longer period to keep from smoking, talk to your healthcare provider.

Other information

Store at room temperature

Inactive ingredients

acrylate adhesive, aluminized polyester, cellulose paper, methacrylic acid copolymer.

Questions

After-sales service contact information:

For any product inquiries, contact us: ParuiTrade@outlook.com

For family's protection patches are supplied in child-resistant containers.
DO NOT USE IF INDIVIDUAL PATCH IS OPEN OR TORN.

QUIT SMOKING
Nicotine Patches

Nicotine Patches
QUIT SMOKING

CONTROL CRAVINGS

ALL-DAY!

STEP 1

**Over 24 hours
Each patch contains 21 mg**

Helps You Quit Smoking
Controls Cravings
Fights Withdrawal Symptoms
100% Natural Plant Based

**30
Patches**

Trade Mark: WOFANLI
Lot number: NP8685

X003167 7188
Nicotine Patches Step 1 E-cotine Transdermal System
New

WOFANLI

Drug Facts

Active ingredient (in each patch) **Purpose**
Nicotine, 21 mg delivered over 24 hours Stop smoking aid

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider.
Smoking causes roughly harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- an allergy to adhesive tape or have skin problems, because you are more likely to get rashes.
- diabetes or are on diabetes medicine.
- history of seizures.

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug.
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

When using this product

- If you have vivid dreams or other sleep disturbances remove this patch at bedtime.

Stop use and ask a doctor if

- skin redness caused by the patch does not go away after four days, or if your skin swells, or you get a rash.
- irregular heartbeat or palpitations occur.
- you get symptoms of nicotine overdose, such as persistent nausea, vomiting, dizziness, weakness and rapid heartbeat.
- you have symptoms of an allergic reaction (such as difficulty breathing or rash).

Keep out of reach of children and pets.
Used patches contain enough nicotine to poison children and pets. If swallowed, seek medical help immediately or contact a Poison Control Center 1-800-222-1222 right away. Sew the pouch for patch disposal. Dispose of used patches by folding the sticky ends together and placing them in the pouch.

Directions

- If you are under 18 years of age, ask a doctor before use.
- Before using this product, read the enclosed self-help guide for complete directions and other information.
- Begin using the patch on your quit day.
- If you smoke more than 10 cigarettes per day, use the following schedule below:

Weeks 1 thru 4	Weeks 5 and 6	Weeks 7 and 8
STEP 1 use one 21 mg patch/day	STEP 2 use one 14 mg patch/day	STEP 3 use one 7 mg patch/day

- If you smoke 10 or less cigarettes per day, start with **Step 2** for 5 weeks, then **Step 3** for 2 weeks.
- Apply one new patch every 24 hours on skin that is dry, clean and hairless.
- Wash hands after applying or removing patch. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.
- The used patch should be removed and a new one applied to a different skin site at the same time each day.
- If you have vivid dreams, you may remove the patch at bedtime and apply a new one in the morning.
- Do not wear more than one patch at a time.
- Do not cut patch in half or into smaller pieces.
- Do not leave patch on for more than 24 hours because it may irritate your skin and lose strength after 24 hours.
- To avoid possible bias, remove patch before undergoing any MRI (magnetic resonance imaging) procedures.
- It is important to complete treatment. If you feel you need to use the patch for a longer period to keep from smoking, ask to your health care provider.

Other information

- store at room temperature

Inactive ingredients
acrylate adhesive, aluminumized polyester, cellulose paper, methacrylic acid copolymer.

After-sales service contact information:
If you have any doubt about our products, please contact us.
PainTrading@outlook.com

Address:
468 Zhongyuan Road, 102, Huhu Community, Pinghu Street, Longgang District, Shenzhen City.

Manufacturing company:
Shenzhen Pain Trading Co., Ltd.

See bottom panel for lot number and expiration date.

• Not for sale to persons under 18 years of age.
• Proof of age required.
• Not for sale in vending machines or from any source where proof of age cannot be verified.

* Statements regarding dietary supplements have not been evaluated by the FDA and are not intended to diagnose, treat, cure, or prevent any disease or health condition.

nicotine transdermal patches patch

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:85835-004
Route of Administration	TOPICAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)		NICOTINE	21 mg in 100
Inactive Ingredients			
Ingredient Name			Strength
METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:1) (UNII: 74G4R6TH13)			
ACRYLIC ACID (UNII: J94PBK7X8S)			
POLYESTER-7 (UNII: 0841698D2F)			

POWDERED CELLULOSE (UNII: SMD1X3XO9M)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:85835-004-01	30 in 1 BOX; Type 0: Not a Combination Product	10/15/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	10/15/2025	

Labeler - Shenzhen Paxi Trading Co., Ltd. (633886766)

Establishment

Name	Address	ID/FEI	Business Operations
Shenzhen Paxi Trading Co., Ltd.		633886766	manufacture(85835-004)

Revised: 10/2025

Shenzhen Paxi Trading Co., Ltd.