

VICKS DAYQUIL AND VICKS NYQUIL SEVERE COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride, and doxylamine succinate
The Procter & Gamble Manufacturing Company

Vicks[®] DayQuil[™] Severe and Vicks[®] NyQuil Severe Cold and Flu Convenience Pack

DayQuil Severe Cold & Flu

Drug Facts

Active ingredients (in each 15 mL)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Guaifenesin 200 mg

Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

Uses

- temporarily relieves common cold/flu symptoms:
 - nasal congestion
 - sinus congestion & pressure
 - cough due to minor throat & bronchial irritation
 - minor aches & pains
 - headache
 - fever
 - sore throat
 - reduces swelling of nasal passages
 - temporarily restores freer breathing through the nose
 - promotes nasal and/or sinus drainage
 - helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4 doses (30 mL each) in 24 hrs, which is the maximum daily amount for this product
- child takes more than 4 doses (15 mL each) in 24 hrs, which is the maximum daily amount for this product
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy Alert:

Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning:

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each 15 mL tablespoon contains:**sodium 47 mg
- store at no greater than 25°C and do not refrigerate.

Inactive ingredients

citric acid, FD&C Yellow No. 6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum

Questions?

1-800-362-1683

NyQuil Severe Cold & Flu

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg
Dextromethorphan HBr 20 mg
Doxylamine succinate 12.5 mg
Phenylephrine HCl 10 mg

Purpose

Pain reliever/fever reducer
Cough suppressant
Antihistamine
Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- minor aches & pains
- headache
- fever
- sore throat
- runny nose & sneezing
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert:

Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning:

If sore throat is severe, persists for more than 2 days, is accompanied or followed by

fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as

for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each 30 mL dose cup contains:**sodium 81 mg
- store at no greater than 25°C and do not refrigerate

Inactive ingredients

alcohol, citric acid, D&C Yellow No. 10, FD&C Yellow No. 6, FD&C Green No. 3, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose

Questions?

1-800-362-1683

TAMPER EVIDENT: Do not use if printed shrinkband on bottle is broken or missing.

Dist. by Procter & Gamble, Cincinnati OH 45202

PRINCIPAL DISPLAY PANEL - Convenience Pack

MAX

STRENGTH

DAY & NIGHT PACK

VICKS®

**DayQuil™
SEVERE**

COLD & FLU

Acetaminophen, Guaifenesin, Phenylephrine HCl, Dextromethorphan HBr

Headache, Fever, Sore Throat Minor Aches & Pains

Chest Congestion, Thins & Loosens Mucus

Nasal Congestion, Sinus Pressure

Cough

Non-Drowsy

Alcohol Free

NyQuil™

SEVERE

COLD & FLU

Acetaminophen, Phenylephrine HCl, Doxylamine Succinate, Dextromethorphan HBr

Headache, Fever, Sore Throat Minor Aches & Pains

Nasal Congestion, Sinus Pressure

Sneezing, Runny Nose

Cough

Nighttime Relief

Alcohol 10%

2 bottles - 1 DAYQUIL/1 NYQUIL 12 FL OZ (354 mL) EACH; TOTAL 24 OZ (708 mL)



VICKS DAYQUIL AND VICKS NYQUIL SEVERE COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride, and doxylamine succinate kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37000-045
---------------------	----------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37000-045-24	1 in 1 PACKAGE	07/31/2014	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity

DayQuil SEVERE

Drug Facts

Active ingredients (in each 30 mL)

Dextromethorphan HBr 20 mg
 Doxylamine succinate 15 mg
 Phenylephrine HCl 10 mg

Purpose
 Pain reliever/fever reducer
 Cough suppressant
 Antihistamine
 Nasal decongestant

Uses
 Temporarily relieves common cold/flu symptoms:
 • nasal congestion
 • sinus congestion & pressure
 • cough due to minor throat & bronchial irritation
 • cough to help you sleep
 • minor aches & pains
 • headache
 • fever
 • sore throat
 • reduces swelling of nasal passages
 • temporarily relieves nasal breathing through the nose
 • promotes nasal and/or sinus drainage

Warnings
 Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
 • more than 4 doses in 24 hours, which is the maximum daily amount for this product
 • with other drugs containing acetaminophen
 • 3 or more 800 mg tablets every 4 to 6 hours while using this product
 • for 3 weeks after stopping the 800 mg tablet. If you do not know if your prescription drug contains acetaminophen, ask a doctor or pharmacist before taking this product.

Do not use
 • with any other drug containing acetaminophen (prescription or over-the-counter), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) medicine for depression, psychiatric or emotional condition, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have
 • liver disease • heart disease • high blood pressure
 • thyroid disease • diabetes
 • a breathing problem or chronic cough that lasts or worsens with smoking, asthma, chronic bronchitis, or emphysema
 • trouble emptying the stomach (nausea)
 • a stomach/intestinal diet

Ask a doctor or pharmacist before use if you are
 • taking medicines or tranquilizers • taking the blood thinning drug warfarin

When using this product
 • do not use more than directed • excitability may occur, especially in children
 • marked dizziness may occur • avoid alcoholic drinks
 • do not drink or drive a motor vehicle or operate machinery
 • alcohol, caffeine, and tranquilizers may increase drowsiness

Stop use and ask a doctor if
 • you get nervous, dizzy or sleepless
 • your nose becomes dry or itchy
 • your eyes become red or itchy
 • your nose becomes dry or itchy
 • your nose becomes dry or itchy
 • your nose becomes dry or itchy
 • your nose becomes dry or itchy

Directions
 • Use only as directed • only use the dose cap provided
 • do not exceed 4 doses per 24 hrs
 • adults & children 12 yrs & over 1 30 mL every 4 hrs
 • children 6 to under 12 yrs 1 15 mL every 4 hrs
 • children 4 to under 6 yrs ask a doctor
 • children under 4 yrs do not use

Other information
 • each 30 mL contains sodium 31 mg
 • does not contain any artificial colors or flavors

Inactive ingredients acetaminophen, citric acid, D&B Yellow No. 10, FD&C Green No. 3, FD&C Yellow No. 6, Ferric glycolate, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium hydroxide, sodium chloride, sorbitol, sucrose

Questions? • 1-800-361-1883

NyQuil SEVERE

Drug Facts

Active ingredients (in each 15 mL)

Dextromethorphan HBr 10 mg
 Doxylamine succinate 7.5 mg
 Phenylephrine HCl 5 mg

Purpose
 Pain reliever/fever reducer
 Cough suppressant
 Antihistamine
 Nasal decongestant

Uses
 Temporarily relieves common cold/flu symptoms:
 • nasal congestion • sinus congestion & pressure
 • cough due to minor throat & bronchial irritation
 • cough to help you sleep • minor aches & pains • headache • fever • sore throat
 • reduces swelling of nasal passages
 • temporarily relieves nasal breathing through the nose
 • promotes nasal and/or sinus drainage
 • helps loosen phlegm (mucus) and thick bronchial secretions to rid the bronchial passageways of mucus and make coughs more productive

Warnings
 Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
 • more than 4 doses in 24 hours, which is the maximum daily amount for this product
 • with other drugs containing acetaminophen
 • 3 or more 800 mg tablets every 4 to 6 hours while using this product
 • for 3 weeks after stopping the 800 mg tablet. If you do not know if your prescription drug contains acetaminophen, ask a doctor or pharmacist before taking this product.

Do not use
 • with any other drug containing acetaminophen (prescription or over-the-counter), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) medicine for depression, psychiatric or emotional condition, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have
 • liver disease • heart disease • high blood pressure
 • thyroid disease • diabetes
 • a breathing problem or chronic cough that lasts or worsens with smoking, asthma, chronic bronchitis, or emphysema
 • trouble emptying the stomach (nausea)
 • a stomach/intestinal diet

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product, do not use more than directed.

Stop use and ask a doctor if
 • you get nervous, dizzy or sleepless
 • your nose becomes dry or itchy
 • your eyes become red or itchy
 • your nose becomes dry or itchy
 • your nose becomes dry or itchy
 • your nose becomes dry or itchy

Directions
 • Use only as directed • only use the dose cap provided
 • do not exceed 4 doses per 24 hrs
 • adults & children 12 yrs & over 1 30 mL every 4 hrs
 • children 6 to under 12 yrs 1 15 mL every 4 hrs
 • children 4 to under 6 yrs ask a doctor
 • children under 4 yrs do not use

Other information
 • each 15 mL contains sodium 17 mg
 • does not contain any artificial colors or flavors

Inactive ingredients acetaminophen, citric acid, FD&C Yellow No. 6, Ferric glycolate, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sodium hydroxide, sodium phosphate, sucrose

Questions? • 1-800-361-1883

MAX STRENGTH

DayQuil SEVERE

THE DAYTIME, COUGHING, ACHING, STUFFY HEAD, FEVER, SORE THROAT, CONGESTED CHEST, POWER THROUGH YOUR DAY, MEDICINE.

MAX STRENGTH

DayQuil SEVERE

Acetaminophen, Guaifenesin, Phenylephrine HCl, Doxylamine Succinate

DAY & NIGHT PACK

NyQuil SEVERE

Acetaminophen, Phenylephrine HCl, Doxylamine Succinate, Doxylamine Succinate

MAX STRENGTH

NyQuil SEVERE

THE NIGHTTIME, SNIFFLING, SNEEZING, COUGHING, ACHING, STUFFY HEAD, FEVER, BEST SLEEP WITH A COLD, MEDICINE.

Headache, Fever, Sore Throat, Minor Aches & Pains
 Chest Congestion, Thins & Loosens Mucus
 Nasal Congestion, Sinus Pressure
 Cough

Non-Drowsy
 Alcohol Free

Nighttime Relief
 Alcohol 10%

PARENTS:
 Learn how to use medicine safely.
www.StopMedicineAbuse.org

DIET. BY PROCTER & GAMBLE, CINCINNATI OH 45202

P&G
www.pg.com
www.inits.com
 Patents: www.pg.com/patents

2 BOTTLES - 1 DAYQUIL/1 NYQUIL 12 FL OZ (354 mL) EACH; TOTAL 24 FL OZ (708 mL)



Part 1	1 BOTTLE, PLASTIC	354 mL
Part 2	1 BOTTLE, PLASTIC	354 mL

Part 1 of 2

VICKS DAYQUIL SEVERE COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride solution

Product Information

Item Code (Source)	NDC:37000-810
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 15 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	orange	Score	
Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37000-810-12	354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/22/2013	

Part 2 of 2

VICKS NYQUIL SEVERE COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, and phenylephrine hydrochloride solution

Product Information

Item Code (Source)	NDC:37000-815
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
ALCOHOL (UNII: 3K9958V90M)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	

SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

Product Characteristics

Color	green	Score	
Shape		Size	
Flavor	ANISE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37000-815-12	354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/21/2014	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/31/2014	

Labeler - The Procter & Gamble Manufacturing Company (004238200)

Revised: 10/2024

The Procter & Gamble Manufacturing Company