

PAIN RELIEF CHILDRENS- acetaminophen suspension
Dolgenercorp, Inc. (DOLLAR GENERAL & REXALL)

Drug Facts

Active ingredient (in each 5 mL)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily reduces fever
- temporarily relieves minor aches and pains due to:
 - the common cold
 - flu
 - headache
 - sore throat
 - toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this

product

Ask a doctor before use if your child has

liver disease.

Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin.

When using this product

- **do not exceed recommended dosage (see overdose warning)**

Stop use and ask a doctor if

- new symptoms occur
- redness or swelling is present
- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days

These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not take more than directed (see overdose warning)**
- **shake well before using**
- mL= milliliter
- use only the enclosed dosing cup designed for use with this product. Do not use any other dosing device.
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give more than 5 days unless directed by a doctor
- find the right dose on chart below. If possible, use weight to dose, use age.

Weight (lb)	Age (year)	Dose (mL)
under 24	under 2	ask a doctor
24-35	2-3	5 mL
36-47	4-5	7.5 ml
48-59	6-8	10 mL
60-71	9-10	12.5 mL
72-95	11	15 mL

Other information

- **each 5 mL contains:** sodium 2 mg
- store between 20-25°C (68-77°F)
- do not refrigerate
- see bottom panel for lot number and expiration date

Inactive ingredients

acesulfame potassium, anhydrous citric acid, butylparaben, carboxymethylcellulose sodium, FD&C red #40, flavors, glycerin, high fructose corn syrup, microcrystalline cellulose, propylene glycol, purified water, sodium benzoate, sorbitol, xanthan gum

Questions or comments?

Call 1-888-309-9030

Principal Display Panel

Compare to active ingredients of Children's Tylenol® Oral Suspension**

For Ages 2 to 11 Years

Children's Pain Relief

Acetaminophen 160 mg per 5 mL

Oral Suspension

Pain Reliever/Fever Reducer

- alcohol free
- aspirin free
- ibuprofen free

Fast, effective relief of children's fever and pain

Cherry Flavor

FL OZ (mL)

*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Tylenol® Oral Suspension

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING

DISTRIBUTED BY DOLGENCORP, LLC

100 MISSION RIDGE

GOODLETSVILLE, TN 37072

Product Label

Drug Facts (continued)

Find the right dose on chart below. If possible, use weight to dose; otherwise, use age.

Weight (lb)	Age (year)	Dose (mL)**
under 24	under 2	ask a doctor
24-35	2-3	5 mL
36-47	4-5	7.5 mL
48-59	6-8	10 mL
60-71	9-10	12.5 mL
72-95	11	15 mL

*or as directed by a doctor

Other information

- each 5 mL contains sodium 2 mg
- store between 20-25°C (68-77°F) do not refrigerate
- see bottom panel for lot number and expiration date

Inactive ingredients acacia, polyethylene glycol, butylparaben, carbocymethylcellulose sodium, citric acid, FD&C red #40, flavors, glycerin, high fructose corn syrup, microcrystalline cellulose, propylene glycol, purified water, sodium benzoate, sorbitol, xanthan gum

Questions or comments?
Call 1-888-309-9030

*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Tylenol® Oral Suspension.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY OLD EAST MAIN CO.
100 MISSION RIDGE
GOODLETTSVILLE, TN 37072

100%
Satisfaction
Guaranteed!
(888) 309-9030

PLD-F231D F0007411

Compare to active ingredient of Children's Tylenol® Oral Suspension**

DG health

Compare to active ingredient of Children's Tylenol® Oral Suspension**

DG health

For Ages 2 to 11 Years

Children's

Pain Relief

**Acetaminophen
160 mg per 5 mL
Oral Suspension**

Pain Reliever/Fever Reducer

- Alcohol free
- Aspirin free
- Ibuprofen free



Fast, effective relief of children's fever and pain

Cherry Flavor

8 FL OZ (237 mL)

160mg PER 5 mL

For Ages 2 to 11 Years

Children's

Pain Relief

**Acetaminophen
160 mg per 5 mL
Oral Suspension**

Pain Reliever/Fever Reducer

- Alcohol free
- Aspirin free
- Ibuprofen free



Fast, effective relief of children's fever and pain

Cherry Flavor

8 FL OZ (237 mL)

160mg PER 5 mL

Drug Facts	Purpose
Active ingredient (in each 5 mL) Acetaminophen 160 mg	Pain reliever/fever reducer
Uses	
<ul style="list-style-type: none"> temporarily reduces fever temporarily relieves minor aches and pains due to: <ul style="list-style-type: none"> the common cold flu headache sore throat toothache 	
Warnings	
<p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:</p> <ul style="list-style-type: none"> more than 5 doses in 24 hours, which is the maximum daily amount with other drugs containing acetaminophen <p>Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: <ul style="list-style-type: none"> skin redness blisters rash If a skin reaction occurs, stop use and seek medical help right away. </p> <p>Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p>	
Do not use	
<ul style="list-style-type: none"> with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. if your child is allergic to acetaminophen or any of the inactive ingredients in this product. 	
<p>Ask a doctor before use if your child has liver disease.</p> <p>Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.</p>	
When using this product, do not exceed recommended dose (see Overdose warning).	
Stop use and ask a doctor if	
<ul style="list-style-type: none"> new symptoms occur redness or swelling is present pain gets worse or lasts more than 5 days fever gets worse or lasts more than 3 days These could be signs of a serious condition.	
Keep out of reach of children.	
<p>Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.</p>	
Directions	
<ul style="list-style-type: none"> this product does not contain directions or complete warnings for adult use do not take more than directed (see Overdose warning) shake well before using mL = milliliter use only the enclosed dosing cup designed for use with this product. Do not use any other dosing device. if needed, repeat dose every 4 hours while symptoms last do not give more than 5 times in 24 hours do not give more than 5 doses unless directed by a doctor 	



Lot No:
Exp. Date:

DOLLAR GENERAL Children's Pain Relief Cherry Flavor

PAIN RELIEF CHILDRENS			
acetaminophen suspension			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55910-282
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL	

Inactive Ingredients

Ingredient Name	Strength
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
BUTYLPARABEN (UNII: 3QPI1U3FV8)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-282-08	1 in 1 BOX	04/30/2015	
1		237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	04/30/2015	

Labeler - Dolgencorp, Inc. (DOLLAR GENERAL & REXALL) (068331990)

Revised: 12/2023

Dolgencorp, Inc. (DOLLAR GENERAL & REXALL)