

**VICKS DAYQUIL AND VICKS NYQUIL COLD AND FLU- acetaminophen,
dextromethorphan hydrobromide, phenylephrine hydrochloride, and
doxylamine succinate
The Procter & Gamble Manufacturing Company**

**Vicks® DayQuil™ and Vicks® NyQuil™
Cold & Flu Convenience Pack**

NyQuil Cold & Flu

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg
Dextromethorphan HBr 30 mg
Doxylamine succinate 12.5 mg

Purpose

Pain reliever/ fever reducer
Cough suppressant
Antihistamine

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever
- runny nose and sneezing

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert:

Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening

- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning:

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each 30 mL contains:** sodium 23 mg
- store at no greater than 25°C and do not refrigerate and do not refrigerate

Inactive ingredients

alcohol, citric acid, D&C Yellow No. 10, FD&C Green No. 3, FD&C Yellow No. 6, flavor, glycerin, propylene glycol, purified water, saccharin sodium,, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions?

1-800-362-1683

Dist. by Procter & Gamble, Cincinnati OH 45202

DayQuil Cold & Flu

Drug Facts

Active ingredients (in each 15 mL)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/Fever reducer
Cough suppressant
Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion
- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if

- adults take more than 4 doses (30 mL each) in 24 hours, which is the maximum daily amount for this product
- child take more than 4 doses (15 mL each) in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy Alert:

Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning:

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease

- diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

When using this product,

do not use more than directed.

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough get worse or last more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each 15 mL contains:** sodium 46 mg
- store at no greater than 25°C and do not refrigerate.

Inactive ingredients

citric acid, FD&C Yellow No. 6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum

Questions?

1-800-362-1683

TAMPER EVIDENT: Do not use if printer shrinkband on bottle is broken or missing.

PRINCIPAL DISPLAY PANEL - Convenience Pack

POWERFUL

RELIEF

Day & Night Pack

VICKS®

DayQuil™

COLD & FLU

Acetaminophen, Phenylephrine HCl, Dextromethorphan HBr,

Headache, Fever, Sore Throat, Minor Aches & Pains

Nasal Congestion, Sinus Pressure

Cough

NyQuil™

COLD & FLU

Acetaminophen, Doxylamine Succinate, Dextromethorphan HBr

Headache, Fever, Sore Throat, Minor Aches & Pains

Sneezing, Runny Nose

Cough

Nighttime Relief

Alcohol 10%

2 BOTTLES - 1 DAYQUIL/1 NYQUIL 12 FL OZ (354 mL) EACH; TOTAL 24 FL OZ (708 mL)

DayQuil

NyQuil

DAY & NIGHT PACK

POWERFUL RELIEF **POWERFUL RELIEF** **DAY & NIGHT PACK** **POWERFUL RELIEF**



DayQuil **NyQuil**

THE DAYTIME, COUGHING, ACHING, STUFFY HEAD, FEVER, SORE THROAT, **POWER THROUGH YOUR DAY, MEDICINE.**

THE NIGHTTIME, SNIFFLING, SNEEZING, COUGHING, ACHING, FEVER, **BEST SLEEP WITH A COLD, MEDICINE.**

Headache, Fever, Sore Throat, Minor Aches & Pains

Nasal Congestion, Sinus Pressure, Cough

Non-Drowsy Alcohol Free

2 BOTTLES - 1 DAYQUIL/1 NYQUIL 12 FL OZ (354 mL) EACH; TOTAL 24 FL OZ (708 mL)



Headache, Fever, Sore Throat, Minor Aches & Pains

Sneezing, Runny Nose, Cough

Nighttime Relief Alcohol 10%

PARENTS:
Learn About Your Medicine Above
www.StopMedIn4Steps.org

DIST. BY PROCTER & GAMBLE, CINCINNATI OH 45202

P&G
www.pg.com
www.vicks.com
Patents: www.pg.com/patents

VICKS DAYQUIL AND VICKS NYQUIL COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride, and doxylamine succinate kit

Product Information

Product Type HUMAN OTC DRUG **Item Code (Source)** **NDC:37000-033**

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37000-033-24	1 in 1 PACKAGE; Type 0: Not a Combination Product	08/01/2013	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE, PLASTIC	354 mL
Part 2	1 BOTTLE, PLASTIC	354 mL

Part 1 of 2

VICKS DAYQUIL COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, and phenylephrine hydrochloride liquid

Product Information

Item Code (Source) NDC:69423-788

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	orange	Score	
Shape		Size	
Flavor	APRICOT	Imprint Code	
Contains			

Packaging

Marketing Start Marketing End

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69423-788-12	354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/22/2011	

Part 2 of 2

VICKS NYQUIL COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, and doxylamine succinate liquid

Product Information

Item Code (Source)	NDC:69423-789
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
XANTHAN GUM (UNII: TTV12P4NEE)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

Product Characteristics

Color	green	Score	
Shape		Size	
Flavor	ANISE, MENTHOL	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69423-789-12	354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/01/2010	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/01/2013	

Labeler - The Procter & Gamble Manufacturing Company (004238200)

Revised: 10/2023

The Procter & Gamble Manufacturing Company