

ACID REDUCER - omeprazole tablet, delayed release

Bi-Mart

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Drug Facts

Active ingredient (in each tablet)

Omeprazole delayed-release tablet 20 mg (equivalent to 20.6 mg omeprazole magnesium)

Purpose

Acid reducer

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert:

Do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking:

- warfarin, clopidogrel or cilostazol (blood-thinning medicines)
- prescription antifungal or anti-yeast medicines
- diazepam (anxiety medicine)
- digoxin (heart medicine)
- tacrolimus or mycophenolate mofetil (immune system medicines)
- prescription antiretrovirals (medicines for HIV infection)
- methotrexate (arthritis medicine)

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor

- swallow whole. Do not chew or crush tablets.

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
- children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20° to 25°c (68° to 77° F) and protect from moisture

Inactive ingredients

crospovidone, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer dispersion, microcrystalline cellulose, polyethylene glycol, polysorbate 80, red iron oxide, silicified microcrystalline cellulose, sodium hydroxide, sodium stearyl fumarate, sugar spheres [which contains liquid glucose, starch (maize) and sucrose], talc, titanium dioxide, triethyl citrate and yellow iron oxide.

Questions?

Call **1-800-935-6737**

*This product is not manufactured or distributed by the owner of the registered trademark Prilosec OTC®

Distributed by:

BI-MART

Eugene, OR 97402

Made in India

Code: TS/DRUGS/22/2009

PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 20 mg (14 Tablet Bottle)

BI-MART NDC 37835-189-14

ACID REDUCER

Omeprazole

Delayed-Release Tablets 20 mg

TREATS FREQUENT HEARTBURN

OCCURRING 2 OR MORE DAYS A WEEK

14 TABLETS

ONE 14-DAYCOURSE OF TREATMENT

Top Ply

BI-MART NDC 37 835-189-14

ACID REDUCER
OMEPRAZOLE
DELAYED-RELEASE TABLETS 20 mg

14 TABLETS TREATS FREQUENT HEARTBURN OCCURRING 2 OR MORE DAYS A WEEK ONE 14-DAY COURSE OF TREATMENT

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

TAMPER EVIDENT - Do not use if seal over bottle opening is broken or missing.

Made in India
Code No.: TS/DRUGS/22/2009 LM-3144

Lift Here

Distributed by:
BI-MART
Eugene, OR 97402
68911-014-11-0

No Varnish

Top Ply (Page #1)

Active ingredient (in each tablet)
Omeprazole delayed-release tablet 20 mg (equivalent to 20.6 mg omeprazole magnesium)

Purpose
Acid reducer

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings **Allergy alert:** Do not use if you are allergic to omeprazole Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain. These may be signs of a serious condition. See your doctor. Ask a doctor before use if you have: ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain Ask a doctor or pharmacist before use if you are taking: ■ warfarin, clopidogrel or cilostazol (blood-thinning medicines) ■ prescription antifungal or anti-yeast medicines ■ diazepam (anxiety medicine) ■ digoxin (heart medicine) ■ tacrolimus or mycophenolate mofetil (immune system medicines) ■ prescription antiretrovirals (medicines for HIV infection) ■ methotrexate (arthritis medicine) Stop use and ask a doctor if: ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain ▶

Back of Top Ply (Page #2)

Bottom Ply

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. **Directions** ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours **14-Day Course of Treatment** ■ swallow 1 tablet with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ swallow whole. Do not chew or crush tablets. **Repeated 14-Day Courses (if needed)** ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor ■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition. **Other information** ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20° to 25°C (68° to 77°F) and protect from moisture

Inactive ingredients crospovidone, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer dispersion, microcrystalline cellulose, polyethylene glycol, polysorbate 80, red iron oxide, silicified microcrystalline cellulose, sodium hydroxide, sodium stearyl fumarate, sugar spheres [which contains liquid glucose, starch (maize) and sucrose], talc, titanium dioxide, triethyl citrate and yellow iron oxide. **Questions?** Call 1-800-935-6737

Base (Page #3)

PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 20 mg Container Carton Label

BI-MART NDC 37835-189-14
ACID REDUCER
Omeprazole Delayed-Release
Tablets 20 mg
TREATS FREQUENT HEARTBURN

**OCCURRING 2 OR MORE DAYS A WEEK
14 TABLETS
ONE 14-DAY COURSE OF TREATMENT**



ACID REDUCER

omeprazole tablet, delayed release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37835-189
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OMEPRAZOLE MAGNESIUM (UNII: 426QFE7XLK) (OMEPRAZOLE - UNII:KG60484QX9)	OMEPRAZOLE	20 mg

Inactive Ingredients

Ingredient Name	Strength
CROSPROVIDONE (35 .MU.M) (UNII: 40UAA97IT9)	
GLYCERYL MONOSTEARATE (UNII: 230OU9XXE4)	
HYDROXYPROPYL CELLULOSE (90000 WAMW) (UNII: UKE75GEA7F)	
HYPROMELLOSE 2910 (5 MPA.S) (UNII: R75537T0T4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
METHACRYLIC ACID - ETHYL ACRYLATE COPOLYMER (1:1) TYPE A (UNII: NX76LV5T8J)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL 4000 (UNII: 4R4HFI6D95)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
SODIUM STEARYL FUMARATE (UNII: 7CV7WJK4UI)	
DEXTROSE, UNSPECIFIED FORM (UNII: IY9XDZ35W2)	
STARCH, CORN (UNII: O8232NY3SJ)	
SUCROSE (UNII: C151H8M554)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
HYDROXYPROPYL CELLULOSE (45000 WAMW) (UNII: 8VAB711C5E)	
HYPROMELLOSE 2910 (6 MPA.S) (UNII: OWZ8WG20P6)	

Product Characteristics

Color	PINK	Score	no score
Shape	RECTANGLE (Oblong)	Size	14mm
Flavor		Imprint Code	Z;69
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37835-189-14	1 in 1 CARTON	06/06/2018	
1		14 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:37835-189-42	3 in 1 CARTON	06/06/2018	
2		14 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End
-----------	---------------------------------	-----------------	---------------

Category	Citation	Date	Date
ANDA	ANDA206877	06/06/2018	

Labeler - Bi-Mart (027630078)

Registrant - Aurohealth LLC (078728447)

Establishment

Name	Address	ID/FEI	Business Operations
Aurobindo Pharma Limited		650381903	ANALYSIS(37835-189) , MANUFACTURE(37835-189)

Revised: 6/2024

Bi-Mart