

**ANTACID AND ANTIGAS- calcium carbonate, magnesium hydroxide,
simethicone liquid
Meijer, Inc.**

mei 655 mylanta classic

Active ingredients (in each 10 mL dose)

Calcium carbonate 800 mg
Magnesium hydroxide 270 mg
Simethicone 80 mg

Purposes

Antacid

Antigas

Uses

relieves

- heartburn
- sour stomach
- acid indigestion
- the symptoms referred to as gas
- upset stomach associated with these symptoms

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are

presently taking a prescription drug.

Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last for more than 2 weeks

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Directions

- shake well before use
- adults and children 12 years of age and older: take 10 mL to 20 mL (1 to 2 doses)

between meals as needed, at bedtime,

or as directed by a doctor

- do not take more than 60 mL (6 doses) in any 24-hour period or use the maximum dosage for more than 2 weeks
- children under 12 years of age: ask a doctor
- measure with dosing cup provided
- mL = milliliter

Other information

- **each 10 mL dose contains:** calcium 320 mg, magnesium 115 mg
- do not freeze
- store at room temperature tightly closed

Inactive ingredients

benzyl alcohol, flavor, purified water, simethicone emulsion, sorbic acid, sorbitol solution, sucralose, xanthan gum

Questions or comments?

1-800-540-3765

package Label



NDC 79481-0655-1

Compare to the Mylanta®
Classic Active
ingredients*

Antacid & Antigas

FAST RELIEF

- HEARTBURN
- ACID INDIGESTION
- PRESSURE
- BLOATING

CITRUS-MINT FLAVOR

Liquid Antacid +
Anti-Gas

12 FL OZ (355 mL) 

225-07522-0 REV 1025

Drug Facts

**TAMPER-EVIDENT: DO NOT USE IF
IMPRINTED NECKBAND IS MISSING OR BROKEN.**

Active ingredients (in each 10 mL dose)

Calcium carbonate 800 mg.....	Antacid
Magnesium hydroxide 270 mg.....	Antacid
Simethicone 80 mg	Antigas

Purposes

Uses relieves • heartburn • sour stomach
• acid indigestion • the symptoms referred to as gas
• upset stomach associated with these symptoms

Warnings

Ask a doctor before use if you have

- kidney disease • a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are
presently taking a prescription drug. Antacids may interact
with certain prescription drugs.

Stop use and ask a doctor if symptoms last for more than 2 weeks

**If pregnant or breast-feeding, ask a health professional
before use.**

Keep out of reach of children.

Directions

- shake well before use
- adults and children 12 years of age and older: take 10 mL
to 20 mL (1 to 2 doses) between meals as needed, at
bedtime, or as directed by a doctor
- do not take more than 60 mL (6 doses) in any 24-hour
period or use the maximum dosage for more than 2 weeks
- children under 12 years of age: ask a doctor
- measure with dosing cup provided • mL=milliliter

Other information

- each 10 mL dose contains:
calcium 320 mg, magnesium 115 mg
- do not freeze • store at room temperature tightly closed

Inactive ingredients benzyl alcohol, flavor, purified
water, simethicone emulsion, sorbic acid, sorbitol solution,
sucralose, xanthan gum

Questions or comments? 1-800-540-3765

*This product is not manufactured or distributed by Kenvue Inc., owner of
the registered trademark Mylanta®. PID 5692498

DIST. BY MEIJER
DISTRIBUTION, INC.
GRAND RAPIDS, MI 49544
www.meijer.com
925-07522-0 REV GC1025

7 08820 00016 2

ANTACID AND ANTIGAS

calcium carbonate, magnesium hydroxide, simethicone liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:79481-0655
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CALCIUM CARBONATE (UNII: H0G9379FGK) (CALCIUM CATION -	CALCIUM CARBONATE	800 mg

UNII:2M83C4R6ZB)	CALCIUM CARBONATE	in 10 mL
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	80 mg in 10 mL
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838)	MAGNESIUM HYDROXIDE	270 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
BENZYL ALCOHOL (UNII: LKG8494WBH)	
SORBIC ACID (UNII: X045WJ989B)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	
WATER (UNII: 059QF0KO0R)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	CITRUS (citrus mint)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79481-0655-1	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	12/01/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	12/01/2025	

Labeler - Meijer, Inc. (006959555)

Registrant - GCP Laboratories (965480861)

Establishment

Name	Address	ID/FEI	Business Operations
GCP Laboratories		965480861	manufacture(79481-0655) , pack(79481-0655)