

**MUCUS RELIEF ALL IN ONE MAXIMUM STRENGTH- acetaminophen,
dextromethorphan hbr, guaifenesin, phenylephrine hcl liquid
Walgreens**

Drug Facts

Active ingredients (in each 20 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Guaifenesin 400 mg

Phenylephrine HCl 10 mg

Purposes

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - nasal congestion
 - minor aches and pains
 - sore throat
 - headache
 - stuffy nose
 - sinus congestion and pressure
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for children under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- thyroid disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

When using this product,

do not use more than directed.

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- dose as follows or as directed by a doctor
- adults and children 12 years of age and older: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- **each 20 mL contains:** sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

anhydrous citric acid, disodium EDTA,, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Principal Display Panel

Compare to Maximum Strength Mucinex® Fast-Max All-In=One Cold & Flu Active ingredients††

DAYTIME

Cold & Flu

ACETAMINOPHEN 650 mg /

PAIN RELIEVER / FEVER REDUCER

DEXTROMETHORPHAN HBr 20 mg

COUGH SUPPRESSANT

GUAIFENESIN 400 mg / EXPECTORANT

PHENYLEPHRINE HCl 10 mg /

NASAL DECONGESTANT

MAXIMUM STRENGTH

- Relieves aches, fever, sore throat, nasal & chest congestion, sinus congestion & pressure
- Controls cough
- Thins & loosens mucus
- 12 years & older

FL OZ (mL)

††This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® All-in-One Cold & Flu.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY WALGREENS CO.

200 WILMOT RD., DEERFIELD, IL 60015

Package Label

Walgreens

Compare to Maximum Strength Mucinex® Fast-Max® All-In-One Cold & Flu active ingredients!

DAYTIME NDC 0363-0886-06

Cold & Flu

ACETAMINOPHEN 650 mg / PAIN RELIEVER / FEVER REDUCER
 DEXTROMETHORPHAN HBr 20 mg / COUGH SUPPRESSANT
 GUAIFENESIN 400 mg / EXPECTORANT
 PHENYLEPHRINE HCl 10 mg / NASAL DECONGESTANT

MAXIMUM STRENGTH

- Relieves aches, fever, sore throat, nasal & chest congestion, sinus congestion & pressure
- Controls cough
- Thins & loosens mucus
- 12 years & older

6 FL OZ (177 mL) LIQUID

*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Mucinex® Maximum Strength Mucinex® Fast-Max® All-In-One Cold & Flu.
 PLD-A574A LB006719
 ITEM 907517 W10389-10/15-F
 ORG1019-F

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Mucinex® Fast-Max® Maximum Strength All-In-One Cold & Flu.

PARENTS:
 Learn about teen medicine abuse
www.StopMedicineAbuse.org

PLD-A574A LB006719

Drug Facts

Active ingredients (in each 20 mL)	Purposes
Acetaminophen 650 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg.....	Cough suppressant
Guaifenesin 400 mg.....	Expectorant
Phenylephrine HCl 10 mg.....	Nasal decongestant

PEEL CORNER FOR MORE DRUG FACTS

Drug Facts (continued)

Uses

- temporarily relieves these common cold and flu symptoms
 - cough
 - nasal congestion
 - minor aches and pains
 - sore throat
 - headache
 - stuffy nose
 - sinus congestion and pressure
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

- Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:
- more than 4,000 mg of acetaminophen in 24 hours
 - with other drugs containing acetaminophen
 - 3 or more alcoholic drinks daily while using this product
- Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:
- skin reddening
 - blisters
 - rash
- If a skin reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for children under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- thyroid disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

PEEL CORNER FOR MORE DRUG FACTS

Drug Facts (continued)

When using this product, do not use more than directed.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse, or lasts more than 7 days
- fever gets worse, or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period

Drug Facts (continued)

- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

WALGREENS Daytime Cold & Flu

MUCUS RELIEF ALL IN ONE MAXIMUM STRENGTH

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-0886
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT9KYH)	DEXTROMETHORPHAN	20 mg

(DEXTROMETHORPHAN - UNII:7355X3ROTS)	HYDROBROMIDE	in 20 mL
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 20 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYL GALLATE (UNII: 8D45NN7V92)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0886-06	177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/27/2021	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/27/2021	

Labeler - Walgreens (008965063)