

RYCONTUSS- chlorpheniramine maleate, dextromethorphan hydrobromide, phenylephrine hydrochloride liquid
Dextrum Laboratories, Inc

RYCONTUSS

ACTIVE INGREDIENT SECTION

ACTIVE INGREDIENT SECTION

CHLORPHENIRAMINE MALEATE 2MG

DEXTROMETHORPHAN HYDROBROMIDE 10MG

PHENYLEPHRINE HYDROCHLORIDE 5MG



INACTIVE INGREDIENT SECTION

INACTIVE INGREDIENT SECTION:

CITRIC ACID, D&C RED#33, FD&C RED#40, NATURAL FLAVOR, METHYLPARABEN, PROPYLENE GLYCOL, PROPYLPARABEN, PURIFIED WATER, SODIUM CITRATE AND SUCRALOSE.



ASK A DOCTOR SECTION

ASK A DOCTOR SECTION:

ask A DOCTOR OR PHARMACIST IF YOU ARE TAKING SEDATIVES OR TRANQUILIZERS.



DO NOT USE SECTION

DO NOT USE SECTION:

DO NOT USE TO SEDATE A CHILD OR TO MAKE A CHILD SLEEPY.

Do not use if you are now taking a prescription monoamine oxidase inhibitor(MAOI) (certain drugs for depression, psychiatric or emotional condition, or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI ask a doctor or pharmacist before taking this product.

Drug Facts (continued)

These could be signs of a serious condition.
If pregnant or breast feeding ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a poison control center right away.

Directions	
Adults and children 12 or older	2 tsp (10 mL) every 4 hours
Children age 6 - 12	1 tsp (5 mL) every 4 hours
Children younger than 6	Ask a doctor

Other information
• Store at controlled room temperature between 15-30 C (59-86F) • Avoid excessive heat or humidity.

Inactive ingredients
Citric acid, D&G red #33, FD&C red #40, natural flavor, methylparaben, propylene glycol, propylparaben, purified water, sodium citrate and sucrose.

Questions or Comments
Phone: 1-281-548-0016, Labor Hours 8:00 am to 5:00 pm

Manufactured for:
Olanopharma Inc.
4287 Constanza, Ave. Villa Del Carmen, Ponce,
PR 00716, United States. Product of Puerto Rico

Lot #

Exp. Date

NDC 76117 - 001 - 16

RYCONTUSS

Liquid

CHERRY
FLAVOR



ANTITUSSIVE NASAL
DECONGESTANT
AND ANTIHISTAMINE

16 fl oz (1 pint) (473 mL)

Drug Facts

Active Ingredients (in each 5mL teaspoonful)	Purpose
Chlorpheniramine Maleate 2 mg	Antihistamine
Dextromethorphan Hydrobromide 10mg	Cough Suppressant
Phenylephrine Hydrobromide 5mg	Nasal Decongestant

Uses

- For the temporary relief of runny nose, sneezing, itching of the nose or throat, and itchy, watery eyes due to hay fever.
- Temporarily relieves cough due to minor throat and bronchial irritation occurring with the common cold.
- Temporarily restores free breathing through the nose.

Warnings**Do not use**

- To sedate a child or to make a child sleepy.
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional condition, or for 2 weeks after stopping the MAOI drug). If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

ASK A DOCTOR BEFORE IF YOU HAVE:

- Heart disease. • Thyroid disease. • Trouble urinating due to enlarged prostate gland. • Glaucoma. • Cough that occurs with too much phlegm (mucus). • Breathing problems or persistent or chronic cough that last such as occurs with smoking, asthma, chronic bronchitis or emphysema.

ASK A DOCTOR OR PHARMACIST: If you are taking sedatives or tranquilizers.**WHEN USING THIS PRODUCT**

- Do not use more than directed. • May cause marked drowsiness.
- Avoid alcoholic beverages. • Alcohol, sedatives and tranquilizers may increase drowsiness. Be careful when driving a motor vehicle or operating machinery.
- Toxicity may occur, especially with children.

STOP USE AND ASK A DOCTOR IF

- You get nervous, dizzy, or sleepless. • Cough last more than 7 days, comes back, or is accompanied by fever, rash, or persistent headaches.





Pregnancy or breast feeding section

Pregnancy or Breast Feeding Section:

If pregnancy or breaft-feeding, ask a heath professional before use

Drug Facts (continued)

These could be signs of a serious condition.
If pregnant or breast feeding ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a poison control center right away.

Directions	
Adults and children 12 or older	2 tsp (10 mL) every 4 hours
Children age 6 - 12	1 tsp (5 mL) every 4 hours
Children younger than 6	Ask a doctor

Other information
• Store at controlled room temperature between 15-30 C (59-86F) • Avoid excessive heat or humidity.

Inactive ingredients
Citric acid, D&C red #33, FD&C red #40, natural flavor, methylparaben, propylene glycol, propylparaben, purified water, sodium citrate and sucrose.

Questions or Comments
Phone: 1-281-548-0016, Labor Hours 9:00 am to 5:00 pm

Manufactured for:
Olanapharma Inc.
4287 Constanza, Ave. Villa Del Carmen, Ponce,
PR 00716, United States. Product of Puerto Rico

Lot #

Exp. Date

NDC 76117 - 001 - 16

RYCONTUSS

Liquid

CHERRY
FLAVOR



ANTITUSSIVE NASAL
DECONGESTANT
AND ANTIHISTAMINE

16 fl oz (1 pint) (473 mL)

Drug Facts

Active Ingredients (in each 5mL teaspoonful)	Purpose
Chlorpheniramine Maleate 2 mg	Antihistamine
Dextromethorphan Hydrobromide 10mg	Cough Suppressant
Phenylephrine Hydrobromide 5mg	Nasal Decongestant

Uses

- For the temporary relief of runny nose, sneezing, itching of the nose or throat, and itchy, watery eyes due to hay fever.
- Temporarily relieves cough due to minor throat and bronchial irritation occurring with the common cold.
- Temporarily restores free breathing through the nose.

Warnings**Do not use**

- To sedate a child or to make a child sleepy.
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional condition, or for 2 weeks after stopping the MAOI drug). If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

ASK A DOCTOR BEFORE IF YOU HAVE:

- Heart disease. • Thyroid disease. • Trouble urinating due to enlarged prostate gland. • Glaucoma. • Cough that occurs with too much phlegm (mucus). • Breathing problems or persistent or chronic cough that last such as occurs with smoking, asthma, chronic bronchitis or emphysema.

ASK A DOCTOR OR PHARMACIST: If you are taking sedatives or tranquilizers.

WHEN USING THIS PRODUCT

- Do not use more than directed. • May cause marked drowsiness. • Avoid alcoholic beverages. • Alcohol, sedatives and tranquilizers may increase drowsiness. Be careful when driving a motor vehicle or operating machinery. • Usability may occur, especially with children.

STOP USE AND ASK A DOCTOR IF

- You get nervous, dizzy, or sleepless. • Cough last more than 7 days, comes back, or is accompanied by fever, rash, or persistent headaches.



Drug Facts (continued)

These could be signs of a serious condition.
If pregnant or breast feeding ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a poison control center right away.

Directions	
Adults and children 12 or older	2 tsp (10 mL) every 4 hours
Children age 6 - 12	1 tsp (5 mL) every 4 hours
Children younger than 6	Ask a doctor

Other information
• Store at controlled room temperature between 15-30 C (59-86F) • Avoid excessive heat or humidity.

Inactive ingredients
Citric acid, D&G red #33, FD&C red #40, natural flavor, methylparaben, propylene glycol, propylparaben, purified water, sodium citrate and sucrose.

Questions or Comments
Phone: 1-787-548-0016, Labor Hours 8:00 am to 5:00 pm

Manufactured for:
Olanapharma Inc.
4287 Constancia Ave. Villa Del Carmen, Ponce,
PR 00716, United States. Product of Puerto Rico

Lot #

Exp. Date

NDC 76117 - 001 - 16

RYCONTUSS

Liquid

CHERRY
FLAVOR



ANTITUSSIVE NASAL
DECONGESTANT
AND ANTIHISTAMINE

16 fl oz (1 pint) (473 mL)

Drug Facts

Active Ingredients (in each 5mL teaspoonful)	Purpose
Chlorpheniramine Maleate 2 mg	Antihistamine
Dextromethorphan Hydrobromide 10mg	Cough Suppressant
Phenylephrine Hydrobromide 5mg	Nasal Decongestant

Uses

- For the temporary relief of runny nose, sneezing, itching of the nose or throat, and itchy, watery eyes due to hay fever.
- Temporarily relieves cough due to minor throat and bronchial irritation occurring with the common cold.
- Temporarily restores free breathing through the nose.

Warnings**Do not use**

- To sedate a child or to make a child sleepy.
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional condition, or for 2 weeks after stopping the MAOI drug). If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

ASK A DOCTOR BEFORE IF YOU HAVE:

- Heart disease. • Thyroid disease. • Trouble urinating due to enlarged prostate gland. • Glaucoma. • Cough that occurs with too much phlegm (mucus). • Breathing problems or persistent or chronic cough that last such as occurs with smoking, asthma, chronic bronchitis or emphysema.

ASK A DOCTOR OR PHARMACIST: If you are taking sedatives or tranquilizers.**WHEN USING THIS PRODUCT**

- Do not use more than directed. • May cause marked drowsiness. • Avoid alcoholic beverages. • Alcohol, sedatives and tranquilizers may increase drowsiness. Be careful when driving a motor vehicle or operating machinery. • Usability may occur, especially with children.

STOP USE AND ASK A DOCTOR IF

- You get nervous, dizzy, or sleepless. • Cough last more than 7 days, comes back, or is accompanied by fever, rash, or persistent headaches.





STOP USE SECTION.

STOP USE SECTION.

STOP USE OR ASK A DOCTOR IF:

YOU GET NERVOUS, DIZZY, OR SLEEPLESS

COUGH LAST MORE THAN 7 DAYS, COMES BACK, OR IS ACCOMPANIED BY FEVER, RASH OR PERSISTENT HEADACHES.



WARNINGS SECTION

WARNINGS SECTION

DO NOT USE TO SEDATE A CHILD OR TO MAKE A CHILD SLEEPY.

Do not use if you are now taking a prescription monoamine oxidase inhibitor(MAOI) (certain drugs for depression, psychiatric or emotional condition, or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI ask a doctor or pharmacist before taking this product.



INDICATION AND USAGE

INDICATION AND USAGE

Uses for the temporary relief of runny nose, sneezing, itching of the nose or throat, and itchy, watery eyes due to hay fever.

Temporarily relieves cough due to minor throat and bronchial irritation occurring with common cold.

Temporarily restore freer breathing through the nose.

Adults and children 12 or older-----2 tsp (10 ml) every 4 hours

Children age 6 - 12 ----- 1 tsp (5ml) every 4 hours

Children younger than 6. ----- Ask a doctor



Dosage and administration

Dosage and administration:

Adults and children 12 or older-----2 tsp (10 ml) every 4 hours

Children age 6 - 12 ----- 1 tsp (5ml) every 4 hours

Children younger tahn 6. ----- Ask a doctor



Package Label-Principal Display Panel

Package Label-Principal Display Panel

NDC 76117 - 001 - 16

RYCONTUSS

LIQUID

CHERRY FLAVOR

ANTITUSSIVE NASAL

DECONGESTANT AND

ANTIHISTAMINE

16 FL OZ (1 PINT) (473 ML)



RYCONTUSS

chlorpheniramine maleate, dextromethorphan hydrobromide, phenylephrine hydrochloride liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:65852-021
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 5 mL
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg in 5 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID (UNII: 2968PHW8QP)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	

METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
WATER (UNII: 059QF0KO0R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65852-021-16	473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/31/2011	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/31/2011	

Labeler - Dextrum Laboratories, Inc (007392322)

Revised: 10/2025

Dextrum Laboratories, Inc