

LIL DRUG STORE PAIN RELIEF ALL DAY- naproxen sodium tablet
Lil' Drug Store Products, Inc.

All Day Pain Relief, Lil' Drug Store ®

Drug Facts

Active ingredient (in each tablet)

Naproxen sodium 220 mg (naproxen 200 mg) (NSAID) ¹

¹ nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - headache
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert

Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic.
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better.
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose

adults and children 12 years and over	<ul style="list-style-type: none">• take 1 tablet every 8 to 12 hours while symptoms last• for the first dose, you may take 2 tablets within the first hour• do not exceed 2 tablets in any 8- to 12-hour period• do not exceed 3 tablets in a 24-hour period
over age 65	<ul style="list-style-type: none">• do not take more than 1 tablet every 12 hours unless directed by a doctor
children under 12 years	<ul style="list-style-type: none">• ask a doctor

Other information

- read all warnings and directions before use. Keep vial.
- **each tablet contains:** sodium 20 mg
- store at room temperature 20-25°C (68-77°F).
- avoid high humidity and excessive heat above 40°C (104°F).
- **do not use if vial is open or damaged or safety seal has been removed**

Inactive ingredients

croscarmellose sodium, FD&C blue #2, macrogol, magnesium stearate, polyvinyl alcohol, povidone, pregelatinized starch, talc, titanium dioxide

Questions or comments?

call toll-free 1-877-507-6516 (M-F 8 AM-4:30PM CST)

PRINCIPAL DISPLAY PANEL - 220 mg Tablet Vial Label

★ QUALITY GUARANTEED ★

Compare to the Active Ingredient in Aleve®**

All Day
Pain Relief
Naproxen Sodium (NSAID)
Pain Reliever/Fever Reducer, 220 mg

12
Tablets

Lil'
Drug Store®

OPEN HERE FOR DRUG FACTS INFORMATION

Product manufactured for: LIT Drug Store Products, Inc.
 3300 Earhart Lane SW, Cedar Rapids, IA 52404 97404-US-11-14-A
 **This product is not manufactured or distributed by Bayer Health Care LLC
 LIT Drug Store Products, Inc. does not own the Aleve® trademark.

Questions or comments? call toll free 1-877-507-6516
 (M-F 8 AM-4:30PM CST)

Inactive ingredients croscarmellose sodium, FD&C blue #2, microcyst, magnesium stearate, polyvinyl alcohol, polyvidone, pregelatinized starch, etc., titanium dioxide

Other information
 • read all warnings and directions before use. Keep vial removed.
 • each tablet contains sodium 20 mg
 • store at room temperature 20-25°C (68-77°F).
 • avoid high humidity and excessive heat above 40°C (104°F).
 • do not use if vial is open or damaged or safety seal has been removed.

Directions • do not take more than directed
 • the smallest effective dose should be used
 • drink a full glass of water with each dose
 • adults and children
 • take 1 tablet every 8 to 12 hours while symptoms last
 • for the first dose, you may take 2 tablets within the first hour
 • do not exceed 2 tablets in any 8- to 12-hour period
 • do not exceed 3 tablets in 24-hour period
 • do not take more than 1 tablet every 8 hours unless directed by a doctor
 • children under 12 years • ask a doctor

Drug Facts (continued)
 When using this product
 • take with food or milk if stomach upset occurs
 • the risk of heart attack or stroke may increase if you use more than directed or for longer than directed
 Stop use and ask a doctor if
 • you experience any of the following signs of stomach bleeding:
 • feel faint • vomit blood • have bloody or black stools
 • have stomach pain that does not get better
 • pain gets worse or lasts more than 10 days
 • fever gets worse or lasts more than 3 days
 • you have difficulty swallowing
 • it feels like the pills are stuck in your throat
 • redness or swelling is present in the pain area
 • any new symptoms appear
 If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless expressly directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)



★ QUALITY GUARANTEED ★
 Compare to the Active Ingredient in Aleve®**

All Day
Pain Relief
 Naproxen Sodium (NSAID)
 Pain Reliever/Fever Reducer, 220 mg

SAFETY SEALED FOR YOUR PROTECTION
 TEAR HERE TO OPEN

Drug Facts

Active ingredient (in each tablet)	Purpose
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*	Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug	


Uses • temporarily relieves minor aches and pains due to:
 • minor pain of arthritis • muscular aches • backache
 • menstrual cramps • headache • toothache
 • the common cold
 • temporarily reduces fever

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 • hives • facial swelling • asthma (wheezing)
 • shock • skin reddening • rash • blisters
 If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
 • are age 60 or older
 • have had stomach ulcers or bleeding problems
 • take a blood thinning (anticoagulant) or steroid drug
 • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 • have 3 or more alcoholic drinks every day while using this product
 • take more or for a longer time than directed

Do not use
 • if you have ever had an allergic reaction to any other pain reliever/fever reducer
 • right before or after heart surgery

Ask a doctor before use if
 • the stomach bleeding warning applies to you
 • you have a history of stomach problems, such as heartburn
 • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
 • you are taking a diuretic.
 • you have problems or serious side effects from taking pain relievers or fever reducers
 • you have asthma

Ask a doctor or pharmacist before use if you are
 • under a doctor's care for any serious condition
 • taking any other drug



LIL DRUG STORE PAIN RELIEF ALL DAY

naproxen sodium tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:66715-9740
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	blue	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	I3
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-9740-4	12 in 1 VIAL; Type 0: Not a Combination Product	12/31/2018	01/02/2027

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079096	01/27/2011	01/02/2027

Labeler - Lil' Drug Store Products, Inc. (093103646)

Revised: 3/2024

Lil' Drug Store Products, Inc.