

**SOLBAR FIFTY SPF50- solbar fifty spf50 cream**  
**Person and Covey**

-----  
**Solbar Fifty SPF50**

**OTC - PURPOSE SECTION**

Sun screen

**INDICATIONS & USAGE SECTION**

Helps prevent sunburn. If used as directed with other sun protection measures, decreases the risk of skin cancer and early skin aging caused by the sun.

**DOSAGE & ADMINISTRATION SECTION**

Apply liberally and evenly to all sun exposed areas of DRY skin 15 minutes before sun exposure. Reapply after 80 minutes of swimming or sweating and immediately after towel drying. Apply at least every 2 hours. For children under 6 months, ask a physician.

**OTC - KEEP OUT OF REACH OF CHILDREN SECTION**

Yes. If swallowed, get medical help or contact a Poison Control Center right away.

**OTC - ACTIVE INGREDIENT SECTION**

Octocrylene

Ethylhexyl Methoxycinnamate

Oxybenzone

Avobenzone

**INACTIVE INGREDIENT SECTION**

Purified Water

Propylene Glycol Diethylhexanoate

Dimethicone

PVP/Eicosene Copolymer

Stearic Acid

Cetyl Phosphate

Glycerin

Benzyl Alcohol

Cetyl Alcohol  
 Carbomer 1342  
 Triethanolamine

**Warnings**

For external use only. Do not use on damaged or broken skin. Keep out of eyes. Rinse eyes thoroughly with water to remove. Stop use and ask a physician if rash or irritation develops and lasts. Store away from excessive heat and direct sun.

**Package Label; Principal Display Panel**

**Drug Facts**

Active Ingredient	Purpose
Octocrylene 10.0%	Sunscreen
Octinoxate 7.5%	Sunscreen
Oxybenzone 6.0%	Sunscreen
Avobenzone 1.0%	Sunscreen

**Uses** ■ helps prevent sunburn ■ if used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer and early skin aging caused by the sun

**Warnings For external use only**

**Do not use** ■ on damaged or broken skin.

**When using this product**  
 ■ keep out of eyes. Rinse with water to remove.

**Stop use and ask a doctor if** ■ if rash occurs.

**Keep out of reach of children** if swallowed, get medical help or contact Poison Control Center right away.

**Directions** ■ Apply liberally and evenly 15 minutes before sun exposure and massage in gently, allowing it to dry. ■ reapply to dry skin: ■ after 80 minutes of swimming or sweating ■ immediately after towel drying ■ at least every 2 hours. ■ children under 6 months of age: Ask a doctor.

**Sun Protection Measures.** ■ Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad-spectrum SPF of 15 or higher and other sun protection measures including ■ Limit time in the sun, especially from 10 a.m. - 2 p.m. ■ Wear long-sleeve shirts, pants, hats, and sunglasses. ■ Children under 6 months: Ask a doctor.

**Other Information** ■ store away from excessive heat or cold.

**Inactive Ingredients**  
 Water (Aqua), Propylene Glycol Dicaprylate/Dicaprate, Dimethicone, Stearic Acid, PVP (and) VP/Eicosene Copolymer, Cetyl Phosphate, Glycerin, Benzyl Alcohol, Cetyl Alcohol, Sodium Hydroxide, Acrylates/C10-30 Alkyl Acrylate Crosspolymer, Citric Acid

**Questions? Please call 1 (800) 423-2341**

PC Person & Covey, Inc. Glendale, California 91201, U.S.A. **MADE IN USA** 8T  
 personandcovey.com

**FRAGRANCE FREE** Natural and healing formula for broad spectrum UV protection

<b>SOLBAR FIFTY SPF50</b>			
solbar fifty spf50 cream			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0096-0741
<b>Route of Administration</b>	TOPICAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>OCTOCRYLENE</b> (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)	OCTOCRYLENE	0.1 g in 1 g
<b>OCTINOXATE</b> (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	0.08 g in 1 g
<b>OXYBENZONE</b> (UNII: 95OOS7VE0Y) (OXYBENZONE - UNII:95OOS7VE0Y)	OXYBENZONE	0.06 g in 1 g
<b>AVOBENZONE</b> (UNII: G63QQF2NOX) (AVOBENZONE - UNII:G63QQF2NOX)	AVOBENZONE	0.011 g in 1 g

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>PROPYLENE GLYCOL DIETHYLHEXANOATE</b> (UNII: 8D8I9Z0F1Z)	
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)	
<b>EICOSYL POVIDONE (2 EICOSYL BRANCHES/REPEAT)</b> (UNII: XQQ9MKE2BJ)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>CETYL PHOSPHATE</b> (UNII: VT07D6X67O)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>BENZYL ALCOHOL</b> (UNII: LKG8494WBH)	
<b>CETYL ALCOHOL</b> (UNII: 936JST6JCN)	
<b>CARBOMER 1342</b> (UNII: 809Y72KV36)	
<b>TROLAMINE</b> (UNII: 9O3K93S3TK)	

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:0096-0741-04	128 g in 1 BOTTLE; Type 0: Not a Combination Product	06/01/1996	

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M020	06/01/1996	

**Labeler** - Person and Covey (008482473)**Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
Person and Covey		008482473	manufacture(0096-0741)

Revised: 12/2025

Person and Covey