

TETRACYCLINE HYDROCHLORIDE- tetracycline hydrochloride capsule
Preferred Pharmaceuticals Inc.

TETRACYCLINE HYDROCHLORIDE CAPSULES USP
For Oral Use

Rx only

To reduce the development of drug-resistant bacteria and maintain the effectiveness of tetracycline hydrochloride and other antibacterial drugs, tetracycline hydrochloride should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria.

DESCRIPTION

Tetracycline, USP is a yellow, odorless, crystalline powder. Tetracycline is stable in air but exposure to strong sunlight causes it to darken. Its potency is affected in solutions of pH below 2 and is rapidly destroyed by alkali hydroxide solutions. Tetracycline, USP is very slightly soluble in water, freely soluble in dilute acid and in alkali hydroxide solutions, sparingly soluble in alcohol, and practically insoluble in chloroform and in ether. The chemical name for tetracycline hydrochloride is 4-(Dimethylamino)-1,4,4a,5,5a,6,11,12a-octahydro-3,6,10,12,-12a-pentahydroxy-6-methyl-1,11-dioxo-2-naphthacenecarboxamide monohydrochloride.

Each capsule, for oral administration, contains tetracycline hydrochloride USP, 250 mg or 500 mg.

Inactive Ingredients: Lactose, magnesium stearate, and sodium lauryl sulfate.



The 250 mg capsule shell contains D&C yellow no. 10, FD&C yellow no. 6, gelatin, sodium lauryl sulfate, and titanium dioxide. It may also contain benzyl alcohol, butylparaben, D&C red no. 22, edetate calcium disodium, methylparaben, propylparaben, silicon dioxide, and sodium propionate.

The imprinting ink for the 250 mg capsule contains pharmaceutical glaze, and synthetic black iron oxide. It may also contain D&C yellow no. 10 (aluminum lake), dimethylpolysiloxane, distilled water, ethylene glycol monoethyl ether, FD&C blue no. 1 (aluminum lake), FD&C blue no. 2 (aluminum lake), FD&C red no. 40 (aluminum lake), lecithin, n-butyl alcohol, propylene alcohol, and SDA-3A alcohol.

The 500 mg capsule shell contains D&C yellow no. 10, FD&C blue no.1, FD&C red no. 40, gelatin, sodium lauryl sulfate, and titanium dioxide. It may also contain benzyl alcohol, butylparaben, edetate calcium disodium, FD&C yellow no. 6, methylparaben, propylparaben, silicon dioxide, and sodium propionate.

The imprinting ink for the 500 mg capsule contains titanium dioxide. It may also contain dimethyl polysiloxane, distilled water, ethyl alcohol, ethylene glycol monoethyl ether, pharmaceutical glaze, pharmaceutical shellac, and soya lecithin.

Its structural formula is as follows:



CLINICAL PHARMACOLOGY

Tetracyclines are readily absorbed and are bound to plasma protein in varying degrees. They are concentrated by the liver in the bile and excreted in the urine and feces at high concentrations in a biologically active form.

Microbiology

Tetracyclines are primarily bacteriostatic and exert their antimicrobial effect by the inhibition of protein synthesis by binding to the 30S ribosomal subunit. Tetracycline is active against a broad range of gram-negative and gram-positive organisms. The drugs in the tetracycline class have closely similar antimicrobial spectra, and cross-resistance among them is common.

Tetracycline has been shown to be active against most isolates of the following bacteria, both *in vitro* and in clinical infections as described in the **INDICATIONS AND USAGE** section of the package insert.

Gram-negative Bacteria

Acinetobacter species
Bartonella bacilliformis
Brucella species
Campylobacter fetus
Enterobacter aerogenes
Escherichia coli
Francisella tularensis
Haemophilus ducreyi
Haemophilus influenzae
Klebsiella species
Klebsiella granulomatis
Neisseria gonorrhoeae
Shigella species
Vibrio cholera
Yersinia pestis

Gram-positive Bacteria

Bacillus anthracis
Streptococcus pyogenes
Streptococcus pneumoniae
Staphylococcus aureus

Listeria monocytogenes

Anaerobes

Bacteroides species

Clostridium species

Fusobacterium fusiforme

Propionibacterium acnes

Other Bacteria

Actinomyces species

Borrelia recurrentis

Chlamydophila psittaci

Chlamydia trachomatis

Rickettsiae

Treponema pallidum *Treponema pallidum* subspecies *pertenue*

Parasites

Entamoeba species

Balantidium coli

Susceptibility Testing

For specific information regarding susceptibility test interpretive criteria and associated test methods and quality control standards recognized by FDA for this drug, please see: <https://www.fda.gov/STIC>.

INDICATIONS AND USAGE

To reduce the development of drug-resistant bacteria and maintain the effectiveness of tetracycline hydrochloride capsules and other antibacterial drugs, tetracycline hydrochloride capsules should be used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

Tetracycline is indicated in the treatment of infections caused by susceptible strains of the designated organisms in the conditions listed below:

- Upper respiratory tract infections caused by *Streptococcus pyogenes*, *Streptococcus pneumoniae* and *Haemophilus influenzae*. Note: Tetracycline should not be used for streptococcal disease unless the organism has been demonstrated to be susceptible.
- Lower respiratory tract infections caused by *Streptococcus pyogenes*, *Streptococcus pneumoniae*, *Mycoplasma pneumoniae* (Eaton agent, and *Klebsiella* sp.)
- Skin and soft tissue infections caused by *Streptococcus pyogenes*, *Staphylococcus aureus*.
(Tetracyclines are not the drugs of choice in the treatment of any type of staphylococcal infections.)

- Infections caused by rickettsia including Rocky Mountain spotted fever, typhus group infections, Q fever, rickettsialpox.
- Psittacosis caused by *Chlamydophila psittaci*.
- Infections caused by *Chlamydia trachomatis* such as uncomplicated urethral, endocervical or rectal infections, inclusion conjunctivitis, trachoma, and lymphogranuloma venereum.
- Granuloma inguinale caused by *Klebsiella granulomatis*.
- Relapsing fever caused by *Borrelia* sp.
- Bartonellosis caused by *Bartonella bacilliformis*.
- Chancroid caused by *Haemophilus ducreyi*.
- Tularemia caused by *Francisella tularensis*.
- Plague caused by *Yersinia pestis*.
- Cholera caused by *Vibrio cholerae*.
- Brucellosis caused by *Brucella* species (tetracycline may be used in conjunction with an aminoglycoside).
- Infections due to *Campylobacter fetus*.
- As adjunctive therapy in intestinal amebiasis caused by *Entamoeba histolytica*.
- Urinary tract infections caused by susceptible strains of *Escherichia coli*, *Klebsiella*, etc.
- Other infections caused by susceptible gram-negative organisms such as *E. coli*, *Enterobacter aerogenes*, *Shigella* sp., *Acinetobacter* sp., *Klebsiella* sp., and *Bacteroides* sp.
- In severe acne, adjunctive therapy with tetracycline may be useful.

When penicillin is contraindicated, tetracyclines are alternative drugs in the treatment of the following infections:

- Syphilis and yaws caused by *Treponema pallidum* and *pertenue*, respectively,
- Vincent's infection caused by *Fusobacterium fusiforme*,
- Infections caused by *Neisseria gonorrhoeae*,
- Anthrax caused by *Bacillus anthracis*,
- Infections due to *Listeria monocytogenes*,
- Actinomycosis caused by *Actinomyces* species,
- Infections due to *Clostridium* species.

CONTRAINDICATIONS

This drug is contraindicated in persons who have shown hypersensitivity to any of the tetracyclines.

WARNINGS

Tooth Development

The use of drugs of the tetracycline-class during tooth development (last half of pregnancy, infancy and

childhood to the age of 8 years) may cause permanent discoloration of the teeth (yellow-gray-brown).

This adverse reaction is more common during long-term use of the drugs but it has

been observed

following repeated short-term courses. Enamel hypoplasia has also been reported. Tetracycline drugs should not be used in this age group, except for anthrax, unless other drugs are not likely to be effective or are contraindicated.

***Clostridium difficile* Associated Diarrhea**

Clostridium difficile associated diarrhea (CDAD) has been reported with use of nearly all antibacterial

agents, including tetracyclines, and may range in severity from mild diarrhea to fatal colitis. Treatment

with antibacterial agents alters the normal flora of the colon leading to overgrowth of *C. difficile*.

C. difficile produces toxins A and B which contribute to the development of CDAD. Hypertoxin producing strains of *C. difficile* cause increased morbidity and mortality, as these infections can be refractory to antimicrobial therapy and may require colectomy. CDAD must be considered in all patients who present with diarrhea following antibacterial drugs. Careful medical history is necessary since CDAD has been reported to occur over two months after the administration of antibacterial agents.

If CDAD is suspected or confirmed, ongoing use of antibacterial drugs not directed against *C. difficile* need to be discontinued. Appropriate fluid and electrolyte management, protein supplementation, antibacterial treatment of *C. difficile*, and institute surgical evaluation clinically indicated.

Photosensitivity

Photosensitivity manifested by an exaggerated sunburn reaction has been observed in some individuals taking tetracyclines. Advise patients apt to be exposed to direct sunlight or ultraviolet lights that this reaction can occur with tetracycline drugs. Discontinue treatment at the first evidence of skin erythema.

Intracranial Hypertension

Intracranial hypertension (IH, pseudotumor cerebri) has been associated with the use of tetracyclines including Achromycin V. Clinical manifestations of IH include headache, blurred vision, diplopia, and vision loss; papilledema can be found on fundoscopy. Women of childbearing age who are overweight or have a history of IH are at greater risk for developing tetracycline associated IH. Concomitant use of isotretinoin and tetracycline should be avoided because isotretinoin, a systemic retinoid, is also known to cause pseudotumor cerebri.

Although IH typically resolve after discontinuation of treatment, the possibility for permanent visual loss exists. If visual disturbance occurs during treatment, prompt ophthalmologic evaluation is warranted. Since intracranial pressure can remain elevated for weeks after drug cessation patients should be monitored until they stabilize.

Skeletal Development

All tetracyclines form a stable calcium complex in any bone forming tissue. A decrease in fibula growth rate has been observed in premature infants given oral tetracycline in

doses of 25 mg/kg every six hours. This reaction was shown to be reversible when the drug was discontinued.

Results of animal studies indicate that tetracyclines cross the placenta, are found in fetal tissues and can have toxic effects on the developing fetus (often related to retardation of skeletal development). Evidence of embryotoxicity has also been noted in animals treated early in pregnancy. If this drug is used during pregnancy or if the patient becomes pregnant while taking this drug, apprise the patient of the potential hazard to the fetus. Tetracycline drugs should not be used during pregnancy unless absolutely necessary.

Antianabolic Action

The antianabolic action of the tetracyclines may cause an increase in BUN. While this is not a problem in those with normal renal function, in patients with significantly impaired renal function, higher serum levels of tetracycline may lead to azotemia, hyperphosphatemia and acidosis.

Laboratory Monitoring for Long-Term Therapy

In long-term therapy, perform periodic laboratory evaluation of organ systems, including hematopoietic, renal and hepatic studies. If renal impairment exists, even usual oral or parenteral doses may lead to excessive systemic accumulation of the drug and possible liver toxicity. Under such conditions, lower than usual total doses are indicated, and, if therapy is prolonged, serum level determinations of the drug may be advisable.

PRECAUTIONS

General

As with other antibacterials, use of this drug may result in overgrowth of nonsusceptible organisms, including fungi. If superinfection occurs, discontinue antibacterial and institute appropriate therapy.

Treat all infections due to Group A beta-hemolytic streptococci for at least ten days.

Perform incision and drainage or other surgical procedures in conjunction with antibacterial therapy, when indicated.

Prescribing tetracycline in the absence of proven or strongly suspected bacterial infection or a prophylactic indication is unlikely to provide benefit to the patient and increases the risk of the development of drug-resistant bacteria.

Information for Patients

Counsel patients that antibacterial drugs including tetracycline should only be used to treat bacterial infections. They do not treat viral infections (e.g., the common cold). When tetracycline is prescribed to treat a bacterial infection, tell patients that although it is common to feel better early in the course of therapy, the medication should be taken exactly as directed. Skipping doses or not completing the full course of therapy may (1) decrease the effectiveness of the immediate treatment and (2) increase the likelihood that bacteria will develop resistance and will not be treatable by tetracycline or other antibacterial drugs in the future.

Laboratory Tests

In sexually transmitted infections, when coexistent syphilis is suspected, perform dark field examinations before treatment is started and the blood serology repeated monthly for at least four months.

Drug Interactions

Since bacteriostatic drugs may interfere with the bactericidal action of penicillin, it is advisable to avoid giving tetracycline in conjunction with penicillin or other bactericidal antibacterials.

Because the tetracyclines have been shown to depress plasma prothrombin activity, patients who are on anticoagulant therapy may require downward adjustment of their anticoagulant dosage.

The concurrent use of tetracycline and methoxyflurane has been reported to result in fatal renal toxicity.

Absorption of tetracyclines is impaired by antacids containing aluminum, calcium or magnesium and preparations containing iron, zinc, or sodium bicarbonate.

Concurrent use of tetracycline may render oral contraceptives less effective.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term animal studies are currently being conducted to determine whether tetracycline hydrochloride has carcinogenic potential. Some related antibacterials (oxytetracycline, minocycline) have shown evidence of oncogenic activity in rats.

In two *in vitro* mammalian cell assay systems (L 51784y mouse lymphoma and Chinese hamster lung cells), there was evidence of mutagenicity with tetracycline hydrochloride.

Tetracycline hydrochloride had no effect on fertility when administered in the diet to male and female rats at a daily intake of approximately 400 mg/kg/day, roughly 8 times the highest recommended human dose based on body surface area.

Pregnancy

Teratogenic Effects

Pregnancy Category D
(see)

Nonteratogenic Effects

(see)

Pregnant women with renal disease may be more prone to develop tetracycline-associated liver failure.

Labor and Delivery

The effect of tetracyclines on labor and delivery is unknown.

Nursing Mothers

Because of the potential for serious adverse reaction in nursing infants from tetracyclines, a decision should be made whether to discontinue the drug, taking into account the importance of the drug to the mother (see).

Pediatric Use

See and .

ADVERSE REACTIONS

Gastrointestinal:

anorexia, nausea, epigastric distress, vomiting, diarrhea, glossitis, black hairy tongue, dysphagia, enterocolitis, and inflammatory lesions (with *Candida* overgrowth) in the anogenital region.

Esophagitis and esophageal ulceration have been reported in patients receiving particularly the capsule and also the tablet forms of tetracyclines. Most of the patients were reported to have taken medication immediately before going to bed (see).

Teeth:

permanent discoloration of teeth may be caused during tooth development. Enamel hypoplasia has been reported (see).

Skin:

maculopapular and erythematous rashes. Exfoliative dermatitis has been reported. Onycholysis and discoloration of the nails have been reported. Photosensitivity is discussed in .

Renal Toxicity:

an increase in BUN has been reported and is dose related.

Liver:

hepatotoxicity and liver failure have been observed in patients receiving tetracycline and in tetracycline-treated patients with renal impairment.

Hypersensitivity Reactions:

urticaria, angioneurotic edema, anaphylaxis, anaphylactoid purpura, pericarditis, exacerbation of systemic lupus erythematosus, and serum sickness-like reactions, as fever, rash, and arthralgia.

Blood:

hemolytic anemia, thrombocytopenia, thrombocytopenic purpura, neutropenia and eosinophilia have been reported.

When given over prolonged periods, tetracyclines have been reported to produce brown-black microscopic discoloration of thyroid glands. No abnormalities of thyroid function studies are known to occur.

To report SUSPECTED ADVERSE EVENTS, contact Actavis at 1-800-272-5525 or FDA at 1-800-FDA-1088 or <http://www.fda.gov/> for voluntary reporting of adverse reactions.

OVERDOSAGE

In case of overdosage, discontinue medication, treat symptomatically and institute supportive measures. Tetracycline is not dialyzable.

DOSAGE AND ADMINISTRATION

Adults: Usual daily dose, 1 gram as 500 mg twice a day or 250 mg four times a day. Higher doses such as 500 mg four times a day may be required for severe infections or for those infections which do not respond to the smaller doses.

For pediatric patients above eight years of age: Usual daily dose, 10 mg/lb to 20 mg/lb (25 mg/kg to 50 mg/kg) body weight divided in four equal doses.

Administration of adequate amounts of fluid with the capsule formulation of tetracycline is recommended to wash down the drug and reduce the risk of esophageal irritation and ulceration (see **ADVERSE REACTIONS**).

Absorption of tetracycline is impaired by antacids containing aluminum, calcium or magnesium and preparations containing iron, zinc or sodium bicarbonate. Food and some dairy products also interfere with absorption.

When used in streptococcal infections, therapy should be continued for 10 days.

For treatment of brucellosis, 500 mg tetracycline four times a day for three weeks accompanied by streptomycin, 1 gram intramuscularly twice daily the first week and once daily the second week.

For the treatment of syphilis in patients allergic to penicillin, the following dosage of tetracycline is recommended: early syphilis (less than one year's duration), 500 mg four times a day for 15 days. Syphilis of more than one year's duration (except neurosyphilis), 500 mg four times a day for 30 days.

For treatment of gonorrhea, the recommended dose is 500 mg by mouth four times a day for seven days.

Uncomplicated urethral, endocervical or rectal infections in adults caused by *Chlamydia trachomatis*: 500 mg, by mouth, four times a day for at least seven days.

In cases of moderate to severe acne which, in the judgement of the clinician, require long-term treatment, the recommended initial dosage is 1 gram daily in divided doses. When improvement is noted, reduce dosage gradually to maintenance levels ranging from 125 mg to 500 mg daily. In some patients it may be possible to maintain adequate remission of lesions with alternate day or intermittent therapy. Tetracycline therapy of acne should augment the other standard measures known to be of value. Duration of long-term treatment which can safely be recommended has not been established (see **WARNINGS** and **Carcinogenesis, Mutagenesis, Impairment of Fertility**).

Use in Specific Population

In patients with renal impairment (see **WARNINGS**): decrease total dosage by reduction of recommended individual doses and/or by extending time intervals between doses.

HOW SUPPLIED

Tetracycline Hydrochloride Capsules USP are available as:

500 mg: Black/yellow capsules, imprinted with “**WPI**” on cap and “**2235**” on body.
Available in bottles of 30 (NDC 68788-8853-3).

Dispense in a tight, light-resistant container as defined in the USP, with a child-resistant closure (as required).

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

ANIMAL PHARMACOLOGY AND ANIMAL TOXICOLOGY

Hyperpigmentation of the thyroid has been produced by members of the tetracycline class in the following species: in rats by oxytetracycline, doxycycline, minocycline, tetracycline PO₄ and methacycline; in minipigs by doxycycline, minocycline, tetracycline PO₄ and methacycline; in dogs by doxycycline and minocycline; in monkeys by minocycline. Minocycline, tetracycline PO₄, methacycline, doxycycline, tetracycline base, oxytetracycline HCl and tetracycline HCl were goitrogenic in rats fed a low iodine diet. This goitrogenic effect was accomplished by high radioactive iodine uptake. Administration of minocycline also produced a large goiter with high radioiodine uptake in rats fed a relatively high iodine diet. Treatment of various animal species with this class of drugs has also resulted in the induction of thyroid hyperplasia in the following: in rats and dogs (minocycline), in chickens (chlortetracycline) and in rats and mice (oxytetracycline). Adrenal gland hyperplasia has been observed in goats and rats treated with oxytetracycline.

Manufactured by:
Watson Pharma Private Limited
Verna, Salcette Goa 403 722 INDIA

Distributed by:
Actavis Pharma, Inc.
Parsippany, NJ 07054 USA

Rev. A 11/2018

Repackaged By: Preferred Pharmaceuticals Inc.

PRINCIPAL DISPLAY PANEL

NDC 68788-8853-3

Rx Only

Tetracycline HCl Capsules USP

500mg

Generic for Achromycin

Each Capsule Contains: Tetracycline HCL USP 500mg

Pkg Size: Exp Date: #####

Ins:
Mfg: Watson Pharma Private Limited


Prod#:

Warning
Store at 20°- 25°C (68°- 77°F) See USP Controlled Room Temperature, Rx Only. Keep this and all medication out of the reach of children. Capsules are black and yellow, imprinted with WPI 2235

PREFERRED
Pharmaceuticals, Inc.


CAUTION: Federal law PROHIBITS transfer of this drug to any person other than the patient for whom it was prescribed.

Tetracycline HCl Capsules USP
500mg
Qty: Ins:
Lot: Bat:
Prod# (NDC):



Directions English

Take ____ capsule(s)
every ____ hours.



GTIN

SN #####
EXP #####

Tetracycline HCl Capsules USP
500mg
Qty: Ins:
Lot: Bat:
Prod# (NDC):

Instrucciones Espanol:

Toma ____ capsula(s)
cada ____ horas.

Tetracycline HCl Capsules USP
500mg
Qty: Ins:
Lot: Bat:
Prod# (NDC):

Tetracycline HCl Capsules USP
500mg
Qty: Ins:
Lot: Bat:
Prod# (NDC):

Ins:
Mfg: Watson Pharma Private Limited
Prod#:
Warning
Store at 20°- 25°C (68°- 77°F) See USP Controlled Room Temperature, Rx Only. Keep this and all medication out of the reach of children. Capsules are black and yellow, imprinted with WPI 2235

CAUTION: Federal law PROHIBITS transfer of this drug to any person other than the patient for whom it was prescribed.

Tetracycline HCl Capsules USP
500mg
Qty: Ins:
Lot: Bat:
Prod# (NDC):

Directions English

Take ____ capsule(s)
every ____ hours.

GTIN

SN #####
EXP #####

Tetracycline HCl Capsules USP
500mg
Qty: Ins:
Lot: Bat:
Prod# (NDC):

Instrucciones Espanol:

Toma ____ capsula(s)
cada ____ horas.

Tetracycline HCl Capsules USP
500mg
Qty: Ins:
Lot: Bat:
Prod# (NDC):

Tetracycline HCl Capsules USP
500mg
Qty: Ins:
Lot: Bat:
Prod# (NDC):

Ins:
Mfg: Watson Pharma Private Limited
Prod#:
Warning
Store at 20°- 25°C (68°- 77°F) See USP Controlled Room Temperature, Rx Only. Keep this and all medication out of the reach of children. Capsules are black and yellow, imprinted with WPI 2235

CAUTION: Federal law PROHIBITS transfer of this drug to any person other than the patient for whom it was prescribed.

Tetracycline HCl Capsules USP
500mg
Qty: Ins:
Lot: Bat:
Prod# (NDC):

Directions English

Take ____ capsule(s)
every ____ hours.

GTIN

SN #####
EXP #####

Tetracycline HCl Capsules USP
500mg
Qty: Ins:
Lot: Bat:
Prod# (NDC):

Instrucciones Espanol:

Toma ____ capsula(s)
cada ____ horas.

Tetracycline HCl Capsules USP
500mg
Qty: Ins:
Lot: Bat:
Prod# (NDC):

Tetracycline HCl Capsules USP
500mg
Qty: Ins:
Lot: Bat:
Prod# (NDC):

Ins:
Mfg: Watson Pharma Private Limited
Prod#:
Warning
Store at 20°- 25°C (68°- 77°F) See USP Controlled Room Temperature, Rx Only. Keep this and all medication out of the reach of children. Capsules are black and yellow, imprinted with WPI 2235

CAUTION: Federal law PROHIBITS transfer of this drug to any person other than the patient for whom it was prescribed.

Tetracycline HCl Capsules USP
500mg
Qty: Ins:
Lot: Bat:
Prod# (NDC):

Directions English

Take ____ capsule(s)
every ____ hours.

GTIN

SN #####
EXP #####

Tetracycline HCl Capsules USP
500mg
Qty: Ins:
Lot: Bat:
Prod# (NDC):

Instrucciones Espanol:

Toma ____ capsula(s)
cada ____ horas.

Tetracycline HCl Capsules USP
500mg
Qty: Ins:
Lot: Bat:
Prod# (NDC):

Tetracycline HCl Capsules USP
500mg
Qty: Ins:
Lot: Bat:
Prod# (NDC):

Ins:
Mfg: Watson Pharma Private Limited
Prod#:
Warning
Store at 20°- 25°C (68°- 77°F) See USP Controlled Room Temperature, Rx Only. Keep this and all medication out of the reach of children. Capsules are black and yellow, imprinted with WPI 2235

CAUTION: Federal law PROHIBITS transfer of this drug to any person other than the patient for whom it was prescribed.

Tetracycline HCl Capsules USP
500mg
Qty: Ins:
Lot: Bat:
Prod# (NDC):

Directions English

Take ____ capsule(s)
every ____ hours.

GTIN

SN #####
EXP #####

Tetracycline HCl Capsules USP
500mg
Qty: Ins:
Lot: Bat:
Prod# (NDC):

Instrucciones Espanol:

Toma ____ capsula(s)
cada ____ horas.

Tetracycline HCl Capsules USP
500mg
Qty: Ins:
Lot: Bat:
Prod# (NDC):

Tetracycline HCl Capsules USP
500mg
Qty: Ins:
Lot: Bat:
Prod# (NDC):

Ins:
Mfg: Watson Pharma Private Limited
Prod#:
Warning
Store at 20°- 25°C (68°- 77°F) See USP Controlled Room Temperature, Rx Only. Keep this and all medication out of the reach of children. Capsules are black and yellow, imprinted with WPI 2235

CAUTION: Federal law PROHIBITS transfer of this drug to any person other than the patient for whom it was prescribed.

Tetracycline HCl Capsules USP
500mg
Qty: Ins:
Lot: Bat:
Prod# (NDC):

Directions English

Take ____ capsule(s)
every ____ hours.

GTIN

SN #####
EXP #####

Tetracycline HCl Capsules USP
500mg
Qty: Ins:
Lot: Bat:
Prod# (NDC):

Instrucciones Espanol:

Toma ____ capsula(s)
cada ____ horas.

Tetracycline HCl Capsules USP
500mg
Qty: Ins:
Lot: Bat:
Prod# (NDC):

Tetracycline HCl Capsules USP
500mg
Qty: Ins:
Lot: Bat:
Prod# (NDC):

Ins:
Mfg: Watson Pharma Private Limited
Prod#:
Warning
Store at 20°- 25°C (68°- 77°F) See USP Controlled Room Temperature, Rx Only. Keep this and all medication out of the reach of children. Capsules are black and yellow, imprinted with WPI 2235

CAUTION: Federal law PROHIBITS transfer of this drug to any person other than the patient for whom it was prescribed.

Tetracycline HCl Capsules USP
500mg
Qty: Ins:
Lot: Bat:
Prod# (NDC):

Directions English

Take ____ capsule(s)
every ____ hours.

GTIN

SN #####
EXP #####

Tetracycline HCl Capsules USP
500mg
Qty: Ins:
Lot: Bat:
Prod# (NDC):

TETRACYCLINE HYDROCHLORIDE

tetracycline hydrochloride capsule

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:68788-8853(NDC:0591-2475)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TETRACYCLINE HYDROCHLORIDE (UNII: P6R62377KV) (TETRACYCLINE - UNII:F8VB5M810T)	TETRACYCLINE HYDROCHLORIDE	500 mg

Inactive Ingredients

Ingredient Name	Strength
LACTOSE, UNSPECIFIED FORM (UNII: J2B2A4N98G)	
MAGNESIUM STEARATE (UNII: 70097M6I3O)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
BENZYL ALCOHOL (UNII: LKG8494WBH)	
BUTYLPARABEN (UNII: 3QPI1U3FV8)	
EDETATE CALCIUM DISODIUM (UNII: 25IH6R4SGF)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM PROPIONATE (UNII: DK6Y9P42IN)	
DIMETHICONE, UNSPECIFIED (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
ALCOHOL (UNII: 3K9958V90M)	
ETHYLENE GLYCOL MONOETHYL ETHER (UNII: IDK7C2HS09)	
SHELLAC (UNII: 46N107B71O)	
SOYBEAN LECITHIN (UNII: 1DI56QDM62)	

Product Characteristics

Color	black, yellow	Score	no score
Shape	CAPSULE	Size	22mm
Flavor		Imprint Code	WPI;2235
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68788-8853-3	30 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/24/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA061837	03/24/2025	

Labeler - Preferred Pharmaceuticals Inc. (791119022)

Registrant - Preferred Pharmaceuticals Inc. (791119022)

Establishment

Name	Address	ID/FEI	Business Operations
Preferred Pharmaceuticals Inc.		791119022	REPACK(68788-8853)