

**ACID CONTROLLER- famotidine tablet, film coated**  
**Walgreen Company**

-----  
**Walgreen Co. Acid Controller Drug Facts**

**Active ingredient (in each tablet)**

Famotidine 20 mg

**Purpose**

Acid reducer

**Uses**

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

**Warnings**

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- if you have kidney disease, except under the advice and supervision of a doctor
- with other acid reducers

**Ask a doctor before use if you have**

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating, or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

**Stop use and ask a doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- adults and children 12 years and over:
- to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

**Other information**

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°-25°C (68°-77°F)
- protect from moisture and light

**Inactive ingredients**

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, lactose (monohydrate), magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talc, and titanium dioxide

**Questions or comments?**

1-800-719-9260

**Principal Display Panel**

Compare to Maximum Strength Pepcid® AC active ingredient

Acid Controller

FAMOTIDINE TABLETS, 20 mg / ACID REDUCER

MAXIMUM STRENGTH

Just one tablet prevents & relieves heartburn due to acid indigestion

25 TABLETS

ACTUAL SIZE

*Walgreens*

# Acid Controller

FAMOTIDINE TABLETS, 20 mg /  
ACID REDUCER

ORF00318-F

*Walgreens*

Compare to Maximum Strength  
Pepcid® AC active ingredient†

NDC 0363-0701-02

# Acid Controller

FAMOTIDINE TABLETS, 20 mg /  
ACID REDUCER

**MAXIMUM STRENGTH**

- Just one tablet prevents & relieves  
heartburn due to acid indigestion

**25** TABLETS



ACTUAL SIZE

**DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN**

: 19447 94 55

**Drug Facts**

Active ingredient (in each tablet)	Purpose
Famotidine 20 mg.....	Acid reducer

**Uses**

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

**Warnings**

**Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- if you have kidney disease, except under the advice and supervision of a doctor
- with other acid reducers

**Ask a doctor before use if you have**

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**

**Drug Facts (continued)**

- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

**Stop use and ask a doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days

**If pregnant or breast-feeding**, ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- adults and children 12 years and over:
  - to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
  - to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **10 to 60 minutes before** eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

**Other information**

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°-25°C (68°-77° F)
- protect from moisture and light

**Drug Facts (continued)**

**Inactive ingredients** carnauba wax, colloidal silicon dioxide, croscarmellose sodium, lactose (monohydrate), magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talc, and titanium dioxide

**Questions or comments?** 1-800-719-9260

Walgreens Pharmacist Recommended  
Walgreens Pharmacist Survey  
† This product is not manufactured or distributed by Johnson & Johnson - MERCK Consumer Pharmaceuticals Co., distributor of Maximum Strength Pepcid® AC.  
Gluten Free

DISTRIBUTED BY: WALGREEN CO.  
200 WILMOT RD., DEERFIELD, IL 60015  
**100% SATISFACTION GUARANTEE D**  
walgreens.com ©2018Walgreen Co.  
ITEM 587087 W10106-0618-L

3 11917 07592 1

OPEN OTHER END

**Tips for Managing Heartburn**

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed

CONVENIENT RECYCLING TAB

<b>ACID CONTROLLER</b>			
famotidine tablet, film coated			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0363-0701
<b>Route of Administration</b>	ORAL		
<b>Active Ingredient/Active Moiety</b>			
<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>	
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	20 mg	
<b>Inactive Ingredients</b>			
<b>Ingredient Name</b>	<b>Strength</b>		
CARNAUBA WAX (UNII: R12CBM0EIZ)			
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)			
LACTOSE MONOHYDRATE (UNII: EWQ57Q815X)			

<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL</b> (UNII: 3WJQ0SDW1A)	
<b>POLYVINYL ALCOHOL</b> (UNII: 532B59J990)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	8mm
<b>Flavor</b>		<b>Imprint Code</b>	L194
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0701-71	1 in 1 CARTON	09/26/2006	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0363-0701-72	1 in 1 CARTON	06/02/2011	
2		60 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:0363-0701-01	1 in 1 CARTON	10/29/2007	
3		85 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:0363-0701-02	25 in 1 CARTON	09/26/2006	
4		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
5	NDC:0363-0701-39	30 in 1 CARTON	04/10/2015	04/10/2015
5		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
6	NDC:0363-0701-82	1 in 1 CARTON	05/08/2019	
6		200 in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA077351	09/26/2006	

**Labeler** - Walgreen Company (008965063)

Revised: 5/2019

Walgreen Company