

ASPIRIN 325 MG- aspirin tablet, coated
BETTER LIVING BRANDS, LLC

Albertsons Aspirin 325 mg tablets

Drug Facts

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves:
- headache
- muscle pain
- toothache
- menstrual pain
- pain and fever of colds
- minor pain of arthritis

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you

- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - redness or swelling is present
 - fever lasts more than 3 days
 - new symptoms occur
 - ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours
- children under 12 years: consult a doctor

Other information

- store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients corn starch, croscarmellose sodium, hypromellose, microcrystalline cellulose, mineral oil, titanium dioxide

Questions or comments? Call **1-877-290-4008**

MINERAL OIL (UNII: T5L8T28FGP)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
Product Characteristics				
Color	white	Score	no score	
Shape	ROUND	Size	10mm	
Flavor		Imprint Code	TCL011	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-758-53	500 in 1 BOTTLE; Type 0: Not a Combination Product	01/05/2026	
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013		01/05/2026	

Labeler - BETTER LIVING BRANDS, LLC (009137209)

Registrant - TIME CAP LABORATORIES, INC (037052099)

Establishment			
Name	Address	ID/FEI	Business Operations
Time Cap Laboratories, Inc.		037052099	manufacture(21130-758)