

PUVIDA WART REMOVAL SYSTEM- salicylic acid 17% wart removal system liquid

Jiangxi Hemei Pharmaceutical Co., Ltd

84010-210

Active Ingredient

Salicylic acid 17%

Purpose

Wart remover

Use

For the removal of common warts and plantar warts. Common warts have a rough "cauliflower-like" surface and are easily recognizable. Plantar warts are located only on the soles of the feet, tender to touch, with an interrupted footprint pattern.

Warnings

For external use only.

Do not use

on irritated skin or on any area that is infected or reddened. On moles, birthmarks, warts with hair growing from them, genital warts or warts on the face or mucous membranes. If you have diabetes or poor blood circulation.

.

When Using

Avoid contact with eyes. If product gets into the eye, flush with Water for 15 minutes. Non-edible. Cap tightly and store at room temperature, away from heat.

Stop Use

Stop use and ask a doctor if discomfort persists.

Ask Doctor

Stop use and ask a doctor if discomfort persists.

Keep Out Of Reach Of Children

If swallowed, get medical help or contact the Poison Control Center (1-800-222-1222) right away.

Directions

Wash the affected area. May soak wart in warm water for 5 minutes. Dry area thoroughly.

Use a cotton swab to apply a sufficient amount to cover each wart. Let it dry. Repeat this procedure once or twice daily as needed (until wart is removed) for up to 12 weeks.


Other information

Store at room temperature. Avoid excessive heat (37°C/ 99°F). Skin discoloration may occur during or after use.

Inactive ingredients

Water □ Polyethylene Glycol □ Glycerin □ Borneol.

PRINCIPAL DISPLAY PANEL





NET:0.085 oz fl (2.5ml)

liangxi Hemei
l Co., Ltd.
or, Building 1956,
ijiasan Medical
i, Zhangshu, Yichun,
China(CN)



6 9784603692716

KEEP OUT OF REACH OF CHILDREN

If swallowed, get medical help or contact the Poison Control Center (1-800-222-1222) right away.

Directions

Wash the affected area. May soak wart in warm water for 5 minutes. Dry area thoroughly. Use a cotton swab to apply a sufficient amount to cover each wart. Let it dry. Repeat this procedure once or twice daily as needed (until wart is removed) for up to 12 weeks.

Other information

Store at room temperature. Avoid excessive heat (37°C / 99°F). Skin discoloration may occur during or after use.

Inactive ingredients:

Water	53%
Polyethylene Glycol	16%
Glycerin	8%
Borneol	6%

PUVIDA®

PUVIDA®

WART REMOVAL SYSTEM

WART REMOVAL TREATMENT

PRECISION TIP FOR ACCURATE APPLICATION



- ✓Efficient Wart Removal Treatment
- ✓Targets Warts, Not Your Skin

PUVIDA®

Manufacturer: Jiangxi Hemei
Pharmaceutical Co., Ltd.
Address: 4th Floor, Building 1956,
Phase III, Zhangliashan Medical
Incubation Park, Zhangshu, Yichun
Jiangxi 331208, China(CN)



6 974603 691270

Drug Facts

Active ingredient:	Purpose
--------------------	---------

Salicylic acid 17%.....Wart remover

Uses

For the removal of common warts and plantar warts. Common warts have a rough "cauliflower-like" surface and are easily recognizable. Plantar warts are located only on the soles of the feet, tender to touch, with an interrupted footprint pattern.

Warnings

For external use only.

Do not use

on irritated skin or on any area that is infected or reddened. On moles, birthmarks, warts with hair growing from them, genital warts, or warts on the face or mucous membranes. If you have diabetes or poor blood circulation.

When using this product

Avoid contact with eyes.

If product gets into the eye, flush with Water for 15 minutes. Non-edible. Cap tightly and store at room temperature, away from heat.

Stop use and ask a doctor if discomfort persists.

Keep out of reach of children.

If swallowed, get medical help or contact the Poison Control Center (1-800-222-1222) right away.

Directions


Wash the affected area. May soak wart in warm water for 5 minutes. Dry area thoroughly. Use a cotton swab to apply a sufficient amount to cover each wart. Let it dry. Repeat this procedure once or twice daily as needed (until wart is removed) for up to 12 weeks.

Other information

Store at room temperature. Avoid excessive heat (37°C / 99°F). Skin discoloration may occur during or after use.

Inactive ingredients:

Water	53%
Polyethylene Glycol	16%

	0.135 oz fl (4ml)		Polyethylene Glycol 15% Glycerin 8% Borneol 6%

PUVIDA WART REMOVAL SYSTEM

salicylic acid 17% wart removal system liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:84010-210
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety		
Ingredient Name		Strength
SALICYLIC ACID (UNII: O414PZ4LPZ) (SALICYLIC ACID - UNII:O414PZ4LPZ)		17 g in 100 mL

Inactive Ingredients	
Ingredient Name	Strength
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
GLYCERIN (UNII: PDC6A3C0OX)	
BORNEOL (UNII: M89NIB437X)	
WATER (UNII: 059QF0KO0R)	

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84010-210-01	4 mL in 1 TUBE; Type 0: Not a Combination Product	10/30/2025	
2	NDC:84010-210-02	2.5 mL in 1 TUBE; Type 0: Not a Combination Product	10/30/2025	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M028	10/30/2025	

Labeler - Jiangxi Hemei Pharmaceutical Co., Ltd (724892056)

Establishment

Name	Address	ID/FEI	Business Operations
Jiangxi Hemei Pharmaceutical Co., Ltd		724892056	manufacture(84010-210)

Revised: 12/2025

Jiangxi Hemei Pharmaceutical Co., Ltd