

UQORA MAXIMUM UTI PAIN RELIEF- phenazopyridine hydrochloride tablet
Bonafide Health, LLC dba Uqora

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Uqora Maximum UTI Pain Relief

Active ingredient (in each tablet)

Phenazopyridine Hydrochloride 99.5 mg .

Purpose

Urinary Tract Analgesic

Use

Fast relief from urinary pain, burning, urgency and frequency associated with urinary tract infections. Treatment should not exceed 2 days; see Directions.

Warnings

Do not exceed recommended dosage

Do not use: if you have Glucose-6-Phosphate Dehydrogenase (G6PD) deficiency unless approved by your physician.

Ask doctor before use if you have

- kidney disease
- allergies to foods, preservatives or dyes
- had a hypersensitive reaction to phenazopyridine

When using this product

- stomach upset may occur, taking this product with or after meals may reduce stomach upset
- your urine will become reddish-orange in color. This is not harmful, but care should be taken to avoid staining clothing or other items.

■ Long-term administration of phenazopyridine hydrochloride has induced neoplasia in rats (large intestine) and mice (liver). Although no association between phenazopyridine hydrochloride and human neoplasia has been reported, adequate epidemiological studies along these lines have not been conducted.

Stop use and ask doctor if

- your symptoms last for more than 2 days

■ you suspect you are having an adverse reaction to the medication

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children. In case of an overdose, get medical help or contact a Poison Control Center right away 1-800-222-1222.

Directions

Adults and children 12 years and over:

■ take 2 tablets 3 times daily with a full glass of water, with or after meals as needed

■ **Do not use for more than 2 days (12 tablets) without consulting a doctor.**

Children under 12 years:

■ consult a doctor

Other information

■ this product may stain contact lenses ■ this product can interfere with laboratory tests including urine, glucose (sugar), and ketones tests ■ store at room temperature 15°-30°C (59°-86°F) in a dry place and protect from light ■ Tamper evident: tablets sealed in blisters. Do not use if blister foil or seal is open or damaged

Inactive ingredients

corn starch, croscarmellose sodium, hypromellose, lactose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinylpyrrolidone, pregelatinized starch, silicon dioxide, sodium starch glycolate, talc and triacetin

MAX STRENGTH MOST POWERFUL DOSE WITHOUT RX

uqora

FOUNDED BY A UTI SUFFERER

UTI PAIN RELIEF

MAX STRENGTH

FAST TARGETED RELIEF FOR

Burning

Pain & discomfort at the source unlike general pain relievers

Urgency

2 Day Supply

12 tablets

99.5 mg Phenazopyridine Hydrochloride

Not intended to replace medical care

WE'RE UQORA, THE URINARY HEALTH EXPERTS

SCIENCE-BACKED

UTI EDUCATION

A TEAM THAT CARES

SCAN TO LEARN MORE ABOUT UQORA





MAX STRENGTH MOST POWERFUL DOSE WITHOUT RX

UTI PAIN RELIEF

MAX STRENGTH

FAST TARGETED RELIEF FOR

- ✓ Burning
- ✓ Pain & discomfort at the source
- ✓ Urgency
- unlike general pain relievers

2 DAY SUPPLY

12 tablets 99.5 mg Phenazopyridine Hydrochloride

Not intended to replace medical care.

DISCOVER OUR URINARY TRACT DRINK MIXES



Dual-action **Flush Advanced** designed for women with recurrent urinary tract challenges



Flush, America's #1 urinary tract drink mix*

*Circana, LLC, Total US - Multi Outlet, Urinary Tract Drink Mix, Dollar Sales, 52 WE Apr 20, 2025

Drug Facts

Active ingredient (in each tablet) Phenazopyridine Hydrochloride 99.5 mg. **Purpose** Urinary Tract Analgesic

Use Fast relief from urinary pain, burning, urgency and frequency associated with urinary tract infections. Treatment should not exceed 2 days; see Directions

Warnings
Do not exceed recommended dosage
Do not use: if you have Glucose-6-Phosphate Dehydrogenase (G6PD) deficiency unless approved by your physician
Ask a doctor before use if you have: ■ kidney disease ■ allergies to foods, preservatives or dyes ■ had a hypersensitive reaction to phenazopyridine
When using this product: ■ stomach upset may occur, taking this product with or after meals may reduce stomach upset ■ your urine will become reddish-orange in color. This is not harmful, but care should be taken to avoid staining clothing or other items.
Stop use and ask a doctor if: ■ your symptoms last for more than 2 days ■ you suspect you are having an adverse reaction to the medication
■ Long-term administration of phenazopyridine hydrochloride has induced neoplasia in rats (large intestine) and mice (liver). Although no association between phenazopyridine hydrochloride and human neoplasia has been reported, adequate epidemiological studies along these lines have not been conducted.
If pregnant or breast feeding, ask a health professional before use.
Keep out of reach of children. In case of an overdose, get medical help or contact a Poison Control Center right away 1-800-222-1222.

Directions Adults and children 12 years and over: ■ take 2 tablets 3 times daily with a full glass of water, with or after meals as needed ■ Do not use for more than 2 days (12 tablets) without consulting a doctor.
Children under 12 years: ■ consult a doctor

Other information ■ this product may stain contact lenses ■ this product can interfere with laboratory tests including urine, glucose (sugar), and ketones tests ■ store at room temperature 15°-30°C (59°-86°F) in a dry place and protect from light ■ Tamper evident: tablets sealed in blisters. Do not use if blister foil or seal is open or damaged

Inactive ingredients corn starch, croscarmellose sodium, hypromellose, lactose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinylpyrrolidone, pregelatinized starch, silicon dioxide, sodium starch glycolate, talc and triacetin.

⚠ WARNING: This product can expose you to phenazopyridine hydrochloride, which is known to the state of California to cause cancer. For more information, visit www.P65Warnings.ca.gov

To report an adverse event call (888) 313-1372
© 2025 Uqora. Uqora is a trademark of Uqora.
Distributed by Uqora
4250 Executive Square, Suite 200, San Diego, CA 92037
support@uqora.com

WE'RE UQORA, THE URINARY HEALTH EXPERTS

SCIENCE-BACKED

UTI EDUCATION

A TEAM THAT CARES

SCAN TO LEARN MORE ABOUT UQORA





MAX STRENGTH MOST POWERFUL DOSE WITHOUT RX

UTI PAIN RELIEF

MAX STRENGTH

FAST TARGETED RELIEF FOR

- ✓ Burning
- ✓ Pain & discomfort at the source
- ✓ Urgency
- unlike general pain relievers

4 DAY SUPPLY

24 tablets 99.5 mg Phenazopyridine Hydrochloride

Not intended to replace medical care.

DISCOVER OUR URINARY TRACT DRINK MIXES



Dual-action **Flush Advanced** designed for women with recurrent urinary tract challenges



Flush, America's #1 urinary tract drink mix*

*Circana, LLC, Total US - Multi Outlet, Urinary Tract Drink Mix, Dollar Sales, 52 WE Apr 20, 2025

Drug Facts

Active ingredient (in each tablet) Phenazopyridine Hydrochloride 99.5 mg. **Purpose** Urinary Tract Analgesic

Use Fast relief from urinary pain, burning, urgency and frequency associated with urinary tract infections. Treatment should not exceed 2 days; see Directions

Warnings
Do not exceed recommended dosage
Do not use: if you have Glucose-6-Phosphate Dehydrogenase (G6PD) deficiency unless approved by your physician
Ask a doctor before use if you have: ■ kidney disease ■ allergies to foods, preservatives or dyes ■ had a hypersensitive reaction to phenazopyridine
When using this product: ■ stomach upset may occur, taking this product with or after meals may reduce stomach upset ■ your urine will become reddish-orange in color. This is not harmful, but care should be taken to avoid staining clothing or other items.
Stop use and ask a doctor if: ■ your symptoms last for more than 2 days ■ you suspect you are having an adverse reaction to the medication
■ Long-term administration of phenazopyridine hydrochloride has induced neoplasia in rats (large intestine) and mice (liver). Although no association between phenazopyridine hydrochloride and human neoplasia has been reported, adequate epidemiological studies along these lines have not been conducted.
If pregnant or breast feeding, ask a health professional before use.
Keep out of reach of children. In case of an overdose, get medical help or contact a Poison Control Center right away 1-800-222-1222.

Directions Adults and children 12 years and over: ■ take 2 tablets 3 times daily with a full glass of water, with or after meals as needed ■ Do not use for more than 2 days (12 tablets) without consulting a doctor.
Children under 12 years: ■ consult a doctor

Other information ■ this product may stain contact lenses ■ this product can interfere with laboratory tests including urine, glucose (sugar), and ketones tests ■ store at room temperature 15°-30°C (59°-86°F) in a dry place and protect from light ■ Tamper evident: tablets sealed in blisters. Do not use if blister foil or seal is open or damaged

Inactive ingredients corn starch, croscarmellose sodium, hypromellose, lactose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinylpyrrolidone, pregelatinized starch, silicon dioxide, sodium starch glycolate, talc and triacetin.

⚠ WARNING: This product can expose you to phenazopyridine hydrochloride, which is known to the state of California to cause cancer. For more information, visit www.P65Warnings.ca.gov

To report an adverse event call (888) 313-1372
© 2025 Uqora. Uqora is a trademark of Uqora.
Distributed by Uqora
4250 Executive Square, Suite 200, San Diego, CA 92037
support@uqora.com

UQORA MAXIMUM UTI PAIN RELIEF

phenazopyridine hydrochloride tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:73712-114
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PHENAZOPYRIDINE HYDROCHLORIDE (UNII: 0EWG668W17) (PHENAZOPYRIDINE - UNII:K2J09EMJ52)	PHENAZOPYRIDINE HYDROCHLORIDE	99.5 mg

Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A)	
STARCH, CORN (UNII: 08232NY3SJ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
TALC (UNII: 7SEV7J4R1U)	
POVIDONE (UNII: FZ989GH94E)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
TRACETIN (UNII: XHX3C3X673)	
LACTOSE (UNII: J2B2A4N98G)	
MAGNESIUM STEARATE (UNII: 70097M6130)	

Product Characteristics

Color	brown	Score	no score
Shape	OVAL	Size	9mm
Flavor		Imprint Code	p99
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73712-114-24	1 in 1 CARTON	10/27/2025	
1		24 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:73712-114-12	1 in 1 CARTON	10/27/2025	
2		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		10/27/2025	

Labeler - Bonafide Health, LLC dba Uqora (118327455)

Establishment

Name	Address	ID/FEI	Business Operations
Pharbest Pharmaceuticals, Inc.		557054835	manufacture(73712-114)

Revised: 10/2025

Bonafide Health, LLC dba Uqora