

OXY-GARD- hydrogen peroxide, lactic acid, dodecyl benzene sulfonic acid solution
Ecolab Inc.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

WARNING

KEEP OUT OF REACH OF CHILDREN

FOR INDUSTRIAL USE ONLY

NOT FOR HUMAN USE

Causes eye irritation

Wash skin thoroughly after handling. Do not mix with bleach or other chlorinated products – will cause chlorine gas.

DIRECTIONS:

IMPORTANT: Do not further dilute with water or mix with any other teat dips. If product in dip cup becomes visibly dirty, discard contents and replenish with fresh product. Do not reuse or return any unused product to the original container.

Udder Prep: When using an udder wash step before milking, make sure to wash teats with appropriate udder wash solution using proper cleaning procedures. Teats should then be dried with single-service towels.

Directions for Teat Dipping

Pre-Milk Dipping: Before each cow is milked, and using fresh Oxy-Gard, dip each teat full-length into the teat dip cup. Wipe teats after dipping using single-service towels to avoid contamination of milk.

Post-Milk Dipping: Using fresh Oxy-Gard, dip each teat full-length into the teat dip cup. Do not wipe. Allow to air dry. Do not turn cows out in freezing weather until the product is completely dry.

Directions for Teat Spraying

Pre-Milk Spraying: Before each cow is milked, and using fresh Oxy-Gard, spray entire teat. Wipe teats after dipping using single-service towels to avoid contamination of milk.

Post-Milk Spraying: Using fresh Oxy-Gard, spray entire teat immediately after each milking. Do not wipe. Allow to air dry. Do not turn cows out in freezing weather until the product is completely dry.

Expanded Usage: When freshening cows, begin dipping teats twice daily for about 10 days before calving.

PRECAUTION: Oxy-Gard is not intended to cure or help the healing of chapped or irritated teats. As with any germicide, irritation or sensitization may occur in sensitive animals. In case of teat irritation or chapping, have the condition examined and, if necessary, treated by a veterinarian.

Consult your Ecolab representative for specific use instructions and recommended dispensing equipment.

Read other label for directions for use, first aid and precautionary statements

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention.

READ SAFETY DATA SHEET (SDS) BEFORE USING THIS PRODUCT

EMERGENCY HEALTH INFORMATION: 1 800 328 0026. If located outside the United States and Canada, call collect 1 651 222 5352 (number is in the US).

Principal Display Panel / Representative Label

ECOLAB

Oxy-Gard™

Sanitizing Teat Dip

An aid in reducing the spread of organisms which may cause mastitis

ACTIVE INGREDIENTS:

Hydrogen Peroxide.....	0.5%
Lactic Acid.....	1.7%
Dodecyl Benzene Sulfonic Acid.....	0.5%

INERT INGREDIENTS:..... 97.3%

(contains glycerin, sorbitol)

TOTAL:..... 100.0%

766581/5300/1016

18.9 L (5 US GAL)

6301973

CAUTION

VENTED CONTAINER.

OPEN OR VENT

WITH CARE.

Ecolab · 1 Ecolab Place · St Paul MN 55102 USA · tel: 1 800 392 3392

© 2016 Ecolab USA Inc. · All rights reserved

Made in U.S.A.

www.ecolab.com · 766585/5300/1016

ECOLAB®



PRECAUCION
CONTENEDOR
VENTILADO.
ABRA O VENTILE
CON CUIDADO.

Oxy-Gard™

Sanitizing Teat Dip Desinfectante de pezones por inmersión

· An aid in reducing the spread of organisms which may cause mastitis | Ayuda a evitar la proliferación de organismos que pueden causar mastitis

ACTIVE INGREDIENTS:

Hydrogen Peroxide..... 0.5%

Lactic Acid 1.7%

Dodecyl Benzene Sulfonic Acid 0.5%

INERT INGREDIENTS:..... 97.3%

(contains glycerin, sorbitol)

TOTAL:..... 100.0%

Warning | Advertencia

KEEP OUT OF REACH OF CHILDREN | MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS

FOR INDUSTRIAL USE ONLY | PARA USO INDUSTRIAL SOLAMENTE

NOT FOR HUMAN USE | NO APTO PARA USO EN SERES HUMANOS

Read other label for directions for use, first aid and precautionary statements

Lea la otra etiqueta para direcciones de uso, primeros auxilios y medidas preventivas

766581/5300/1016



Teat Dip
Baño de tetina



Teat Spray
Rocedor de tetina

KEEP FROM
FREEZING
NO DEJE QUE
SE CONGEE

CAUTION
VENTED CONTAINER.
OPEN OR VENT
WITH CARE.

18.9 L (5 US GAL)
6301973



Oxy-Gard™

Sanitizing Teat Dip

FOR INDUSTRIAL USE ONLY

DIRECTIONS:

IMPORTANT: Do not further dilute with water or mix with any other teat dips. If product in dip cup becomes visibly dirty, discard contents and replenish with fresh product. Do not reuse or return any unused product to the original container.

Udder Prep: When using an udder wash step before milking, make sure to wash teats with appropriate udder wash solution using proper cleaning procedures. Teats should then be dried with single-service towels.

Directions for Teat Dipping

Pre-Milk Dipping: Before each cow is milked, and using fresh Oxy-Gard, dip each teat full-length into the teat dip cup. Wipe teats after dipping using single-service towels to avoid contamination of milk.

Post-Milk Dipping: Using fresh Oxy-Gard, dip each teat full-length into the teat dip cup. Do not wipe. Allow to air dry. Do not turn cows out in freezing weather until the product is completely dry.

Directions for Teat Spraying

Pre-Milk Spraying: Before each cow is milked, and using fresh Oxy-Gard, spray entire teat. Wipe teats after dipping using single-service towels to avoid contamination of milk.

Post-Milk Spraying: Using fresh Oxy-Gard, spray entire teat immediately after each milking. Do not wipe. Allow to air dry. Do not turn cows out in freezing weather until the product is completely dry.

Expanded Usage: When freshening cows, begin dipping teats twice daily for about 10 days before calving.

PRECAUTION: Oxy-Gard is not intended to cure or help the healing of chapped or irritated teats. As with any germicide, irritation or sensitization may occur in sensitive animals. In case of teat irritation or chapping, have the condition examined and, if necessary, treated by a veterinarian.

Consult your Ecolab representative for specific use instructions and recommended dispensing equipment.

KEEP OUT OF REACH OF CHILDREN

NOT FOR HUMAN USE

Warning

Causes eye irritation.

Wash skin thoroughly after handling. Do not mix with bleach or other chlorinated products - will cause chlorine gas.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/ attention.

READ SAFETY DATA SHEET (SDS) BEFORE USING THIS PRODUCT

EMERGENCY HEALTH INFORMATION: 1 800 328 0026. If located outside the United States and Canada, call collect 1 651 222 5352 (number is in the US).

Ecolab - 1 Ecolab Place - St Paul MN 55102 USA - tel: 1 800 392 3392
 © 2016 Ecolab USA Inc. - All rights reserved | Todos los derechos reservados
 Made in U.S.A. | Hecho en EE.UU.
 www.ecolab.com - 766585/5300/1016

Desinfectante de pezones por inmersión

PARA USO INDUSTRIAL SOLAMENTE

INSTRUCCIONES:

IMPORTANTE: No lo vuelva a diluir con agua ni lo mezcle con otros productos para baño de pezones. Si el producto del recipiente para realizar el baño está visiblemente sucio, desechelo y rellene el recipiente con producto fresco. No vuelva a utilizar el producto ni vuelva a colocar producto sin usar en el contenedor original.

Preparado para ubres: Si se realiza un paso de limpieza de ubres antes del ordeño, asegúrese de limpiar los pezones con una solución para limpieza de ubres adecuada y procedimientos de limpieza apropiados. Los pezones deben secarse con toallas de uso único.

Instrucciones para el sellado de pezones

Sellado del pezón previo al ordeño: Antes de ordeñar cada vaca, bañe toda la superficie de cada pezón en un recipiente que contenga Oxy-Gard fresco. Limpie los pezones con toallas de uso único luego del baño para evitar la contaminación de la leche.

Sellado del pezón posterior al ordeño: Bañe toda la superficie de cada pezón en un recipiente que contenga Oxy-Gard fresco. No utilice paños. Deje secar al aire. No deje las vacas en un clima helado hasta que el producto se haya secado por completo.

Instrucciones para el rociado de pezones

Rociado previo al ordeño: Antes de ordeñar cada vaca, rocíe toda la superficie de cada pezón con Oxy-Gard fresco. Limpie los pezones con toallas de uso único luego del baño para evitar la contaminación de la leche.

Rociado de pezones posterior al ordeño: Inmediatamente después del ordeño, rocíe toda la superficie de cada pezón con Oxy-Gard fresco. No utilice paños. Deje secar al aire. No deje las vacas en un clima helado hasta que el producto se haya secado por completo.

Uso ampliado: Si la vaca va a parir, comience a realizar el baño dos veces al día unos 10 días antes del parto.

PRECAUCIÓN: Oxy-Gard no está pensado para curar ni ayudar a sanar los pezones agrietados o irritados. Como cualquier germicida, el producto puede causar irritación o sensibilización en los animales sensibles. Si los pezones están agrietados o irritados, encargue a un veterinario que examine la condición y la trate si es necesario.

Consulte con su representante de Ecolab para obtener instrucciones específicas sobre el uso y el equipo dispensador recomendado.

MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS

NO ES PARA USO HUMANO

Advertencia

Provoca irritación ocular.

Lave la piel minuciosamente después del manejo. No se mezcle con el blanqueo o otros productos tratados con cloro - causará el gas de cloro.

EN CASO DE CONTACTO CON LOS OJOS: Enjuagar con agua cuidadosamente durante varios minutos. Quitar las lentes de contacto cuando estén presentes y pueda hacerse con facilidad. Proseguir con el lavado. Si la irritación ocular persiste, consultar a un médico.

LEA LA FICHA DE DATOS DE SEGURIDAD ANTES DE USAR EL PRODUCTO

INFORMACIÓN DE EMERGENCIA SOBRE SALUD: 1 800 328 0026. Si está fuera de los Estados Unidos y Canadá, llame a cobro revertido al 1 651 222 5352 (número en los EE.UU.).

OXY-GARD

hydrogen peroxide, lactic acid, dodecyl benzene sulfonic acid solution

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:47593-119
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HYDROGEN PEROXIDE (UNII: BBX060AN9V) (HYDROGEN PEROXIDE - UNII:BBX060AN9V)	HYDROGEN PEROXIDE	5 mg in 1 mL
DODECYLBENZENESULFONIC ACID (UNII: 60NSK897G9) (DODECYLBENZENESULFONIC ACID - UNII:60NSK897G9)	DODECYLBENZENESULFONIC ACID	5 mg in 1 mL
LACTIC ACID (UNII: 33X04XA5AT) (LACTIC ACID - UNII:33X04XA5AT)	LACTIC ACID	17 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
SORBITOL (UNII: 506T60A25R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47593-119-16	18930 mL in 1 PAIL		
2	NDC:47593-119-17	56780 mL in 1 DRUM		
3	NDC:47593-119-21	208200 mL in 1 DRUM		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		01/09/1990	

Labeler - Ecolab Inc. (006154611)