

**VICKS DAYQUIL NYQUIL HOT REMEDY COLD AND FLU- acetaminophen,
dextromethorphan hbr, diphenhydramine hcl
Procter & Gamble Manufactura S de RL de CV**

**Vicks® @ DayQuil™ HOT REMEDY Vicks® NyQuil™ HOT REMEDY
COLD & FLU**

Vicks® DayQuil™ HOT REMEDY COLD & FLU

Drug Facts

DayQuil™

Active Ingredient (in each Packet)

Acetaminophen 500 mg

Dextromethorphan HBr 20 mg

Purpose

Headache, Fever, Sore Throat, Minor Aches & Pains

Cough

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert:

Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening

- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning:

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- do not exceed 6 doses per 24 hrs
- Dissolve contents of one packet into 8 oz. of hot water and stir briskly; sip while hot. Consume entire drink within 10-15 minutes
- If using a microwave, add contents of one packet to 8 oz. of cool water, stir briskly before and after heating. Do not overheat

Age	Dose
adults & children 12 yrs & over	one packet (dose) every 4 hours
children under 12 yrs	do not use

Other information

- **each packet contains:** potassium 10 mg
- phenylketonurics: contains phenylalanine 68 mg per dose
- store at no greater than 25°C (77°F)

Inactive ingredients:

acesulfame potassium, aspartame, carboxymethylcellulose, citric acid, D&C Yellow No. 10, FD&C Blue No. 1, FD&C Red No. 40, flavor, sodium citrate, sucrose, tricalcium phosphate

Questions?

1-800-362-1683

MADE IN MEXICO

DIST. BY PROCTER & GAMBLE,

CINCINNATI OH 45202

Vicks® NyQuil™ HOT REMEDY COLD & FLU

Drug Facts

NyQuil™

Active Ingredient (in each Packet)

Acetaminophen 650 mg

Diphenhydramine HCl 25 mg

Purpose

Pain reliever/fever reducer

Antihistamine/Cough suppressant

Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- minor aches & pains
- sore throat
- fever
- headache
- cough due to minor sore throat & bronchial irritation
- cough to help you sleep
- runny nose
- sneezing

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert:

Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning:

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- with any other product containing diphenhydramine, even one used on skin.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (Certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- excitability may occur especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- do not exceed 6 doses per 24 hrs
- Dissolve contents of one packet into 8 oz. of hot water and stir briskly; sip while hot. Consume entire drink within 10-15 minutes
- If using a microwave, add contents of one packet to 8 oz. of cool water, stir briskly before and after heating. Do not overheat

Age	Dose
adults & children 12 yrs & over	one packet (dose) every 4 hours
children under 12 yrs	do not use

Other information

- **each packet contains:**potassium 10 mg
- phenylketonurics: contains phenylalanine 68 mg per dose
- store at no greater than 25°C (77°F)

Inactive ingredients

acesulfame potassium, aspartame, carboxymethylcellulose, citric acid, D&C Yellow No. 10, FD&C Blue No. 1, FD&C Red No. 40, flavor, sodium citrate, sucrose, tricalcium phosphate

Questions?

1-800-362-1683

TAMPER EVIDENT: DO NOT USE IF SACHET IS OPENED OR TORN

Keep outer package for complete product information.

MADE IN MEXICO

**DIST. BY PROCTER & GAMBLE,
CINCINNATI OH 45202**

PRINCIPAL DISPLAY PANEL - 6 packet package

NEW FORMULAS

VALUE PACK

Vicks[®] DayQuil[™] HOT REMEDY

COLD & FLU

Acetaminophen, Dextromethorphan HBr

Headache, Fever, Sore Throat, Minor Aches & Pains

Cough

Non-Drowsy

Vicks[®] NyQuil[™] HOT REMEDY

COLD & FLU

Acetaminophen, Diphenhydramine HCl

Headache, Fever, Sore Throat, Minor Aches & Pains

Cough, Sneezing, Runny Nose

Nighttime Relief

HONEY LEMON FLAVORED

3 DAYQUIL PACKETS; 3 NYQUIL PACKETS; 6 TOTAL PACKETS



Drug Facts	NyQuil™	Drug Facts	DayQuil™
Active Ingredients (in each Packet)	Purpose	Active Ingredients (in each Packet)	Purpose
Acetaminophen 500 mg	Pain reliever/Fever reducer	Acetaminophen 500 mg	Pain reliever/Fever reducer
Diphenhydramine HCl 25 mg	Antihistamine/Cough suppressant	Dextromethorphan HBr 20 mg	Cough suppressant
Uses Temporarily relieves common cold/flu symptoms: • minor aches & pains • sore throat • fever • headache • cough due to minor throat & bronchial irritation • cough to help you sleep • runny nose • sneezing		Uses Temporarily relieves common cold/flu symptoms: • cough due to minor throat & bronchial irritation • sore throat • headache • minor aches & pains • fever	
Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: • more than 6 doses in 24 hours, which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin redness • blisters • rash If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 7 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.		Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: • more than 6 doses in 24 hours, which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin redness • blisters • rash If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 7 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.	
Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • with any other product containing diphenhydramine, even one used on skin • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist.		Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist.	

Drug Facts (continued)	NyQuil™	Drug Facts (continued)	DayQuil™
Ask a doctor before taking this product. Ask a doctor before use if you have • liver disease • glaucoma • cough that occurs with too much phlegm (mucus) • a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema • trouble urinating due to enlarged prostate gland Ask a doctor or pharmacist before use if you are • taking sedatives or tranquilizers • taking the blood thinning drug warfarin When using this product • excitability may occur, especially in children • marked drowsiness may occur • alcohol, sedatives, and tranquilizers may increase drowsiness • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery Stop use and ask a doctor if • pain or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.		Know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. Ask a doctor before use if you have • liver disease • cough that occurs with too much phlegm (mucus) • persistent or chronic cough such as occurs with smoking, asthma, or emphysema Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin. Stop use and ask a doctor if • pain or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	
Directions • take only as directed • do not exceed 6 doses per 24 hrs • Dissolve contents of one packet into 8 oz. of hot water and stir briskly; sip while hot. Consume entire drink within 10-15 minutes. • If using a microwave, add contents of one packet to 8 oz. of cool water, stir briskly before and after heating. Do not overheat. Age adults & children 12 yrs & over one packet (dose) every 4 hours children under 12 yrs do not use Other information • each packet contains: potassium 10 mg • phenylephrine: contains phenylephrine 58 mg per dose • store at no greater than 25°C (77°F) Inactive ingredients acacia, potassium, aspartame, carboxymethylcellulose, citric acid, D&G Yellow No. 10, FD&C Blue No. 1, FD&C Red No. 40, flavor, sodium citrate, sucrose, tricalcium phosphate Questions? 1-800-362-1683		Directions • take only as directed • do not exceed 6 doses per 24 hrs • Dissolve contents of one packet into 8 oz. of hot water and stir briskly; sip while hot. Consume entire drink within 10-15 minutes. • If using a microwave, add contents of one packet to 8 oz. of cool water, stir briskly before and after heating. Do not overheat. Age adults & children 12 yrs & over one packet (dose) every 4 hours children under 12 yrs do not use Other information • each packet contains: potassium 10 mg • phenylephrine: contains phenylephrine 58 mg per dose • store at no greater than 25°C (77°F) Inactive ingredients acacia, potassium, aspartame, carboxymethylcellulose, citric acid, D&G Yellow No. 10, FD&C Blue No. 1, FD&C Red No. 40, flavor, sodium citrate, sucrose, tricalcium phosphate Questions? 1-800-362-1683	

VICKS DAYQUIL NYQUIL HOT REMEDY COLD AND FLU

acetaminophen, dextromethorphan hbr, diphenhydramine hcl kit

Product Information

Product Type HUMAN OTC DRUG **Item Code (Source)** NDC:58933-554

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58933-554-16	1 in 1 PACKAGE; Type 0: Not a Combination Product	06/11/2025	
2	NDC:58933-554-06	1 in 1 PACKAGE; Type 0: Not a Combination Product	06/11/2025	

Quantity of Parts		
Part #	Package Quantity	Total Product Quantity
Part 1	1 PACKET	5 g
Part 2	1 PACKET	5 g

Part 1 of 2

VICKS NYQUIL HOT REMEDY COLD AND FLU

acetaminophen, diphenhydramine hcl powder, for solution

Product Information	
Item Code (Source)	NDC:58933-553
Route of Administration	ORAL

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg in 5 g
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 5 g

Inactive Ingredients	
Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
TRICALCIUM PHOSPHATE (UNII: K4C08XP666)	
SUCROSE (UNII: C151H8M554)	
ASPARTAME (UNII: Z0H242BBR1)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
CARBOXYMETHYLCELLULOSE (UNII: 05JZI7B19X)	

Product Characteristics			
Color	white	Score	
Shape		Size	
Flavor	HONEY (Honey Lemon)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58933-553-01	5 g in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/11/2025	

Part 2 of 2

VICKS DAYQUIL HOT REMEDY COLD AND FLU

acetaminophen, dextromethorphan hbr powder, for solution

Product Information

Item Code (Source)	NDC:58933-552
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 5 g
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg in 5 g

Inactive Ingredients

Ingredient Name	Strength
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
SUCROSE (UNII: C151H8M554)	
TRICALCIUM PHOSPHATE (UNII: K4C08XP666)	
CARBOXYMETHYLCELLULOSE (UNII: 05JZI7B19X)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
ASPARTAME (UNII: Z0H242BBR1)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	

Product Characteristics

Color	white	Score	
-------	-------	-------	--

Shape		Size		
Flavor	HONEY (Honey Lemon)	Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58933-552-01	5 g in 1 PACKET; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M012	06/11/2025	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M012	06/11/2025	

Labeler - Procter & Gamble Manufactura S de RL de CV (812807550)