

ZILACTIN-B- benzocaine gel
ASCENT CONSUMER PRODUCTS, INC

Zilactin B

Drug Facts

Active ingredient

Benzocaine 10%

Purpose

Oral pain reliever

Use

temporarily relieves pain caused by:

- canker sores
- minor mouth sores and gum irritations
- denture and brace pain

Warnings

Flammable:

- keep away from fire or flame

Allergy Alert:

- do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other “caine” anesthetics

Stop use and ask a physician if

- sore mouth symptoms do not improve in 7 days
- swelling, rash or fever develops
- irritation, pain or redness persists or worsens
- apply only to affected area
- do not exceed recommended dosage
- avoid contact with the eyes
- do not use for more than 7 days unless directed by a physician or dentist

Keep out of reach of children. If swallowed, get medical help or contact a poison control center immediately.

Directions

adults and children 2 years and older	Dry affected area. Apply a thin coat of gel with cotton swab or clean finger up to 4 times daily. Allow to dry 30-60 seconds.
children under 12 years	adult supervision should be given in the use of this product
children under 2 years	do not use, consult a physician or dentist

Other information

- do not peel off protective film. Attempting to peel off film may result in skin irritation or tenderness. To remove film, first apply another coat of Zilactin-B Long Lasting Mouth Sore Gel to film, and immediately wipe the area with a moist gauze pad or tissue.
- contains alcohol 65% by volume
- store at 15-30°C (59-86°F)

Inactive ingredients

Boric acid, hydroxypropylcellulose, propylene glycol, purified water, salicylic acid, SD alcohol 38-B, tannic acid

Call 1-888-547-5492

Distributed by Ascent Consumer Products



Oral Pain
Reliever

zilactin-B[®]

ORAL PAIN RELIEVER

6 Hour Canker & Mouth Sore Relief

• Bioadhesive
Promotes Healing

• Eat & Drink Pain-Free
for up to 6 Hours



LOVE THIS STUFF

My dental hygienist recommends Zilactin-B for
my mouth ulcers. It has been a life saver! — D. Ziegler



Feel It.
Heal It.[™]

NET WT. .25 OZ. (7g)

FPO
UPC# 6-46011-55032-7

X XXXXX XXXXX X

Distributed by:
Ascent Consumer Products
Melville, NY 11747 USA ©2024
Questions? 1-888-547-5492
www.zilactin.com

ACP# B5503-101-01

Drug Facts	
Active ingredient Benzocaine 10% Purpose Oral pain reliever	
Uses Temporarily relieves pain caused by: ■ canker sores ■ minor mouth sores and gum irritations ■ denture and braces pain	
Warnings ■ Medicallabeling warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used the product before. Stop use and seek immediate medical attention if you or a child in your care develops pale, gray, or blue colored skin (cyanosis), headache, rapid heart rate, shortness of breath, dizziness or lightheadedness, fatigue or lack of energy.	
Flammable: ■ keep away from fire or flame	
Allergy Alert: ■ do not use this product if you have a history of allergy to local anesthetics such as procaine, butabaine, benzocaine or other "caine" anesthetics	
Stop use and ask a physician if: ■ some mouth symptoms do not improve in 7 days ■ swelling, rash or liver develops ■ irritation, pain or redness persists or worsens ■ do not exceed recommended dosage ■ avoid contact with the eyes ■ do not use for more than 7 days unless directed by a physician or dentist	
Keep out of reach of children. If swallowed, get medical help or contact a poison control center immediately.	
Do not use: ■ for teething ■ in children under 3 years of age	
Directions remove cap and cut open tip of tube on scoremark adults and children 3 years and older Dry affected area. Apply a thin coat of gel with cotton swab or clean finger up to 4 times daily. Allow to dry 30-60 seconds. adult supervision should be given in the use of this product	
children under 12 years	do not use
children under 3 years	do not use
Other information ■ do not peel off protective film. Attempting to peel off film may result in skin irritation or tenderness. To remove film, first apply another coat of Zilactin-B Long Lasting Mouth Sore Gel to film, and immediately wipe the area with a moist gauze pad or tissue. ■ Zilactin-B may darken the gel's appearance over time, but its effectiveness is unchanged. ■ contains alcohol 5% by volume ■ store at 15-30°C (59-86°F)	
Inactive ingredients benzoic acid, hydroxypropylcellulose, propylene glycol, purified water, salicylic acid, SD alcohol 3B-9, stearic acid	

Long Lasting
Mouth Sore Gel

zilactin-B

ZILACTIN-B

benzocaine gel

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:42829-550
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	0.1 g in 1 g

Inactive Ingredients

Ingredient Name	Strength
BORIC ACID (UNII: R57ZHV85D4)	
HYDROXYPROPYL CELLULOSE (TYPE M) (UNII: U3JF91U133)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SALICYLIC ACID (UNII: O414PZ4LPZ)	
ALCOHOL (UNII: 3K9958V90M)	
PEPPERMINT OIL (UNII: AV092KU4JH)	
TANNIC ACID (UNII: 28F9E0DJY6)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:42829-550-32	1 in 1 CARTON	11/06/2025	
1		7 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M022	11/06/2025	

Labeler - ASCENT CONSUMER PRODUCTS, INC (078396381)