# GOOD NEIGHBOR PHARMACY CHILDRENS PAIN AND FEVER-

# acetaminophen suspension

### Amerisource Bergen

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

# Amerisource Bergen Children's Pain & Fever Drug Facts

### Active ingredient (in each 5 mL)

Acetaminophen 160 mg

# Purpose

Pain reliever/fever reducer

#### Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to:
- the common cold
- flu
- headache
- sore throat
- toothache

### **Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### Do not use

• with any other drug containing acetaminophen (prescription or nonprescription). If you are not

sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

• if your child has ever had an allergic reaction to this product or any of its ingredients

# Ask a doctor before use if your child has

liver disease

# Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin

### When using this product

do not exceed recommended dose (see overdose warning)

# Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

### Keep out of reach of children.

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

#### **Directions**

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see overdose warning)
- shake well before using
- mL = milliliter
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child's dose into the dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)*	
under 24	under 2 years	ask a doctor	
24-35	2-3 years	5 mL	
36-47	4-5 years	7.5 mL	
48-59	6-8 years	10 mL	
60-71	9-10 years	12.5 mL	
72-95	11 years	15 mL	

<sup>\*</sup>or as directed by a doctor

**Attention:** use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

### Other information

- each 5 mL contains: sodium 3 mg
- store at 20-25°C (68-77°F)
- do not use if printed neckband is broken or missing

# **Inactive ingredients**

anhydrous citric acid, calcium sulfate, carrageenan, FD&C red #40, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, propylparaben, purified water, sodium benzoate, sorbitol solution, tribasic sodium phosphate

### Questions or comments?

1-800-719-9260

# **Principal Display Panel**

Compare to Children's Tylenol® Oral Suspension active ingredient

FOR AGES 2 TO 11 YEARS

children's Pain & Fever

acetaminophen 160 mg per 5 mL

oral suspension

Fever Reducer – Pain Reliever


Aspirin Free

Ibuprofen Free

Alcohol Free

Strawberry Flavor

4 fl oz (118 mL)





#### Drug Facts

Active inaredient (in each 5 ml.)

Pain reliever/fever reduc

Purpose

#### Uses temporarily:

- USES temperature.

  If reduces fever

  If relieves minor aches and pains due to:

  If the common cold

  If the landache

  sore throat

  I toothache

### Warnings

warming: This product contains acetaminophen. Severe liver damage may occur if your child takes ■ more than 5 doses in 24 hours, which is the maximu

■ more than is uneen use and adja ancent adja ancent adja ancent.
■ with other drugs containing acetaminophen
Allergy alert. Acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin reddening ■ bisters ■ rash
fa skin reaction occurs, stop use and seek medical help

If a skin reaction occurs, stop use and seek medical he right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headoche, rash, nausea, or vomiting, consult a doctor promptly.

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has liver disease

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

When using this product do not exceed recommended dose (see overdose warning)

#### Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days fever gets worse or lasts more than 3 days
- new symptoms occur
  redness or swelling is present
  These could be signs of a serious condition.

Keep out of reach of children.
Overdose warning: In case of overdose, get medical
help or contact a Poison Control Center right away
(1-800-222-1222). Quick medical attention is critical ven if you do not notice any signs or symptoms.

#### Drug Facts (continued)

#### Directions

- this product does not contain directions or complete warnings for adult use
   do not give more than directed (see overdose

- do not give more than curected sees created warning)
   shake well before using
   ml. = millitier
   if ind right dose on chart below. If possible, use weight to dose; otherwise, use age.
   remove the child protective cap and squeeze your child's dose into the dosing cup
   receat dose every 4 hours white symptoms last

■ repeat dose every 4 nours write symptoms ia	31
■ do not give more than 5 times in 24 hours	

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL
72-95	11 years	15 mL

\*or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

#### Other information

- each 5 mL contains: sodium 3 mg
   store at 20-25°C (68-7°F)
   do not use if printed neckband is broken or missing

#### Inactive ingredients

Intactive Ingredients
anhydrous citic acid, calcium sulfate, carrageenan,
FD&C red #40, flavor, glycerin, high fructose com
syrup, hydrosyethyl cellulose, microcrystalline cellulose
and carboxymethylcellulose sodium, propylene glycol,
propyleratein, purified water, sodium benzeate,
sorbitol solution, tribasic sodium phosphate

#### Questions or comments?

\*\*Good Neighbor Pharmacy® Children's Pain & Fever is not manufactured or distributed by Johnson & Johnson Con Inc., distributor of Children's Tylenol® Oral Suspension.

#### **GLUTEN FREE**

Distributed By AmeriscurceBergen 1300 Morris Drive, Chesterbrook, PA 19087 Questions or Concerns? www.mygnp.com



Children's Tylenol





acetaminophen 160 mg per 5 mL oral suspension

#### Fever Reducer – Pain Reliever

Aspirin Free Ibuprofen Free Alcohol Free

Strawberry Flavor







DIRECTIONS: STEP 1. Shake v before using




STEP 2. Remove child protective cap



STEP 3. Squeeze your child's dose into the dosing cup



IMPORTANT: Please see dosing chart or ask a doctor for your child's correct dose.





# GOOD NEIGHBOR PHARMACY CHILDRENS PAIN AND FEVER

acetaminophen suspension

### **Product Information**

HUMAN OTC DRUG NDC:46122-212 Product Type Item Code (Source)

Route of Administration

ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINO PHEN (UNII: 36209 ITL9 D) (ACETAMINO PHEN - UNII: 36209 ITL9 D)	ACETAMINOPHEN	160 mg in 5 mL	

Inactive Ingredients	
Ingredient Name	Strength
ANHYDRO US CITRIC ACID (UNII: XF417D3PSL)	
CALCIUM SULFATE, UNSPECIFIED FORM (UNII: WAT0 DDB505)	
CARRAGEENAN (UNII: 5C69 YCD2YJ)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
MICRO CRYSTALLINE CELLULO SE (UNII: OP1R32D61U)	
CARBO XYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K679 OBS311)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SODIUM PHO SPHATE, TRIBASIC (UNII: A752Q30A6X)	

Product Characteristics			
Color	RED	Score	
Shape		Size	
Flavor	STRAWBERRY	Imprint Code	
Contains			

Packaging			
# Item Code	Package Description	<b>Marketing Start Date</b>	Marketing End Date
1 NDC:46122-212-26	1 in 1 CARTON	02/01/2011	
1	118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	02/01/2011	

# Labeler - Amerisource Bergen (007914906)

Revised: 12/2019 Amerisource Bergen