

**TRIPLE ANTIBIOTIC- bacitracin zinc, neomycin sulfate, polymyxin b,
pramoxine hcl ointment
Meijer Distribution Inc**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Meijer Distribution, Inc. Triple Antibiotic Drug Facts

Active ingredients (in each gram)

Bacitracin zinc 500 units

Neomycin sulfate equivalent to 3.5 mg of neomycin base

Polymyxin B 10,000 units as polymyxin B sulfate

Pramoxine HCl 10 mg

Purpose

First aid antibiotic

External analgesic

Uses

first aid to help prevent infection and for temporary relief of pain or discomfort in minor:

- cuts
- scrapes
- burns

Warnings

For external use only.

Do not use

- if you are allergic to any of the ingredients
- in the eyes
- over large areas of the body

Ask a doctor before use if you have

- deep or puncture wounds
- animal bites
- serious burns

Stop use and ask a doctor if

- you need to use longer than 1 week
- condition persists or gets worse
- symptoms persist for more than 1 week, or clear up and occur again within a few days
- rash or other allergic reaction develops

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- adults and children 2 years of age and older:
- clean the affected area
- apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
- may be covered with a sterile bandage
- children under 2 years of age: ask a doctor

Other information

- store at 20-25°C (68-77°F)

Inactive ingredients

cocoa butter, cottonseed oil, olive oil, petrolatum, sodium pyruvate, tocopheryl acetate

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

MAXIMUM STRENGTH

triple antibiotic

Bacitracin Zinc | Neomycin Sulfate | Polymyxin B Sulfate | Pramoxine HCl

+ PAIN | ITCH | SCAR

24 HOUR Infection Protection

Multi-Action Ointment

Meijer®

Compare to Neosporin® + Pain Itch Scar active ingredients

MAXIMUM STRENGTH

triple antibiotic

Bacitracin Zinc | Neomycin Sulfate | Polymyxin B Sulfate | Pramoxine HCl

+ PAIN | ITCH | SCAR

NET WT 0.5 OZ (14 g)

First Aid Antibiotic/Pain Relieving Ointment

MAXIMUM STRENGTH
triple antibiotic

Bacitracin Zinc | Neomycin Sulfate | Polymyxin B Sulfate | Pramoxine HCl

+ PAIN | ITCH | SCAR

24 HOUR Infection Protection
Multi-Action Ointment

Powerful Pain & Itch Relief for
Cuts | Scrapes | Burns

meijer

MAXIMUM STRENGTH
triple antibiotic

Bacitracin Zinc | Neomycin Sulfate | Polymyxin B Sulfate | Pramoxine HCl

+ PAIN | ITCH | SCAR

NET WT 0.5 OZ (14 g)

First Aid Antibiotic/Pain Relieving Ointment

NDC 41250-430-58
Compare to Neosporin®
+ Pain Itch Scar
active ingredients*

Code Area

*This product is not affiliated with, manufactured by, or produced by the makers of Neosporin®.

DIST. BY MEIJER DISTRIBUTION, INC.
GRAND RAPIDS, MI 49544
www.meijer.com



Drug Facts

Active ingredients (in each gram)

Bacitracin zinc 500 units.....First aid antibiotic
Neomycin sulfate equivalent to 3.5 mg
of neomycin base.....First aid antibiotic
Polymyxin B 10,000 units as
polymyxin B sulfate.....First aid antibiotic
Pramoxine HCl 10 mg.....External analgesic

Purpose

Uses first aid to help prevent infection and for temporary relief of pain or discomfort in minor:
■ cuts ■ scrapes ■ burns

Warnings

For external use only.

Do not use
■ if you are allergic to any of the ingredients
■ in the eyes
■ over large areas of the body

Ask a doctor before use if you have

■ deep or puncture wounds
■ animal bites
■ serious burns

Drug Facts (continued)

Stop use and ask a doctor if

■ you need to use longer than 1 week
■ condition persists or gets worse
■ symptoms persist for more than 1 week, or clear up and occur again within a few days
■ rash or other allergic reaction develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

■ adults and children 2 years of age and older:
■ clean the affected area
■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
■ may be covered with a sterile bandage
■ children under 2 years of age: ask a doctor

Other information ■ store at 20-25°C (68-77°F)

Inactive ingredients cocoa butter, cottonseed oil, olive oil, petrolatum, sodium pyruvate, tocopheryl acetate

Questions or comments? 1-800-719-9260

Code Area

TRIPLE ANTIBIOTIC

bacitracin zinc, neomycin sulfate, polymyxin b, pramoxine hcl ointment

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-433
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BACITRACIN ZINC (UNII: 89Y4M234ES) (BACITRACIN - UNII:58H6RWO52I)	BACITRACIN	500 [USP'U] in 1 g
NEOMYCIN SULFATE (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)	NEOMYCIN	3.5 mg in 1 g
POLYMYXIN B SULFATE (UNII: 19371312D4) (POLYMYXIN B - UNII:J2VZ07J96K)	POLYMYXIN B	10000 [USP'U] in 1 g
PRAMOXINE HYDROCHLORIDE (UNII: 88AYB867L5) (PRAMOXINE - UNII:068X84E056)	PRAMOXINE HYDROCHLORIDE	10 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
COCOA BUTTER (UNII: 512OYT1CRR)	
OLIVE OIL (UNII: 6UYK2W1W1E)	
PETROLATUM (UNII: 4T6H12BN9U)	
SODIUM PYRUVATE (UNII: POD38AIF08)	
COTTONSEED OIL (UNII: H3E878020N)	
.ALPHA.-TOCOPHEROL ACETATE (UNII: 9E8X80D2L0)	

Product Characteristics

Color	YELLOW (Pale)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-433-58	1 in 1 CARTON	10/03/2017	
1		14 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

Labeler - Meijer Distribution Inc (006959555)

Revised: 6/2022

Meijer Distribution Inc