

**TRIPLE ANTIBIOTIC- bacitracin zinc, neomycin sulfate, polymyxin b,  
pramoxine hcl ointment  
Meijer Distribution Inc**

-----  
**Meijer Distribution, Inc. Triple Antibiotic Drug Facts**

**Active ingredients (in each gram)**

Bacitracin zinc 500 units

Neomycin sulfate equivalent to 3.5 mg of neomycin base

Polymyxin B 10,000 units as polymyxin B sulfate

Pramoxine HCl 10 mg

**Purpose**

First aid antibiotic

External analgesic

**Uses**

first aid to help prevent infection and for temporary relief of pain or discomfort in minor:

- cuts
- scrapes
- burns

**Warnings**

**For external use only.**

**Do not use**

- if you are allergic to any of the ingredients
- in the eyes
- over large areas of the body

**Ask a doctor before use if you have**

- deep or puncture wounds
- animal bites
- serious burns

**Stop use and ask a doctor if**

- you need to use longer than 1 week

- condition persists or gets worse
- symptoms persist for more than 1 week, or clear up and occur again within a few days
- rash or other allergic reaction develops

### **Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### **Directions**

- adults and children 2 years of age and older:
- clean the affected area
- apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
- may be covered with a sterile bandage
- children under 2 years of age: ask a doctor

### **Other information**

- store at 20-25°C (68-77°F)

### **Inactive ingredients**

cocoa butter, cottonseed oil, olive oil, petrolatum, sodium pyruvate, tocopheryl acetate

### **Questions or comments?**

**1-800-719-9260**

### **Package/Label Principal Display Panel**

meijer®

MAXIMUM STRENGTH

Triple Antibiotic

COMPARE TO NEOSPORIN® + PAIN ITCH SCAR ACTIVE INGREDIENTS

First Aid Antibiotic | Pain Relieving Ointment

24 HOUR INFECTION PROTECTION

POWERFUL PAIN AND ITCH RELIEF FOR MINOR CUTS, SCRAPES, AND BURNS

Minimize the Appearance of Scars

+ Pain • Itch • Scar

NET WT 0.5 OZ (14 g)

meijer.

MAXIMUM STRENGTH

# Triple Antibiotic

First Aid Antibiotic | Pain Relieving Ointment

NDC 41250-433-58

COMPARE TO NEOSPORIN®  
+ PAIN ITCH SCAR  
ACTIVE INGREDIENTS\*

24 HOUR  
INFECTION  
PROTECTION

POWERFUL PAIN AND ITCH RELIEF FOR  
MINOR CUTS, SCRAPES, AND BURNS

Minimizes the  
Appearance of Scars

+ Pain · Itch · Scar

NET WT 0.5 OZ (14 g)



how2recycle.info

PID 4144856



www.meijer.com/satisfaction

DIST. BY MEIJER DISTRIBUTION INC.  
GRAND RAPIDS, MI 49544  
www.meijer.com

\*This product is not affiliated with, manufactured by, or produced by the  
makers of Neosporin®.

Questions or comments? 1-800-719-9260

**Drug Facts (continued)**  
Inactive ingredients: cocoa butter, cottonseed oil, olive oil,  
petroleum, sodium pyruvate, tocopheryl acetate

**Drug Facts (continued)**  
Apply a small amount of this product in an amount equal to the  
surface area of the tip of a finger) on the area 1 to 3 times daily.  
■ may be covered with a sterile bandage  
■ children under 2 years of age: ask a doctor

**Other information** ■ store at 20-25°C (68-77°F)

**Drug Facts (continued)**  
Keep out of reach of children. If swallowed, get medical help or  
contact a Poison Control Center right away (1-800-222-1222).  
**Directions**  
■ adult & children 2 years of age and older:  
■ clean the affected area

**Drug Facts (continued)**  
Do not use  
■ If you are allergic to any of the ingredients  
■ in the eyes ■ over large areas of the body  
Ask a doctor before use if you have  
■ deep or puncture wounds  
■ animal bites ■ serious burns  
Stop use and ask a doctor if  
■ you need to use longer than 1 week  
■ you have persistent or get worse  
■ symptoms persist for more than 1 week, or clear up and occur again  
within a few days  
■ rash or other allergic reaction develops

**Drug Facts**  
**Active ingredients (in each gram)**  
Bacitracin zinc 500 units, Neomycin sulfate equivalent to  
3.5 mg of neomycin base, Polymyxin B 10,000 units  
Polymyxin B sulfate, Pramoxine HCl 10 mg.  
External analgesic:  
Uses this aid to help prevent infection and for temporary relief of pain  
or discomfort in minor: ■ cuts ■ scrapes ■ burns  
**Warnings**  
For external use only.

## TRIPLE ANTIBIOTIC

bacitracin zinc, neomycin sulfate, polymyxin b, pramoxine hcl ointment

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41250-433
<b>Route of Administration</b>	TOPICAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BACITRACIN ZINC</b> (UNII: 89Y4M234ES) (BACITRACIN - UNII:58H6RWO52I)	BACITRACIN	500 [USP'U] in 1 g
<b>NEOMYCIN SULFATE</b> (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)	NEOMYCIN	3.5 mg in 1 g
<b>POLYMYXIN B SULFATE</b> (UNII: 19371312D4) (POLYMYXIN B - UNII:J2VZ.07J96K)	POLYMYXIN B	10000 [USP'U] in 1 g
<b>PRAMOXINE HYDROCHLORIDE</b> (UNII: 88AYB867L5) (PRAMOXINE - UNII:068X84E056)	PRAMOXINE HYDROCHLORIDE	10 mg in 1 g

## Inactive Ingredients

Ingredient Name	Strength
<b>COCOA BUTTER</b> (UNII: 512OYT1CRR)	
<b>OLIVE OIL</b> (UNII: 6UYK2W1W1E)	
<b>PETROLATUM</b> (UNII: 4T6H12BN9U)	
<b>SODIUM PYRUVATE</b> (UNII: POD38AIF08)	
<b>COTTONSEED OIL</b> (UNII: H3E878020N)	
<b>.ALPHA.-TOCOPHEROL ACETATE</b> (UNII: 9E8X80D2L0)	

## Product Characteristics

<b>Color</b>	YELLOW (Pale)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-433-58	1 in 1 CARTON	10/03/2017	
1		14 g in 1 TUBE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M004	10/03/2017	

