

TOPCARE OMEPRAZOLE- omeprazole tablet, orally disintegrating, delayed release

Topco Associates LLC

Topco Associates LLC. Omeprazole Drug Facts

Active ingredient (in each tablet)

Omeprazole 20 mg

Purpose

Acid reducer

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert:

- do not use if you are allergic to omeprazole.
- omeprazole may cause severe skin reactions. Symptoms may include:
 - skin reddening
 - blisters
 - rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss

- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- take 1 tablet before eating in the morning
- **do not crush or chew tablets**
- place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- do not take this medicine with alcohol

Repeated 14-Day Course (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.

- store at 20-25°C (68-77°F); keep product out of high heat and moisture
- take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

Inactive ingredients

amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, crospovidone, ferric oxide, flavor, hypromellose, hypromellose phthalate, maize maltodextrin, mannitol, microcrystalline cellulose, propylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Questions or comments?

1-800-719-9260: weekdays 7:30 AM to 5:00 PM EST

Package/Label Principal Display Panel

TopCare® health

COMPARE TO PRILOSEC OTC®

MELTech®

Melts In Your Mouth

MELTS IN YOUR MOUTH • DISSOLVES WITHOUT WATER

Omeprazole

DELAYED RELEASE ORALLY DISINTEGRATING TABLETS 20 mg

ACID REDUCER

24 HR

Treats Frequent Heartburn!

STRAWBERRY FLAVOR

DISSOLVE TABS

actual size

42 TABLETS

THREE 14-DAY COURSES OF TREATMENT • MAY TAKE 1 TO 4 DAYS FOR FULL EFFECT



Safety Feature –
Do not use if printed tablet
blister unit is open or torn.

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking



NDC 36800-601-55

+TopCare
health

COMPARE TO
PRILOSEC OTC®*

MELTech
Melts In Your Mouth

MELTS IN YOUR MOUTH • DISSOLVES WITHOUT WATER

Omeprazole

DELAYED RELEASE ORALLY DISINTEGRATING TABLETS 20 mg

ACID REDUCER



Treats Frequent
Heartburn!



STRAWBERRY
FLAVOR
DISSOLVE TABS



actual size

42 TABLETS

THREE 14-DAY COURSES OF TREATMENT • MAY TAKE 1 TO 4 DAYS FOR FULL EFFECT

1Y056 888 06

Drug Facts**Active ingredient (in each tablet)**

Omeprazole 20 mg

Purpose
Acid reducer**Use** ■ treats frequent heartburn (occurs 2 or more days a week)
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect**Warnings****Allergy alert:**

- do not use if you are allergic to omeprazole.
- omeprazole may cause severe skin reactions. Symptoms may include:
 - skin reddening
 - blisters
 - rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck, or shoulders; or lightheadedness
- frequent chest pain

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Drug Facts (continued)**Directions** ■ for adults 18 years of age and older

- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- take 1 tablet before eating in the morning
- do not crush or chew tablets
- place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- do not take this medicine with alcohol

Repeated 14-Day Course (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor

■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information ■ read the directions and warnings before use

- keep the carton. It contains important information.
- store at 20-25°C (68-77°F); keep product out of high heat and moisture
- take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

Inactive ingredients amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, crospovidone, ferric oxide, flavor, hypromellose, hypromellose phthalate, maize maltodextrin, mannitol, microcrystalline cellulose, propylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate**Questions or comments?** 1-800-719-9260 : weekdays 7:30 AM to 5:00 PM EST


Treats Frequent Heartburn!

ACID REDUCER

DELAYED RELEASE ORALLY DISINTEGRATING TABLETS 20 mg

Omeprazole

MELTS IN YOUR MOUTH • DISSOLVES WITHOUT WATER



DISTRIBUTED BY TOPCO ASSOCIATES LLC
ELK GROVE VILLAGE, IL 60007
©TOPCO PERA0223
QUESTIONS? 1-888-423-0139
topcare@topco.com
www.topcarebrand.com
MADE IN ISRAEL



Scan here for more information or call 1-888-423-0139

QUALITY GUARANTEED

This TopCare® product is laboratory tested to guarantee its highest quality. Your total satisfaction is guaranteed.

*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

TOPCARE OMEPRAZOLE

omeprazole tablet, orally disintegrating, delayed release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:36800-601
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OMEPRAZOLE (UNII: KG60484QX9) (OMEPRazole - UNII:KG60484QX9)	OMEPRazole	20 mg

Inactive Ingredients

Ingredient Name	Strength
DIMETHYLAMINOETHYL METHACRYLATE - BUTYL METHACRYLATE - METHYL METHACRYLATE COPOLYMER (UNII: 905HNO1SIH)	
ASCORBIC ACID (UNII: PQ6CK8PD0R)	
CETYL ALCOHOL (UNII: 936JST6JCN)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	

MANNITOL (UNII: 3OWL53L36A)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SODIUM STEARATE (UNII: QU7E2XA9TG)	
SODIUM STEARYL FUMARATE (UNII: 7CV7WJK4UI)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	RED (reddish)	Score	no score
Shape	ROUND	Size	9mm
Flavor	STRAWBERRY	Imprint Code	20
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-601-74	14 in 1 CARTON	04/02/2018	06/01/2023
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:36800-601-55	3 in 1 CARTON	04/02/2018	
2		14 in 1 CARTON		
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA209400	04/02/2018	

Labeler - Topco Associates LLC (006935977)