

**COLD AND FLU RELIEF- acetaminophen, dextromethorphan hbr, doxylamine succinate capsule, liquid filled  
Dolgenercorp, Inc. (DOLLAR GENERAL & REXALL)**

-----

**Drug Facts**

**Active ingredients (in each softgel)**

**Acetaminophen 325 mg**

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

**Purpose**

**Pain reliever/Fever Reducer**

Cough suppressant

Antihistamine

**Uses**

- temporarily relieves common cold and flu symptoms:
  - sore throat
  - headache
  - minor aches and pains
  - fever
  - runny nose and sneezing
  - cough due to minor throat and bronchial irritation

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks everyday while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists more than 2 days, is

accompanied or followed by fever, headache rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

### **Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

### **When using this product**

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

### **Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (0-800-222-1222) right away. Quick medical attention is critical for adults as well as

for children even if you do not notice any signs or symptoms.

### **Directions**

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 6 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- **when using other Daytime or Nighttime products, carefully read each label to insure correct dosing**

### **Other information**

- store between 15-30°C (59-86°F)
- avoid excessive heat

### **Inactive ingredients**

D&C yellow #10, FD&C blue #1, gelatin, glycerin, lecithin, light mineral oil, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

### **Questions or comments?**

Call **1-888-309-9030**

### **Principal Display Panel**

Compare to the active ingredients in Vicks® Nyquil® Cold & Flu LiquiCaps®\*\*

Night Time

Cold and Flu Relief

Acetaminophen- Pain Reliever/Fever Reducer

Dextromethorphan HBr- Cough Suppressant

Doxylamine Succinate- Antihistamine

Relieves:

- Aches, fever, sore throat
- Cough
- Runny nose, sneezing
  
- Multi-symptom

Softgel†

(†Liquid Filled Capsules)

\*\*This product is not manufactured or distributed by the Procter & Gamble Company.

Vicks® , NyQuil® , and LiquiCaps® are registered trademarks of the Procter & Gamble Company

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

DISTRIBUTED BY OLD EAST MAIN CO.

100 MISSION RIDGE

GOODLETTSVILLE, TN 37072

**Package Label**



Compare to the active ingredients of Vicks® NyQuil® Cold & Flu LiquiCaps®\*\*\*

# Night Time Cold & Flu Relief



**Acetaminophen - Pain Reliever/Fever Reducer**  
**Dextromethorphan HBr - Cough Suppressant**  
**Doxylamine Succinate - Antihistamine**

Relieves:

- Aches, fever, sore throat
- Cough
- Runny nose, sneezing



**8 Softgels<sup>†</sup>**

(\*Liquid Filled Capsules)

\*\*\*AMPER EVIDENCE: DO NOT USE IF CARTON IS OPENED OR IF BLISTER LIMIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

lot: \_\_\_\_\_  
 Date: \_\_\_\_\_

<p><b>Drug Facts (continued)</b></p> <p><b>Warnings</b></p> <p><b>Liver warning:</b> This product contains acetaminophen. Severe liver damage may occur if you take:</p> <ul style="list-style-type: none"> <li>■ more than 4,000 mg of acetaminophen in 24 hours</li> <li>■ with other drugs containing acetaminophen</li> <li>■ 3 or more alcoholic drinks every day while using this product</li> </ul> <p><b>Allergy alert:</b> Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash</p> <p>If a skin reaction occurs, stop use and seek medical help right away.</p> <p><b>Sore throat warning:</b> If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</li> <li>■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</li> </ul> <p><b>Ask a doctor before use if you have</b> ■ liver disease ■ glaucoma ■ trouble urinating due to an enlarged prostate gland ■ a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis or emphysema</p> <p><b>Ask a doctor or pharmacist before use if you are</b></p> <ul style="list-style-type: none"> <li>■ taking the blood thinning drug warfarin</li> <li>■ taking sedatives or tranquilizers</li> </ul> <p><b>When using this product</b> ■ do not exceed recommended dosage ■ excitability may occur, especially in children</p>	<p><b>Drug Facts (continued)</b></p> <p><b>Directions</b> ■ do not take more than directed (see Overdose warning)</p> <ul style="list-style-type: none"> <li>■ do not take more than 4 doses in 24 hours</li> <li>■ adults and children 12 years and over: take 2 softgels with water every 6 hours</li> <li>■ swallow whole; do not crush, chew, or dissolve</li> <li>■ children under 12 years: do not use</li> </ul> <p>to ensure correct dosing</p> <p>when using other Daytime or Nighttime products, carefully read each label</p> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>■ store between 15-30°C (59-86°F) ■ avoid excessive heat</li> </ul> <p><b>Inactive ingredients</b> D&amp;C yellow #10, FD&amp;C blue #1, gelatin, glycerin, lecithin, light mineral oil, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink</p> <p><b>Questions or comments?</b>          Call 1-888-309-9030</p>
<p><b>Drug Facts (continued)</b></p> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>■ pain or cough gets worse or lasts more than 7 days</li> <li>■ fever gets worse or lasts more than 3 days</li> <li>■ redness or swelling is present ■ new symptoms occur</li> <li>■ cough comes back or occurs with rash or headache that lasts.</li> </ul> <p>These could be signs of a serious condition.</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.</p> <p><b>Overdose warning:</b> Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p>	<p><b>Drug Facts (continued)</b></p> <p><b>Uses</b> ■ temporarily relieves common cold and flu symptoms: ■ minor aches and pains ■ headache ■ sore throat ■ runny nose and sneezing</p> <p>■ cough due to minor throat and bronchial irritation</p> <p><b>Antihistamine</b></p>
<p><b>Drug Facts (continued)</b></p> <p><b>Active ingredients (in each softgel)</b></p> <ul style="list-style-type: none"> <li>Acetaminophen 325 mg</li> <li>Pain reliever/fever reducer</li> <li>Dextromethorphan HBr 15 mg</li> <li>Cough suppressant</li> <li>Doxylamine succinate 6.25 mg</li> <li>Antihistamine</li> </ul>	

• Multi-symptom

Actual Softgel Size

Lot N  
Exp. |

\*\*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, NyQuil®, and LiquiCaps® are registered trademarks of The Procter & Gamble Company.

DISTRIBUTED BY OLD EAST MAIN CO.  
100 MISSION RIDGE  
GOODLETTSVILLE, TN 37072



## DOLLAR GENERAL HEALTH Nighttime Cold and Flu Relief

### COLD AND FLU RELIEF

acetaminophen, dextromethorphan hbr, doxylamine succinate capsule, liquid filled

#### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55910-910
<b>Route of Administration</b>	ORAL		

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

#### Inactive Ingredients

Ingredient Name	Strength
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>LECITHIN, SOYBEAN</b> (UNII: 1DI56QDM62)	
<b>LIGHT MINERAL OIL</b> (UNII: N6K5787QVP)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	

#### Product Characteristics

<b>Color</b>	green	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	21mm
<b>Flavor</b>		<b>Imprint Code</b>	PC10
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-910-08	8 in 1 CARTON	04/30/2023	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/30/2023	

**Labeler** - Dolgencorp, Inc. (DOLLAR GENERAL & REXALL) (068331990)

Revised: 3/2024

Dolgencorp, Inc. (DOLLAR GENERAL &amp; REXALL)