


NYSTATIN- nystatin suspension
Kesin Pharma Corporation

NYSTATIN ORAL SUSPENSION, USP
Rx only

DESCRIPTION

Nystatin is an antimycotic polyene antibiotic obtained from *Streptomyces noursei*.
Structural formula:



Nystatin Oral Suspension, USP for oral administration, is cherry/mint flavored, containing 100,000 USP Nystatin Units per mL. Inactive ingredients: cherry flavor, dibasic sodium phosphate heptahydrate, disodium edetate, glycerin, methylparaben, monobasic sodium phosphate monohydrate, peppermint oil, propylparaben, sodium benzoate, sodium hexametaphosphate, and sucrose.

CLINICAL PHARMACOLOGY

Pharmacokinetics

Gastrointestinal absorption of nystatin is insignificant. Most orally administered nystatin is passed unchanged in the stool. In patients with renal insufficiency receiving oral therapy with conventional dosage forms, significant plasma concentrations of nystatin may occasionally occur.

Microbiology

Nystatin is both fungistatic and fungicidal *in vitro* against a wide variety of yeasts and yeast-like fungi. *Candida albicans* demonstrates no significant resistance to nystatin *in vitro* on repeated subculture in increasing levels of nystatin; other *Candida species* become quite resistant. Generally, resistance does not develop *in vivo*. Nystatin acts by binding to sterols in the cell membrane of susceptible *Candida species* with a resultant change in membrane permeability allowing leakage of intracellular components. Nystatin exhibits no appreciable activity against bacteria, protozoa, or viruses.

INDICATIONS AND USAGE

Nystatin Oral Suspension is indicated for the treatment of candidiasis in the oral cavity.

CONTRAINDICATIONS

The preparation is contraindicated in patients with a history of hypersensitivity to any of its components.

PRECAUTIONS

General

This medication is not to be used for the treatment of systemic mycoses. Discontinue treatment if sensitization or irritation is reported during use.

Carcinogenesis, Mutagenesis, Impairment of Fertility

No long-term animal studies have been performed to evaluate carcinogenic potential. There also have been no studies to determine mutagenicity or whether this medication affects fertility in males or females.

Pregnancy

Teratogenic Effects Category C

Animal reproduction studies have not been conducted with nystatin oral suspension. It is also not known whether nystatin oral suspension can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Nystatin oral suspension should be given to a pregnant woman only if clearly needed.

Nursing Mothers

It is not known whether nystatin is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when nystatin is administered to a nursing woman.

Pediatric Use

See **DOSAGE AND ADMINISTRATION**.

ADVERSE REACTIONS

Nystatin is well tolerated even with prolonged therapy. Oral irritation and sensitization have been reported. (See **PRECAUTIONS: General**).

Gastrointestinal: Diarrhea (including one case of bloody diarrhea), nausea, vomiting, gastrointestinal upset/disturbances.

Dermatologic: Rash, including urticaria has been reported rarely. Stevens-Johnson syndrome has been reported very rarely.

Other: Tachycardia, bronchospasm, facial swelling, and non-specific myalgia have also been rarely reported.

To report SUSPECTED ADVERSE REACTIONS, contact Kesin Pharma at 1-833-537-4679 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

OVERDOSAGE

Oral doses of nystatin in excess of five million units daily have caused nausea and gastrointestinal upset. There have been no reports of serious toxic effects of superinfections (see **CLINICAL PHARMACOLOGY, Pharmacokinetics**).

DOSAGE AND ADMINISTRATION

INFANTS: 2 mL (200,000 units) four times daily (in infants and young children, use dropper to place one-half of dose in each side of mouth and avoid feeding for 5 to 10 minutes).

NOTE: Limited clinical studies in premature and low birth weight infants indicate that 1 mL four times daily is effective.

CHILDREN AND ADULTS: 4 to 6 mL (400,000 to 600,000 units) four times daily (one-half of dose in each side of mouth). The preparation should be retained in the mouth as long as possible before swallowing.

Continue treatment for at least 48 hours after perioral symptoms have disappeared and cultures demonstrate eradication of *Candida albicans*.

HOW SUPPLIED

Nystatin Oral Suspension, USP 100,000 USP Nystatin Units per mL, is available as a cherry-mint flavored, yellow, ready-to-use suspension, supplied in the following oral dosage forms:

NDC 81033-015-05: 5 mL unit-dose cup

NDC 81033-015-01: Case containing 100 unit-dose cups of 5 mL each

NDC 81033-015-54: Carton containing 50 unit-dose cups of 5 mL each

Rx Only

STORAGE AND HANDLING

SHAKE WELL BEFORE USE

Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15° C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature].

AVOID FREEZING

Pharmacists: Dispense in a tight, light-resistant container as defined in the USP.

Distributed by:

Kesin Pharma

Oldsmar, FL

34677

Revised: 11/2025

PI-015-V01



PRINCIPAL DISPLAY PANEL

NDC: 81033-015-54

Nystatin Oral Suspension, USP

500,000 units per 5 mL

Delivers 5 mL

50 x 5 mL

NDC: 81033-015-54

Nystatin Oral Suspension, USP

500,000 units per 5 mL

Delivers 5 mL



(01) 00381033015545

(21) 0000000000000000

(17) 00-NOV-0000

(10) K0000000

Rx Only

Packaged by and Distributed by
Kesin Pharma Corporation - Oldsmar, FL

FOR INSTITUTIONAL USE ONLY

Nystatin Oral Suspension, USP

500,000 units per 5 mL

50 x 5 mL Unit Dose Cups

EXP: 00-NOV-0000 LOT: K0000000

Store at 68°F to 77°F (20°C to 25°C)



See Insert
Internal Use Only R.01



0000000000000000

PRINCIPAL DISPLAY PANEL

NDC: 81033-015-01

Nystatin Oral Suspension, USP

500,000 units per 5 mL

Delivers 5 mL

Case = 100 UD Cups (Do Not Break Case)

NDC: 81033-015-01

Product Name: **Nystatin Oral Suspension, USP**
500,000 units per 5 mL



Rx Only

Delivers: 5 mL Case = 100 UD Cups (Do Not Break Case)

Storage: 68°F to 77°F (20°C to 25°C)

EXP: 00-NOV-0000 LOT: K0000000 QTY: 1



(17) 001100 (10) K0000000 (30) 1



(01) 003810331015019 (21) 00000000000000



Packaged by and Distributed by:
Kesin Pharma Corporation - Oldsmar, FL

Internal Use Only R.01



00000000000000

NYSTATIN

nystatin suspension

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:81033-015
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NYSTATIN (UNII: BDF1O1C72E) (NYSTATIN - UNII: BDF1O1C72E)	NYSTATIN	100000 U in 1 mL

Inactive Ingredients

Ingredient Name	Strength
EDETATE DISODIUM (UNII: 7FLD91C86K)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
HEXASODIUM HEXAMETAPHOSPHATE (UNII: N40N91DW96)	
SODIUM PHOSPHATE, DIBASIC, HEPTAHYDRATE (UNII: 70WT22SF4B)	
SODIUM PHOSPHATE, MONOBASIC, MONOHYDRATE (UNII: 593YOG76RN)	

GLYCERIN (UNII: PDC6A3C0OX)				
METHYLPARABEN (UNII: A2I8C7HI9T)				
PROPYLPARABEN (UNII: Z8IX2SC1OH)				
SUCROSE (UNII: C151H8M554)				
PEPPERMINT OIL (UNII: AV092KU4JH)				
Product Characteristics				
Color	yellow	Score		
Shape		Size		
Flavor	CHERRY (MINT)	Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:81033-015-01	100 in 1 CASE	11/24/2025	
1	NDC:81033-015-05	5 mL in 1 CUP, UNIT-DOSE; Type 0: Not a Combination Product		
2	NDC:81033-015-54	50 in 1 CARTON	11/24/2025	
2	NDC:81033-015-05	5 mL in 1 CUP, UNIT-DOSE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA062832	08/14/2020		

Labeler -
Kesin Pharma Corporation (117447816)

Establishment			
Name	Address	ID/FEI	Business Operations
Kesin Pharma Corporation		117447816	label(81033-015) , pack(81033-015)