

STOMACH RELIEF- bismuth subsalicylate liquid
Shield Pharmaceuticals Corp

Stomach Relief

Drug Facts

Active ingredient (in each 30 mL Dose)

Bismuth subsalicylate 525 mg

Purpose

Upset stomach reliever and antidiarrheal

Uses

relieves

- traveler's diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
- heartburn
- indigestion
- nausea
- gas
- belching
- fullness

Warnings

Reye's syndrome:

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- shake well before use
- only use dose cup provided
- adults and children 12 years and over:
- 30 mL (1 dose) every $\frac{1}{2}$ hour or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea
- 30mL (1 dose) every $\frac{1}{2}$ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (240 mL) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- **each 30 mL dose contains:** magnesium 3 mg, sodium 4 mg
- salicylate 261 mg
- low sodium
- protect from freezing
- avoid excessive heat (over 104°F or 40°C)

TAMPER EVIDENT: Do not use if printed shrinkband is missing or broken.

Inactive ingredients

FD&C red #33, FD&C yellow #6, gelatin, glycerin, lecithin, polyethylene glycol-400, polysorbate 20, polysorbate 80, povidone, purified water, sorbitan, sorbitol

solution, titanium dioxide

Questions?

1-800-373-6981

Distributed by:

Shield Pharmaceuticals Corp. Ronkonkoma, NY 11779

PRINCIPAL DISPLAY PANEL

Label

Your Brand

Compare to the active ingredient in Pepto-Bismol® Liquicaps™ NDC 0000-000-0

Stomach Relief

Bismuth Subsalicylate

Upset Stomach Reliever/
Antidiarrheal

Soothing Relieves of 5 Symptoms:

- Nausea
- Heartburn
- Indigestion
- Upset Stomach
- Diarrhea

4 fl. oz. (118 mL)

TAMPER EVIDENT: DO NOT USE IF SAFETY SEAL AROUND THE BOTTLE OR UNDER THE CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each 30 mL dose) Purpose

Bismuth subsalicylate 525 mg Upset stomach reliever and anti-diarrheal

Uses

■ relieves: ■ travelers' diarrhea ■ diarrhea ■ upset stomach due to overindulgence in food and drink, including: ■ heartburn ■ indigestion ■ nausea ■ gas ■ belching ■ fullness

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Drug Facts (continued under label)

This product is not manufactured or distributed by Procter & Gamble, or any of the registered trademarks Pepto-Bismol® Liquicaps™.

Made in India

LOT: EXP:

Drug Facts (continued under label)

Distributed by:
Shield Pharmaceuticals
Ronkonkoma, NY 11779

Directions

■ shake well before use
■ only use dose cup provided
adults and children 12 years and over:
■ 30 mL (1 dose) every $\frac{1}{2}$ hour or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea
■ 30 mL (1 dose) every $\frac{1}{2}$ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
■ do not exceed 8 doses (240 mL) in 24 hours
■ use until diarrhea stops but not more than 2 days
■ children under 12 years, ask a doctor
■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

■ **each 30 mL dose contains:** magnesium 3 mg, sodium 4 mg
■ salicylate 261 mg ■ low sodium ■ protect from freezing
■ avoid excessive heat (over 104°F or 40°C)

TAMPER EVIDENT: Do not use if printed shrinkband is missing or broken.

Inactive ingredients FD&C red #33, FD&C yellow #6, gelatin, glycerin, lecithin, polyethylene glycol 400, polysorbate 20, polysorbate 80, povidone, purified water, sorbitan, sorbitol solution, titanium dioxide

Questions or comments? 1-800-373-6981

Drug Facts (continued)

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use if you have

- an ulcer ■ a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever ■ mucus in the stool
- the stool and/or tongue may occur

Ask a doctor or pharmacist before use if you are taking any drug for

- anticoagulation (thinning the blood) ■ diabetes ■ gout
- arthritis

When using this product

- a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 3 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

■ shake well before use
■ only use dose cup provided
adults and children 12 years and over:
■ 30 mL (1 dose) every $\frac{1}{2}$ hour or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea
■ 30 mL (1 dose) every $\frac{1}{2}$ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
■ do not exceed 8 doses (240 mL) in 24 hours
■ use until diarrhea stops but not more than 2 days
■ children under 12 years, ask a doctor
■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

■ **each 30 mL dose contains:** magnesium 3 mg, sodium 4 mg
■ salicylate 261 mg ■ low sodium ■ protect from freezing
■ avoid excessive heat (over 104°F or 40°C)

TAMPER EVIDENT: Do not use if printed shrinkband is missing or broken.

Inactive ingredients FD&C red #33, FD&C yellow #6, gelatin, glycerin, lecithin, polyethylene glycol 400, polysorbate 20, polysorbate 80, povidone, purified water, sorbitan, sorbitol solution, titanium dioxide

Questions or comments? 1-800-373-6981

STOMACH RELIEF

bismuth subsalicylate liquid**Product Information**

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDc:83059-0126
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ, BISMUTH CATION - UNII:ZS9CD1I8YE)	BISMUTH SUBSALICYLATE	262 mg in 118 mL

Inactive Ingredients

Ingredient Name	Strength
FD&C RED NO. 3 (UNII: PN2ZH5LOQY)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C00X)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POLYSORBATE 20 (UNII: 7T1F30V5YH)	
POLYSORBATE 80 (UNII: 60ZP39ZG8H)	
WATER (UNII: 059QF0KOOR)	
POVIDONE (UNII: FZ989GH94E)	
SORBITAN (UNII: 6092ICV9RU)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	pink	Score	
Shape		Size	
Flavor	WINTERGREEN	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDc:83059-0126-4	118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/26/2026	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date

Labeler - Shield Pharmaceuticals Corp (118724924)

Revised: 1/2026

Shield Pharmaceuticals Corp