

**PAIN RELIEF- acetaminophen tablet**  
**Meijer Distribution Inc**

-----  
**Meijer Distribution, Inc. Pain Relief Drug Facts**

**Active ingredient (in each tablet)**

Acetaminophen 325 mg

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
- the common cold
- headache
- minor pain of arthritis
- backache
- muscular aches
- toothache
- premenstrual and menstrual cramps
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or

pharmacist.

- if the user has ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if the user**

- has liver disease
- is a child with pain of arthritis

### **Ask a doctor or pharmacist before use if the user is**

taking the blood thinning drug warfarin

### **Stop use and ask a doctor if**

- pain gets worse or lasts for more than 10 days (for adults) or 5 days (for children)
- fever gets worse or lasts for more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

### **Directions**

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none"><li>• take 2 tablets every 4 to 6 hours while symptoms last</li><li>• do not take more than 10 tablets in 24 hours</li><li>• do not use for more than 10 days unless directed by a doctor</li></ul>
children 6-11 years	<ul style="list-style-type: none"><li>• take 1 tablet every 4 to 6 hours while symptoms last</li><li>• do not take more than 5 tablets in 24 hours</li><li>• do not use for more than 5 days unless directed by a doctor</li></ul>
children under 6 years	ask a doctor

### **Inactive ingredients**

croscarmellose sodium\*, povidone, pregelatinized starch, stearic acid \*may contain this ingredient

**Questions or comments?**

**1-800-719-9260**

**Principal Display Panel**

meijer®

REGULAR STRENGTH

Pain Relief

COMPARE TO TYLENOL® ACTIVE INGREDIENT

ACETAMINOPHEN

PAIN RELIEVER | FEVER REDUCER

For Ages 6 Years And Over

Actual Size

100 TABLETS

325 mg each



**Drug Facts**

**Active ingredient (in each tablet)**  
Acetaminophen 325 mg... Pain reliever/fever reducer

**Uses**  
temporarily relieves minor aches and pains due to:  
the common cold  
minor pain of arthritis  
muscular aches  
toothache  
menstrual and menstrual cramps  
temporarily reduces fever

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if:  
adult takes more than 4,000 mg of acetaminophen in 24 hours  
child takes more than 5 doses in 24 hours  
taken with other drugs containing acetaminophen  
adult has 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:  
skin redness  
hives  
rash  
If a skin reaction occurs, stop use and seek medical help right away.  
**Do not use**  
with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
if the user has ever had an allergic reaction to this product or any of its ingredients

40378 6E C4

DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING

8 49086 98066 2  
1 3733 98066 2

PID 2855830

**Drug Facts (continued)**

Ask a doctor before use if the user:  
has liver disease  
is a child with pain of arthritis  
Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin  
Stop use and ask a doctor if:  
new symptoms occur  
pain gets worse or lasts more than 10 days (for adults) or 5 days (for children)  
fever gets worse or lasts more than 3 days  
redness or swelling is present  
These could be signs of a serious condition.  
If pregnant or breast-feeding, ask a health professional before use.  
Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

do not take more than directed (see overdose warning)

adults and children 12 years and over	take 2 tablets every 4 to 6 hours while symptoms last do not take more than 10 tablets in 24 hours do not use for more than 10 days unless directed by a doctor
children 6-11 years	take 1 tablet every 4 to 6 hours while symptoms last do not take more than 5 tablets in 24 hours do not use for more than 5 days unless directed by a doctor
children under 6 years	ask a doctor

**Drug Facts (continued)**  
**Inactive ingredients** croscarmellose sodium, povidone, pregelatinized starch, stearic acid \*may contain this ingredient  
**Questions or comments?**  
1-800-71-9-9280  
Store at 20-25°C (68-77°F).  
This product is not manufactured or distributed by KeVive Inc., owner of the registered trademark "Tylenol".  
DIST. BY MEIER DISTRIBUTION, INC., GRAND RAPIDS, MI 49544  
www.meier.com  
www.meier.com/our-quality-guarantee  
satisfaction

meijer. REGULAR STRENGTH Pain Relief ACETAMINOPHEN PAIN RELIEVER FEVER REDUCER

COMPARE TO TYLENOL® ACTIVE INGREDIENT\*\*

NDC 41250-403-78

100 TABLETS 325 mg each Actual Size

meijer. REGULAR STRENGTH Pain Relief ACETAMINOPHEN PAIN RELIEVER | FEVER REDUCER

COMPARE TO TYLENOL® ACTIVE INGREDIENT\*\*

NDC 41250-403-78

100 TABLETS 325 mg each Actual Size

For Ages 6 Years And Over

meijer. REGULAR STRENGTH Pain Relief ACETAMINOPHEN PAIN RELIEVER FEVER REDUCER

COMPARE TO TYLENOL® ACTIVE INGREDIENT\*\*

NDC 41250-403-78

100 TABLETS 325 mg each Actual Size

meijer. REGULAR STRENGTH Pain Relief ACETAMINOPHEN PAIN RELIEVER | FEVER REDUCER

COMPARE TO TYLENOL® ACTIVE INGREDIENT\*\*

NDC 41250-403-78

100 TABLETS 325 mg each Actual Size

For Ages 6 Years And Over

**PAIN RELIEF**  
acetaminophen tablet

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-403
Route of Administration	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

**Inactive Ingredients**

Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

**Product Characteristics**

Color	WHITE	Score	no score
Shape	ROUND (beveled edge)	Size	10mm
Flavor		Imprint Code	325MG;L403
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-403-78	1 in 1 CARTON	06/17/2024	
1		100 in 1 BOTTLE; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	06/17/2024	

**Labeler** - Meijer Distribution Inc (006959555)

Revised: 6/2024

Meijer Distribution Inc