

GLENMAX PEB DM FORTE- dextromethorphan hydrobromide, phenylephrine hydrochloride, and brompheniramine maleate syrup
Glendale Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Glenmax PEB DM FORTE

Drug Facts

| Active ingredients (in each teaspoonful) | Purpose |
|---|---------------------------------|
| Brompheniramine Maleate 4 mg | Antihistamine |
| Dextromethorphan Hydrobromide 20 mg | Antitussive (cough suppressant) |
| Phenylephrine Hydrochloride 10 mg | Nasal Decongestant |

Uses

temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- cough due to minor throat and bronchial irritation
- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes
- nasal congestion
- temporarily restores freer breathing through the nose congestion
- reduces swelling of nasal passages

Warnings

Do not exceed recommended dosage.

Do not use this product

- to sedate a child or make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- a cough that lasts or is chronic such as occurs with smoking, asthma or emphysema
- a cough that occur with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking any other nasal decongestant or stimulant
- taking sedatives or tranquilizers

When using this product

- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- cough or nasal congestion persists for more than 1 week, tends to recur, or is accompanied by a fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions**Do not exceed 4 dosage in a 24-hour period.**

| | |
|---|-----------------------------|
| Adults and children 12 years of age and over: | 1 teaspoonful every 4 hours |
| Children under 12 years of age: | Consult a physician |

Other information

Store at 59°-86°F (15°-30°C) [see USP for Controlled Room Temperature]

Inactive ingredients

Fruit gum flavor, citric acid, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol.

Questions? Comments?

To report a serious adverse event or obtain product information, Call 1-630-530-7000.

Distributed by:

Glendale Inc

Villa Park, IL 60181

PRINCIPAL DISPLAY PANEL - 473 mL Bottle Label

NDC 70147-0223-16

Glenmax

PEB DM FORTE

Antihistamine/Antitussive

Nasal Decongestant

**Each teaspoonful for oral
administration contains:**

Brompheniramine Maleate 4 mg

Dextromethorphan HBr 20 mg

Phenylephrine HCl 10 mg

SUGAR FREE / DYE FREE

ALCOHOL FREE

Fruit Gum Flavored Liquid

Tamper evident by foil seal under cap.

Do not use if foil seal is broken or missing.

Distributed by:

Glendale Inc

Villa Park, IL 60181

16 fl oz. (473 mL)

NDC 70147-0223-16

Glenmax

PEB DM FORTE

Antihistamine/Antitussive
Nasal Decongestant

Each teaspoonful for oral administration contains:

Brompheniramine Maleate..... 4 mg
Dextromethorphan HBr20 mg
Phenylephrine HCl10 mg

**SUGAR FREE / DYE FREE
ALCOHOL FREE**

Fruit Gum Flavored Liquid

Tamper evident by foil seal under cap.
Do not use if foil seal is broken or missing.



16 fl oz. (473 mL)

Drug Facts

Active ingredients (in each teaspoonful)

| | Purpose |
|-------------------------------------|---------------------------------|
| Brompheniramine Maleate 4 mg | Antihistamine |
| Dextromethorphan Hydrobromide 20 mg | Antitussive (cough suppressant) |
| Phenylephrine Hydrochloride 10 mg | Nasal Decongestant |

Uses temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- cough due to minor throat and bronchial irritation
- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes
- nasal congestion
- temporarily restores freer breathing through the nose
- reduces swelling of nasal passages

Warnings

Do not exceed recommended dosage.

Do not use this product

- to sedate a child or make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- a cough that lasts or is chronic such as occurs with smoking, asthma or emphysema
- a cough that occur with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Do not exceed 4 dosage in a 24-hour period.

| | |
|---|-----------------------------|
| Adults and children 12 years of age and over: | 1 teaspoonful every 4 hours |
| Children under 12 years of age: | Consult a physician |

Other information

Store at 59°-86°F (15°-30°C) [see USP for Controlled Room Temperature]

Inactive ingredients

Fruit gum flavor, citric acid, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol.

Questions? Comments?

To report a serious adverse event or obtain product information, Call 1-630-530-7000.

ASK a doctor or pharmacist before use if you are

■ taking any other nasal decongestant or stimulant ■ taking sedatives or tranquilizers

When using this product

■ marked drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery ■ excitability may occur, especially in children

Stop use and ask a doctor if

■ nervousness, dizziness, or sleeplessness occur ■ cough or nasal congestion persists for more than 1 week, tends to recur, or is accompanied by a fever, rash, or persistent headache. These could be signs of a serious condition.

Rev. 10/15

GLENMAX PEB DM FORTE

dextromethorphan hydrobromide, phenylephrine hydrochloride, and brompheniramine maleate syrup

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|----------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:70 147-223 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|----------------------------------|------------------|
| DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 20 mg in 5 mL |
| PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV) | PHENYLEPHRINE HYDROCHLORIDE | 10 mg in 5 mL |
| BROMPHENIRAMINE MALEATE (UNII: IXA7C9ZN03) (BROMPHENIRAMINE - UNII:H57G17P2FN) | BROMPHENIRAMINE MALEATE | 4 mg in 5 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| WATER (UNII: 059QF0KO0R) | |
| SACCHARIN SODIUM (UNII: SB8ZUX40TY) | |
| SODIUM BENZOATE (UNII: OJ245FE5EU) | |
| SORBITOL (UNII: 506T60A25R) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |

Product Characteristics

| | | | |
|-----------------|-------|---------------------|--|
| Color | | Score | |
| Shape | | Size | |
| Flavor | FRUIT | Imprint Code | |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|------------------------------|---------------------|--|----------------------|--------------------|
| 1 | NDC:70 147-223-16 | 473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | | |
| Marketing Information | | | | |
| | Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
| | OTC MONOGRAPH FINAL | part341 | 12/05/20 15 | |

Labeler - Glendale Inc (079987961)

Revised: 12/2015

Glendale Inc