

ACETAMINOPHEN- acetaminophen tablet
Grocery Outlet, Inc.

1001-GBR-2026-0218

Drug Facts

Active ingredient (in each tablet)

Acetaminophen 325 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If

you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if the user is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if the user has liver disease.

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days in adults
- pain gets worse or lasts more than 5 days in children under 12 years
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none">▪ take 2 tablets every 4 to 6 hours while symptoms last▪ do not take more than 10 tablets in 24 hours▪ do not use for more than 10 days unless directed by a doctor
children 6-11 years	<ul style="list-style-type: none">▪ take 1 tablet every 4 to 6 hours while symptoms last▪ do not take more than 5 tablets in 24 hours▪ do not use for more than 5 days unless directed by a doctor
children under 6 years	<ul style="list-style-type: none">▪ ask a doctor

Other information

- store between 20-25°C (68-77°F)
- retain carton for complete product information

Inactive ingredients

povidone, pregelatinized starch, sodium starch glycolate, stearic acid

Questions or comments?

1-844-705-4384

PRINCIPAL DISPLAY PANEL

NDC 85828-102-02

Compare to the active ingredient in Tylenol® Regular Strength Tablets*

REGULAR STRENGTH

Acetaminophen

TABLETS, 325 mg

Pain Reliever / Fever Reducer

Contains No Aspirin

GO

Actual Size

100 TABLETS

INK AND COATING FREE
FOR LOT AND
EXPIRATION STAMPING



**DO NOT USE IF IMPRINTED SEAL
UNDER CAP IS BROKEN OR MISSING**

Consumer HealthCare, distributor of Tylenol Regular Strength Tablets. This product is not to be used or dispensed in any manner.

Drug Facts

Active Ingredient (in each tablet)
Acetaminophen 325 mg.....Pain reliever/fever reducer

Purpose

Uses ■ temporarily relieves minor aches and pains due to:
■ the common cold ■ headache ■ backache
■ minor pain of arthritis ■ toothache ■ muscular aches
■ premenstrual and menstrual cramps
■ temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if:
■ adult takes more than 4,000 mg of acetaminophen in 24 hours
■ child takes more than 5 doses in 24 hours, which is the maximum daily amount
■ taken with other drugs containing acetaminophen
■ adult has 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin redness ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if the user is allergic to acetaminophen or any of the inactive ingredients in this product
Ask a doctor before use if the user has liver disease. ▶

Drug Facts (continued)

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin

Stop use and ask a doctor if
■ pain gets worse or lasts more than 10 days in adults
■ pain gets worse or lasts more than 5 days in children under 12 years
■ fever gets worse or lasts more than 3 days
■ new symptoms occur ■ redness or swelling is present
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ do not take more than directed (see overdose warning)
adults and children 12 years and over
■ take 2 tablets every 4 to 6 hours while symptoms last
■ do not take more than 10 tablets in 24 hours
■ do not use for more than 10 days unless directed by a doctor
children 6-11 years
■ take 1 tablet every 4 to 6 hours while symptoms last
■ do not take more than 5 tablets in 24 hours
■ do not use for more than 5 days unless directed by a doctor
children under 6 years
■ ask a doctor

Drug Facts (continued)

Other information

■ store between 20-25°C (68-77°F)
■ retain carton for complete product information

Inactive ingredients powders, pregelatinized starch, sodium starch glycolate, stearic acid

Questions or comments? 1-844-705-4384

DISTRIBUTED BY:
Greener Queer, Inc.
Emeryville, CA 94608
www.greenerqueer.com



NDC 85828-102-02
Compare to active ingredient in Tylenol® Regular Strength Tablets*

**REGULAR STRENGTH
Acetaminophen**

TABLETS, 325 mg
Pain Reliever / Fever Reducer
Contains No Aspirin



NDC 85828-102-02
Compare to active ingredient in Tylenol® Regular Strength Tablets*

**REGULAR STRENGTH
Acetaminophen**

TABLETS, 325 mg
Pain Reliever / Fever Reducer
Contains No Aspirin



ACETAMINOPHEN

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:85828-102
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

Inactive Ingredients

Ingredient Name	Strength
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	10mm
Flavor		Imprint Code	M2A3;57344
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:85828-102-02	1 in 1 CARTON	02/18/2026	
1		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/18/2026	

Labeler - Grocery Outlet, Inc. (029161585)

Revised: 2/2026

Grocery Outlet, Inc.