

IBUPROFEN MINIS- ibuprofen capsule, liquid filled
Grocery Outlet, Inc.

Grocery Outlet Ibuprofen Capsules 200 mg (Mini liquid capsules)

Drug Facts

Active ingredient (in each capsule)

Solubilized ibuprofen equal to
200 mg ibuprofen (NSAID)*
(present as the free acid and potassium salt)
*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/Fever reducer

Uses

-temporarily relieves minor aches and pains due to:

- headache
- toothache
- backache
- menstrual cramps
- the common cold
- muscular aches
- minor pain of arthritis

-temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

take with food or milk if stomach upset occurs

Stop use and ask a doctor if

-you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better

-you have symptoms of heart problems or stroke:

- chest pain
- trouble breathing

- weakness in one part or side of body
- slurred speech
- leg swelling

-pain gets worse or lasts more than 10 days
-fever gets worse or lasts more than 3 days
-redness or swelling is present in the painful area
-any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

-

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- each capsule contains: **potassium 20 mg**
- read all warnings and directions before use
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F).

Inactive ingredients

ammonium hydroxide, FD&C green no.3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, potassium hydroxide, propylene glycol, purified water, shellac, sorbitan monooleate, sorbitol sorbitan

Questions or comments?

Call **1-877-290-4008**

Ibuprofen Minis

SMALLER CAPSULES
SAME STRENGTH

SOLUBILIZED IBUPROFEN CAPSULES, 200 mg
Pain Reliever / Fever Reducer (NSAID)

GO

80 SOFTGELS¹
¹Liquid filled capsules

NDC 85828-700-80

DO NOT USE IF SEAL UNDER BOTTLE CAP IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING.

RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Drug Facts

Active Ingredient (in each capsule) Purpose
Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)¹ Pain reliever/fever reducer (present as the free acid and potassium salt)
nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to:
■ headache ■ toothache ■ backache ■ menstrual cramps
■ the common cold ■ muscular aches ■ minor pain of arthritis
Warnings
temporarily reduces fever

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock
■ skin reddening ■ rash ■ blisters
If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug
NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
■ if pain or fever does not respond to 1 capsule, 2 capsules may be used by a doctor
■ children under 12 years: ask a doctor

Other information ■ each capsule contains: potassium 20 mg ■ read all warnings and directions before use ■ store at 20-25°C (68-77°F) ■ avoid excessive heat above 40°C (104°F)
Questions or comments? Call 1-877-290-4008

DISTRIBUTED BY:
Grocery Outlet, Inc., Emeryville, CA 94608
www.groceryoutlet.com
Made in India. 700T 0126

LOT: GIM005
COUNT 100%
ON US MONEY BACK GUARANTEE

Varnish Omit Area

LOT:
EXP.:

Varnish Omit

Drug Facts (continued)

Do not use
■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
■ right before or after heart surgery

Ask a doctor before use if
■ stomach bleeding warning applies to you
■ you have problems or serious side effects from taking pain relievers or fever reducers
■ you have a history of stomach problems, such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are
■ under a doctor's care for any serious condition
■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
■ taking any other drug

When using this product
■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if
■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood ■ have bloody or black stools
■ have stomach pain that does not get better
■ you have symptoms of heart problems or stroke:
■ chest pain ■ trouble breathing
■ weakness in one part or side of body
■ slurred speech ■ leg swelling
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present in the painful area
■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 30 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Varnish Omit

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
■ do not take more than directed
■ the smallest effective dose should be used
■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
■ if pain or fever does not respond to 1 capsule, 2 capsules may be used
■ do not exceed 6 capsules in 24 hours, unless directed by a doctor
■ children under 12 years: ask a doctor

Other information
■ each capsule contains: potassium 20 mg
■ read all warnings and directions before use
■ store at 20-25°C (68-77°F)
■ avoid excessive heat above 40°C (104°F)

Inactive ingredients
ammonium hydroxide, FD&C green no. 3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, potassium hydroxide, propylene glycol, purified water, shellac, sorbitan monooleate, sorbitol sorbitan

Questions or comments? Call 1-877-290-4008

¹This product is not manufactured or distributed by Haleon US Holdings LLC, owner of the registered trademark Advil[®] and Cabotant Pharma Solutions, Inc., owner of the registered trademark Liqui-Gels[®].

DISTRIBUTED BY:
Grocery Outlet, Inc.
Emeryville, CA 94608
www.groceryoutlet.com
Made in India.
700T 0126

COUNT ON US! 100% MONEY BACK GUARANTEE

Ibuprofen Minis

SMALLER CAPSULES
SAME STRENGTH

SOLUBILIZED IBUPROFEN CAPSULES, 200 mg
Pain Reliever / Fever Reducer (NSAID)

NDC 85828-700-80
Compare to the active ingredient in Advil[®] Liqui-Gels[®] minis¹**

Ibuprofen Minis

SMALLER CAPSULES
SAME STRENGTH

SOLUBILIZED IBUPROFEN CAPSULES, 200 mg
Pain Reliever / Fever Reducer (NSAID)

GO

80 ACTUAL SIZE SOFTGELS¹
¹Liquid filled capsules

DO NOT USE IF SEAL UNDER BOTTLE CAP IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING.



LOT: Varnish Omit
EXP.:

RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Drug Facts

Active Ingredient (in each capsule) Purpose
Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)¹ Pain reliever/fever reducer (present as the free acid and potassium salt)
nonsteroidal anti-inflammatory drug

Uses
■ temporarily relieves minor aches and pains due to:
■ headache ■ toothache ■ backache
■ menstrual cramps ■ the common cold
■ muscular aches ■ minor pain of arthritis
■ temporarily reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters
If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

IBUPROFEN MINIS

ibuprofen capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:85828-700
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
SORBITAN (UNII: 6O92ICV9RU)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
AMMONIA (UNII: 5138Q19F1X)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
GELATIN (UNII: 2G86QN327L)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SORBITAN MONOOLEATE (UNII: 06XEA2VD56)	
SHELLAC (UNII: 46N107B71O)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
SORBITOL (UNII: 506T60A25R)	
WATER (UNII: 059QF0KO0R)	
FERROSO FERRIC OXIDE (UNII: XM0M87F357)	

Product Characteristics

Color	green (Sea green to light green)	Score	no score
Shape	OVAL	Size	13mm
Flavor		Imprint Code	62
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:85828-700-80	1 in 1 CARTON	03/05/2026	
1		80 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079205	03/05/2026	

Labeler - Grocery Outlet, Inc. (029161585)

Registrant - TIME CAP LABORATORIES, INC. (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		925822975	manufacture(85828-700)

Revised: 3/2026

Grocery Outlet, Inc.