

ANTI DIARRHEAL- loperamide hydrochloride tablet, film coated
Publix Super Markets Inc

Publix Super Markets, Inc. Anti Diarrheal Drug Facts

Active ingredient (in each caplet)

Loperamide HCl 2 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Heart alert: Taking more than directed can cause serious heart problems or death

Do not use

if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

- store at 68°-77°F (20°-25°C)
- see end panel for lot number and expiration date

Inactive ingredients

anhydrous lactose, carnauba wax, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch

Principal Display Panel

LOPERAMIDE HYDROCHLORIDE TABLETS, 2 mg

antidiarrheal

ACTUAL SIZE

Controls the symptoms of diarrhea

Anti-diarrheal

12 CAPLETS*

*CAPSULE-SHAPED TABLETS

Compare to Imodium® A-D active ingredient

SEE NEW WARNINGS

22453 63 C8

OPEN OTHER END

000005/04173

LOPERAMIDE HYDROCHLORIDE TABLETS, 2 mg



antidiarrheal

NDC 56062-224-53

LOPERAMIDE HYDROCHLORIDE TABLETS, 2 mg

antidiarrheal



Drug Facts

Active ingredient (in each caplet) Loperamide HCl 2 mg.....Anti-diarrheal

Use Controls symptoms of diarrhea, including Travelers' Diarrhea.

Warnings Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl. Heart alert: Taking more than directed can cause serious heart problems or death. Do not use if you have bloody or black stool.

Ask a doctor before use if you have ever ■ mucus in the stool ■ a history of liver disease ■ a history of abnormal heart rhythm.

Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product the dizziness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if ■ symptoms get worse ■ diarrhea lasts for more than 2 days ■ you get abdominal swelling or bloating. These may be signs of a serious condition.

If pregnant or breastfeeding, ask a health professional before use.

Drug Facts (continued)

Inactive ingredients anhydrous lactose, carnuba wax, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch.

Other information ■ store at 68°-77°F (20°-25°C) ■ see end panel for lot number and expiration date

Keep out of reach of children in case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea ■ find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over: 2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours

children 9-11 years (60-95 lbs) 1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours

children 6-8 years (48-59 lbs) 1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours

children 2-5 years (34 to 47 lbs) ask a doctor


children under 2 years (up to 3 lbs) do not use

DISTRIBUTED BY
PUBLIX SUPER MARKETS, INC.
3300 PUBLIX CORPORATE PARKWAY
LAKELAND, FL 33811
1-888-267-3037 publix.com
PUBLIX GUARANTEE: COMPLETE
SATISFACTION OR YOUR MONEY BACK
Publix.



DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN
**This product is not manufactured or distributed by the owner of the registered trademark Imodium®.





ANTI DIARRHEAL

loperamide hydrochloride tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:56062-224
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	

Product Characteristics

Color	GREEN	Score	2 pieces
Shape	OVAL	Size	10mm
Flavor		Imprint Code	L2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:56062-224-53	12 in 1 CARTON	03/19/2003	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:56062-224-62	24 in 1 CARTON	03/25/2003	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:56062-224-80	1 in 1 CARTON	01/13/2005	12/31/2021
3		96 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075232	03/19/2003	

Labeler - Publix Super Markets Inc (006922009)

Revised: 12/2025

Publix Super Markets Inc