

**DAYTIME AND NIGHTTIME COLD AND FLU- acetaminophen,  
dextromethorphan hbr, phenylephrine hcl, acetaminophen,  
dextromethorphan hbr, doxylamine succinate  
Grocery Outlet, Inc.**

-----  
**Grocery Outlet Daytime and Nighttime Cold & Flu Liquid**

***Nighttime Cold & Flu***

***Drug Facts***

***Active ingredients (in each 30 mL)***

Acetaminophen 650 mg  
Dextromethorphan HBr 30 mg  
Doxylamine succinate 12.5 mg

***Purpose***

Pain reliever/ fever reducer  
Cough suppressant  
Antihistamine

***Uses***

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

■ more than 4 doses in 24 hours, which is the maximum daily amount for this product.

■ with other drugs containing acetaminophen

■ 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions.

Symptoms may include:

■ skin reddening

■ blisters

■ rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more

than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Liver warning:**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

### **Allergy alert:**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

### **Sore throat warning:**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

### **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

### **When using this product**

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

- alcohol, sedatives, and tranquilizers may increase drowsiness

### **Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

### **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

### **When using this product**

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

### **Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

■ cough comes back or occurs with rash or headache that lasts.  
These could be signs of a serious condition.

**If pregnant or breast-feeding**,ask a health professional before use.

**Keep out of reach of children.**In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

---

adults & children 12 yrs & over	30 mL every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

---

### **Other information**

- **each 30 mL tablespoon contains:**sodium 52 mg
- store at no greater than 25°C and do not refrigerate

### **Inactive ingredients**

alcohol, anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium saccharin, sorbitol solution, sucralose, xanthan gum

### **Questions or comments?**

Call **1-877-290-4008**

### **Daytime Cold & Flu Drug Facts**

#### **Active ingredients (in each 15 mL)**

Acetaminophen 325 mg  
Dextromethorphan HBr 10 mg  
Phenylephrine HCl 5 mg

#### **Purpose**

Pain reliever/Fever reducer  
Cough suppressant

## Nasal decongestant

### **Uses**

temporarily relieves common cold/flu symptoms:

- nasal congestion
- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever

### **Warnings**

**Liver warning:**This product contains acetaminophen. Severe liver damage may occur if

■ adult takes more than 4 doses (30 mL each) in 24 hrs, which is the maximum daily amount for this product

■ child takes more than 4 doses (15 mL each) in 24 hrs, which is the maximum daily amount for this product

■ taken with other drugs containing acetaminophen

■ adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:**Acetaminophen may cause severe skin reactions.

Symptoms may include:

■ skin reddening

■ blisters

■ rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:**If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Liver warning:**

This product contains acetaminophen. Severe liver damage may occur if

- adults take more than 4 doses (30 mL each) in 24 hours, which is the maximum daily amount for this product
- child take more than 4 doses (15 mL each) in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

### **Allergy alert:**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

**Sore throat warning:**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin.

**When using this product,**

**do not use more than directed.**

**Stop use and ask a doctor if**

- You get nervous, dizzy, or sleepless.
- Pain, nasal congestion or cough get worse or last more than 5 days (children) or 7 days (adults)
- Fever gets worse or last more than 3 days
- Redness or swelling is present
- New symptoms occur
- Cough comes back, or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

## Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

---

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

---

### Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin.

### When using this product, do not use more than directed.

#### Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough get worse or last more than 5 days (children) and 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Other information**

- each 15 mL contains: sodium 56 mg
- store at no greater than 25°C and do not refrigerate.

**Inactive ingredients**

anhydrous citric acid, FD&C yellow no. 6, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium chloride, sodium citrate, sodium saccharin, sorbitol solution, sucralose, xanthan gum

**Questions or comments?**

Call **1-877-290-4008**

**PRINCIPAL DISPLAY PANEL - Convenience Pack**

<p>Compare to the active ingredients in Vicks® DayQuil® Cold &amp; Flu Liquid*</p> <h2>Daytime</h2> <p><b>COLD &amp; FLU</b></p> <p><b>Acetaminophen,</b> Dextromethorphan HBr, Phenylephrine HCl</p> <p>Headache, Fever, Sore Throat, Minor Aches &amp; Pains</p> <p>Cough</p> <p>Nasal Congestion, Sinus Pressure</p> <p>Alcohol Free</p>	<p>NDC XXXXX-XXXX-XXXX</p> <p>Compare to the active ingredients in Vicks® NyQuil® Cold &amp; Flu Cherry Liquid*</p> <h2>Nighttime</h2> <p><b>COLD &amp; FLU</b></p> <p><b>Acetaminophen,</b> Dextromethorphan HBr, Doxylamine succinate</p> <p>Headache, Fever, Sore Throat, Minor Aches &amp; Pains</p> <p>Cough</p> <p>Sneezing, Runny Nose</p> <p>Alcohol 10% Nighttime Relief Cherry</p>	<table border="1"> <thead> <tr> <th>Drug Facts</th> <th>Nighttime Cold &amp; Flu</th> <th>Drug Facts</th> <th>Daytime Cold &amp; Flu</th> </tr> </thead> <tbody> <tr> <td><b>Active ingredients (in each 30 mL)</b></td> <td>Acetaminophen 325 mg Dextromethorphan HBr 10 mg Phenylephrine HCl 5 mg</td> <td><b>Active ingredients (in each 15 mL)</b></td> <td>Acetaminophen 325 mg Dextromethorphan HBr 10 mg Phenylephrine HCl 5 mg</td> </tr> <tr> <td><b>Warnings</b></td> <td>Do not use if you are allergic to any of the ingredients. Do not use if you are pregnant or breastfeeding. Do not use if you are taking other medications that contain acetaminophen, dextromethorphan, or phenylephrine. Do not use if you are taking other medications that contain doxylamine. Do not use if you are taking other medications that contain pseudoephedrine. Do not use if you are taking other medications that contain chlorpheniramine. Do not use if you are taking other medications that contain diphenhydramine. Do not use if you are taking other medications that contain promethazine. Do not use if you are taking other medications that contain cetirizine. Do not use if you are taking other medications that contain loratadine. Do not use if you are taking other medications that contain fexofenadine. Do not use if you are taking other medications that contain levocetirizine. Do not use if you are taking other medications that contain desloratadine. Do not use if you are taking other medications that contain rupatadine. Do not use if you are taking other medications that contain ebastine. Do not use if you are taking other medications that contain cetirizine. Do not use if you are taking other medications that contain loratadine. Do not use if you are taking other medications that contain fexofenadine. Do not use if you are taking other medications that contain levocetirizine. Do not use if you are taking other medications that contain desloratadine. Do not use if you are taking other medications that contain rupatadine. Do not use if you are taking other medications that contain ebastine.</td> <td>Do not use if you are allergic to any of the ingredients. Do not use if you are pregnant or breastfeeding. Do not use if you are taking other medications that contain acetaminophen, dextromethorphan, or phenylephrine. Do not use if you are taking other medications that contain doxylamine. Do not use if you are taking other medications that contain pseudoephedrine. Do not use if you are taking other medications that contain chlorpheniramine. Do not use if you are taking other medications that contain diphenhydramine. Do not use if you are taking other medications that contain promethazine. Do not use if you are taking other medications that contain cetirizine. Do not use if you are taking other medications that contain loratadine. Do not use if you are taking other medications that contain fexofenadine. Do not use if you are taking other medications that contain levocetirizine. Do not use if you are taking other medications that contain desloratadine. Do not use if you are taking other medications that contain rupatadine. Do not use if you are taking other medications that contain ebastine.</td> <td>Do not use if you are allergic to any of the ingredients. Do not use if you are pregnant or breastfeeding. Do not use if you are taking other medications that contain acetaminophen, dextromethorphan, or phenylephrine. Do not use if you are taking other medications that contain doxylamine. Do not use if you are taking other medications that contain pseudoephedrine. Do not use if you are taking other medications that contain chlorpheniramine. Do not use if you are taking other medications that contain diphenhydramine. Do not use if you are taking other medications that contain promethazine. Do not use if you are taking other medications that contain cetirizine. Do not use if you are taking other medications that contain loratadine. Do not use if you are taking other medications that contain fexofenadine. Do not use if you are taking other medications that contain levocetirizine. Do not use if you are taking other medications that contain desloratadine. Do not use if you are taking other medications that contain rupatadine. Do not use if you are taking other medications that contain ebastine.</td> </tr> <tr> <td><b>Directions</b></td> <td>Take 15 mL every 4 to 6 hours as needed. Do not exceed 6 doses per 24 hours. Do not use if you are pregnant or breastfeeding. Do not use if you are taking other medications that contain acetaminophen, dextromethorphan, or phenylephrine. Do not use if you are taking other medications that contain doxylamine. Do not use if you are taking other medications that contain pseudoephedrine. Do not use if you are taking other medications that contain chlorpheniramine. Do not use if you are taking other medications that contain diphenhydramine. Do not use if you are taking other medications that contain promethazine. Do not use if you are taking other medications that contain cetirizine. Do not use if you are taking other medications that contain loratadine. Do not use if you are taking other medications that contain fexofenadine. Do not use if you are taking other medications that contain levocetirizine. Do not use if you are taking other medications that contain desloratadine. Do not use if you are taking other medications that contain rupatadine. Do not use if you are taking other medications that contain ebastine.</td> <td>Take 15 mL every 4 to 6 hours as needed. Do not exceed 6 doses per 24 hours. Do not use if you are pregnant or breastfeeding. Do not use if you are taking other medications that contain acetaminophen, dextromethorphan, or phenylephrine. Do not use if you are taking other medications that contain doxylamine. Do not use if you are taking other medications that contain pseudoephedrine. Do not use if you are taking other medications that contain chlorpheniramine. Do not use if you are taking other medications that contain diphenhydramine. Do not use if you are taking other medications that contain promethazine. Do not use if you are taking other medications that contain cetirizine. Do not use if you are taking other medications that contain loratadine. Do not use if you are taking other medications that contain fexofenadine. Do not use if you are taking other medications that contain levocetirizine. Do not use if you are taking other medications that contain desloratadine. Do not use if you are taking other medications that contain rupatadine. Do not use if you are taking other medications that contain ebastine.</td> <td>Take 15 mL every 4 to 6 hours as needed. Do not exceed 6 doses per 24 hours. Do not use if you are pregnant or breastfeeding. Do not use if you are taking other medications that contain acetaminophen, dextromethorphan, or phenylephrine. Do not use if you are taking other medications that contain doxylamine. Do not use if you are taking other medications that contain pseudoephedrine. Do not use if you are taking other medications that contain chlorpheniramine. Do not use if you are taking other medications that contain diphenhydramine. Do not use if you are taking other medications that contain promethazine. Do not use if you are taking other medications that contain cetirizine. Do not use if you are taking other medications that contain loratadine. Do not use if you are taking other medications that contain fexofenadine. Do not use if you are taking other medications that contain levocetirizine. Do not use if you are taking other medications that contain desloratadine. Do not use if you are taking other medications that contain rupatadine. Do not use if you are taking other medications that contain ebastine.</td> </tr> <tr> <td><b>Other information</b></td> <td>Do not use if you are pregnant or breastfeeding. Do not use if you are taking other medications that contain acetaminophen, dextromethorphan, or phenylephrine. Do not use if you are taking other medications that contain doxylamine. Do not use if you are taking other medications that contain pseudoephedrine. Do not use if you are taking other medications that contain chlorpheniramine. Do not use if you are taking other medications that contain diphenhydramine. Do not use if you are taking other medications that contain promethazine. Do not use if you are taking other medications that contain cetirizine. Do not use if you are taking other medications that contain loratadine. Do not use if you are taking other medications that contain fexofenadine. Do not use if you are taking other medications that contain levocetirizine. Do not use if you are taking other medications that contain desloratadine. Do not use if you are taking other medications that contain rupatadine. Do not use if you are taking other medications that contain ebastine.</td> <td>Do not use if you are pregnant or breastfeeding. Do not use if you are taking other medications that contain acetaminophen, dextromethorphan, or phenylephrine. Do not use if you are taking other medications that contain doxylamine. Do not use if you are taking other medications that contain pseudoephedrine. Do not use if you are taking other medications that contain chlorpheniramine. Do not use if you are taking other medications that contain diphenhydramine. Do not use if you are taking other medications that contain promethazine. Do not use if you are taking other medications that contain cetirizine. Do not use if you are taking other medications that contain loratadine. Do not use if you are taking other medications that contain fexofenadine. Do not use if you are taking other medications that contain levocetirizine. Do not use if you are taking other medications that contain desloratadine. Do not use if you are taking other medications that contain rupatadine. Do not use if you are taking other medications that contain ebastine.</td> <td>Do not use if you are pregnant or breastfeeding. Do not use if you are taking other medications that contain acetaminophen, dextromethorphan, or phenylephrine. Do not use if you are taking other medications that contain doxylamine. Do not use if you are taking other medications that contain pseudoephedrine. Do not use if you are taking other medications that contain chlorpheniramine. Do not use if you are taking other medications that contain diphenhydramine. Do not use if you are taking other medications that contain promethazine. Do not use if you are taking other medications that contain cetirizine. Do not use if you are taking other medications that contain loratadine. Do not use if you are taking other medications that contain fexofenadine. Do not use if you are taking other medications that contain levocetirizine. Do not use if you are taking other medications that contain desloratadine. Do not use if you are taking other medications that contain rupatadine. Do not use if you are taking other medications that contain ebastine.</td> </tr> </tbody> </table>	Drug Facts	Nighttime Cold & Flu	Drug Facts	Daytime Cold & Flu	<b>Active ingredients (in each 30 mL)</b>	Acetaminophen 325 mg Dextromethorphan HBr 10 mg Phenylephrine HCl 5 mg	<b>Active ingredients (in each 15 mL)</b>	Acetaminophen 325 mg Dextromethorphan HBr 10 mg Phenylephrine HCl 5 mg	<b>Warnings</b>	Do not use if you are allergic to any of the ingredients. Do not use if you are pregnant or breastfeeding. Do not use if you are taking other medications that contain acetaminophen, dextromethorphan, or phenylephrine. Do not use if you are taking other medications that contain doxylamine. Do not use if you are taking other medications that contain pseudoephedrine. Do not use if you are taking other medications that contain chlorpheniramine. Do not use if you are taking other medications that contain diphenhydramine. Do not use if you are taking other medications that contain promethazine. Do not use if you are taking other medications that contain cetirizine. Do not use if you are taking other medications that contain loratadine. Do not use if you are taking other medications that contain fexofenadine. Do not use if you are taking other medications that contain levocetirizine. Do not use if you are taking other medications that contain desloratadine. Do not use if you are taking other medications that contain rupatadine. Do not use if you are taking other medications that contain ebastine. Do not use if you are taking other medications that contain cetirizine. Do not use if you are taking other medications that contain loratadine. Do not use if you are taking other medications that contain fexofenadine. Do not use if you are taking other medications that contain levocetirizine. Do not use if you are taking other medications that contain desloratadine. Do not use if you are taking other medications that contain rupatadine. Do not use if you are taking other medications that contain ebastine.	Do not use if you are allergic to any of the ingredients. Do not use if you are pregnant or breastfeeding. Do not use if you are taking other medications that contain acetaminophen, dextromethorphan, or phenylephrine. Do not use if you are taking other medications that contain doxylamine. Do not use if you are taking other medications that contain pseudoephedrine. Do not use if you are taking other medications that contain chlorpheniramine. Do not use if you are taking other medications that contain diphenhydramine. Do not use if you are taking other medications that contain promethazine. Do not use if you are taking other medications that contain cetirizine. Do not use if you are taking other medications that contain loratadine. Do not use if you are taking other medications that contain fexofenadine. Do not use if you are taking other medications that contain levocetirizine. Do not use if you are taking other medications that contain desloratadine. Do not use if you are taking other medications that contain rupatadine. Do not use if you are taking other medications that contain ebastine.	Do not use if you are allergic to any of the ingredients. Do not use if you are pregnant or breastfeeding. Do not use if you are taking other medications that contain acetaminophen, dextromethorphan, or phenylephrine. Do not use if you are taking other medications that contain doxylamine. Do not use if you are taking other medications that contain pseudoephedrine. Do not use if you are taking other medications that contain chlorpheniramine. Do not use if you are taking other medications that contain diphenhydramine. Do not use if you are taking other medications that contain promethazine. Do not use if you are taking other medications that contain cetirizine. Do not use if you are taking other medications that contain loratadine. Do not use if you are taking other medications that contain fexofenadine. Do not use if you are taking other medications that contain levocetirizine. Do not use if you are taking other medications that contain desloratadine. Do not use if you are taking other medications that contain rupatadine. Do not use if you are taking other medications that contain ebastine.	<b>Directions</b>	Take 15 mL every 4 to 6 hours as needed. Do not exceed 6 doses per 24 hours. Do not use if you are pregnant or breastfeeding. Do not use if you are taking other medications that contain acetaminophen, dextromethorphan, or phenylephrine. Do not use if you are taking other medications that contain doxylamine. Do not use if you are taking other medications that contain pseudoephedrine. Do not use if you are taking other medications that contain chlorpheniramine. Do not use if you are taking other medications that contain diphenhydramine. Do not use if you are taking other medications that contain promethazine. Do not use if you are taking other medications that contain cetirizine. Do not use if you are taking other medications that contain loratadine. Do not use if you are taking other medications that contain fexofenadine. Do not use if you are taking other medications that contain levocetirizine. Do not use if you are taking other medications that contain desloratadine. Do not use if you are taking other medications that contain rupatadine. Do not use if you are taking other medications that contain ebastine.	Take 15 mL every 4 to 6 hours as needed. Do not exceed 6 doses per 24 hours. Do not use if you are pregnant or breastfeeding. Do not use if you are taking other medications that contain acetaminophen, dextromethorphan, or phenylephrine. Do not use if you are taking other medications that contain doxylamine. Do not use if you are taking other medications that contain pseudoephedrine. Do not use if you are taking other medications that contain chlorpheniramine. Do not use if you are taking other medications that contain diphenhydramine. Do not use if you are taking other medications that contain promethazine. Do not use if you are taking other medications that contain cetirizine. Do not use if you are taking other medications that contain loratadine. Do not use if you are taking other medications that contain fexofenadine. Do not use if you are taking other medications that contain levocetirizine. Do not use if you are taking other medications that contain desloratadine. Do not use if you are taking other medications that contain rupatadine. Do not use if you are taking other medications that contain ebastine.	Take 15 mL every 4 to 6 hours as needed. Do not exceed 6 doses per 24 hours. Do not use if you are pregnant or breastfeeding. Do not use if you are taking other medications that contain acetaminophen, dextromethorphan, or phenylephrine. Do not use if you are taking other medications that contain doxylamine. Do not use if you are taking other medications that contain pseudoephedrine. Do not use if you are taking other medications that contain chlorpheniramine. Do not use if you are taking other medications that contain diphenhydramine. Do not use if you are taking other medications that contain promethazine. Do not use if you are taking other medications that contain cetirizine. Do not use if you are taking other medications that contain loratadine. Do not use if you are taking other medications that contain fexofenadine. Do not use if you are taking other medications that contain levocetirizine. Do not use if you are taking other medications that contain desloratadine. Do not use if you are taking other medications that contain rupatadine. Do not use if you are taking other medications that contain ebastine.	<b>Other information</b>	Do not use if you are pregnant or breastfeeding. Do not use if you are taking other medications that contain acetaminophen, dextromethorphan, or phenylephrine. Do not use if you are taking other medications that contain doxylamine. Do not use if you are taking other medications that contain pseudoephedrine. Do not use if you are taking other medications that contain chlorpheniramine. Do not use if you are taking other medications that contain diphenhydramine. Do not use if you are taking other medications that contain promethazine. Do not use if you are taking other medications that contain cetirizine. Do not use if you are taking other medications that contain loratadine. Do not use if you are taking other medications that contain fexofenadine. Do not use if you are taking other medications that contain levocetirizine. Do not use if you are taking other medications that contain desloratadine. Do not use if you are taking other medications that contain rupatadine. Do not use if you are taking other medications that contain ebastine.	Do not use if you are pregnant or breastfeeding. Do not use if you are taking other medications that contain acetaminophen, dextromethorphan, or phenylephrine. Do not use if you are taking other medications that contain doxylamine. Do not use if you are taking other medications that contain pseudoephedrine. Do not use if you are taking other medications that contain chlorpheniramine. Do not use if you are taking other medications that contain diphenhydramine. Do not use if you are taking other medications that contain promethazine. Do not use if you are taking other medications that contain cetirizine. Do not use if you are taking other medications that contain loratadine. Do not use if you are taking other medications that contain fexofenadine. Do not use if you are taking other medications that contain levocetirizine. Do not use if you are taking other medications that contain desloratadine. Do not use if you are taking other medications that contain rupatadine. Do not use if you are taking other medications that contain ebastine.	Do not use if you are pregnant or breastfeeding. Do not use if you are taking other medications that contain acetaminophen, dextromethorphan, or phenylephrine. Do not use if you are taking other medications that contain doxylamine. Do not use if you are taking other medications that contain pseudoephedrine. Do not use if you are taking other medications that contain chlorpheniramine. Do not use if you are taking other medications that contain diphenhydramine. Do not use if you are taking other medications that contain promethazine. Do not use if you are taking other medications that contain cetirizine. Do not use if you are taking other medications that contain loratadine. Do not use if you are taking other medications that contain fexofenadine. Do not use if you are taking other medications that contain levocetirizine. Do not use if you are taking other medications that contain desloratadine. Do not use if you are taking other medications that contain rupatadine. Do not use if you are taking other medications that contain ebastine.
Drug Facts	Nighttime Cold & Flu	Drug Facts	Daytime Cold & Flu																			
<b>Active ingredients (in each 30 mL)</b>	Acetaminophen 325 mg Dextromethorphan HBr 10 mg Phenylephrine HCl 5 mg	<b>Active ingredients (in each 15 mL)</b>	Acetaminophen 325 mg Dextromethorphan HBr 10 mg Phenylephrine HCl 5 mg																			
<b>Warnings</b>	Do not use if you are allergic to any of the ingredients. Do not use if you are pregnant or breastfeeding. Do not use if you are taking other medications that contain acetaminophen, dextromethorphan, or phenylephrine. Do not use if you are taking other medications that contain doxylamine. Do not use if you are taking other medications that contain pseudoephedrine. Do not use if you are taking other medications that contain chlorpheniramine. Do not use if you are taking other medications that contain diphenhydramine. Do not use if you are taking other medications that contain promethazine. Do not use if you are taking other medications that contain cetirizine. Do not use if you are taking other medications that contain loratadine. Do not use if you are taking other medications that contain fexofenadine. Do not use if you are taking other medications that contain levocetirizine. Do not use if you are taking other medications that contain desloratadine. Do not use if you are taking other medications that contain rupatadine. Do not use if you are taking other medications that contain ebastine. Do not use if you are taking other medications that contain cetirizine. Do not use if you are taking other medications that contain loratadine. Do not use if you are taking other medications that contain fexofenadine. Do not use if you are taking other medications that contain levocetirizine. Do not use if you are taking other medications that contain desloratadine. Do not use if you are taking other medications that contain rupatadine. Do not use if you are taking other medications that contain ebastine.	Do not use if you are allergic to any of the ingredients. Do not use if you are pregnant or breastfeeding. Do not use if you are taking other medications that contain acetaminophen, dextromethorphan, or phenylephrine. Do not use if you are taking other medications that contain doxylamine. Do not use if you are taking other medications that contain pseudoephedrine. Do not use if you are taking other medications that contain chlorpheniramine. Do not use if you are taking other medications that contain diphenhydramine. Do not use if you are taking other medications that contain promethazine. Do not use if you are taking other medications that contain cetirizine. Do not use if you are taking other medications that contain loratadine. Do not use if you are taking other medications that contain fexofenadine. Do not use if you are taking other medications that contain levocetirizine. Do not use if you are taking other medications that contain desloratadine. Do not use if you are taking other medications that contain rupatadine. Do not use if you are taking other medications that contain ebastine.	Do not use if you are allergic to any of the ingredients. Do not use if you are pregnant or breastfeeding. Do not use if you are taking other medications that contain acetaminophen, dextromethorphan, or phenylephrine. Do not use if you are taking other medications that contain doxylamine. Do not use if you are taking other medications that contain pseudoephedrine. Do not use if you are taking other medications that contain chlorpheniramine. Do not use if you are taking other medications that contain diphenhydramine. Do not use if you are taking other medications that contain promethazine. Do not use if you are taking other medications that contain cetirizine. Do not use if you are taking other medications that contain loratadine. Do not use if you are taking other medications that contain fexofenadine. Do not use if you are taking other medications that contain levocetirizine. Do not use if you are taking other medications that contain desloratadine. Do not use if you are taking other medications that contain rupatadine. Do not use if you are taking other medications that contain ebastine.																			
<b>Directions</b>	Take 15 mL every 4 to 6 hours as needed. Do not exceed 6 doses per 24 hours. Do not use if you are pregnant or breastfeeding. Do not use if you are taking other medications that contain acetaminophen, dextromethorphan, or phenylephrine. Do not use if you are taking other medications that contain doxylamine. Do not use if you are taking other medications that contain pseudoephedrine. Do not use if you are taking other medications that contain chlorpheniramine. Do not use if you are taking other medications that contain diphenhydramine. Do not use if you are taking other medications that contain promethazine. Do not use if you are taking other medications that contain cetirizine. Do not use if you are taking other medications that contain loratadine. Do not use if you are taking other medications that contain fexofenadine. Do not use if you are taking other medications that contain levocetirizine. Do not use if you are taking other medications that contain desloratadine. Do not use if you are taking other medications that contain rupatadine. Do not use if you are taking other medications that contain ebastine.	Take 15 mL every 4 to 6 hours as needed. Do not exceed 6 doses per 24 hours. Do not use if you are pregnant or breastfeeding. Do not use if you are taking other medications that contain acetaminophen, dextromethorphan, or phenylephrine. Do not use if you are taking other medications that contain doxylamine. Do not use if you are taking other medications that contain pseudoephedrine. Do not use if you are taking other medications that contain chlorpheniramine. Do not use if you are taking other medications that contain diphenhydramine. Do not use if you are taking other medications that contain promethazine. Do not use if you are taking other medications that contain cetirizine. Do not use if you are taking other medications that contain loratadine. Do not use if you are taking other medications that contain fexofenadine. Do not use if you are taking other medications that contain levocetirizine. Do not use if you are taking other medications that contain desloratadine. Do not use if you are taking other medications that contain rupatadine. Do not use if you are taking other medications that contain ebastine.	Take 15 mL every 4 to 6 hours as needed. Do not exceed 6 doses per 24 hours. Do not use if you are pregnant or breastfeeding. Do not use if you are taking other medications that contain acetaminophen, dextromethorphan, or phenylephrine. Do not use if you are taking other medications that contain doxylamine. Do not use if you are taking other medications that contain pseudoephedrine. Do not use if you are taking other medications that contain chlorpheniramine. Do not use if you are taking other medications that contain diphenhydramine. Do not use if you are taking other medications that contain promethazine. Do not use if you are taking other medications that contain cetirizine. Do not use if you are taking other medications that contain loratadine. Do not use if you are taking other medications that contain fexofenadine. Do not use if you are taking other medications that contain levocetirizine. Do not use if you are taking other medications that contain desloratadine. Do not use if you are taking other medications that contain rupatadine. Do not use if you are taking other medications that contain ebastine.																			
<b>Other information</b>	Do not use if you are pregnant or breastfeeding. Do not use if you are taking other medications that contain acetaminophen, dextromethorphan, or phenylephrine. Do not use if you are taking other medications that contain doxylamine. Do not use if you are taking other medications that contain pseudoephedrine. Do not use if you are taking other medications that contain chlorpheniramine. Do not use if you are taking other medications that contain diphenhydramine. Do not use if you are taking other medications that contain promethazine. Do not use if you are taking other medications that contain cetirizine. Do not use if you are taking other medications that contain loratadine. Do not use if you are taking other medications that contain fexofenadine. Do not use if you are taking other medications that contain levocetirizine. Do not use if you are taking other medications that contain desloratadine. Do not use if you are taking other medications that contain rupatadine. Do not use if you are taking other medications that contain ebastine.	Do not use if you are pregnant or breastfeeding. Do not use if you are taking other medications that contain acetaminophen, dextromethorphan, or phenylephrine. Do not use if you are taking other medications that contain doxylamine. Do not use if you are taking other medications that contain pseudoephedrine. Do not use if you are taking other medications that contain chlorpheniramine. Do not use if you are taking other medications that contain diphenhydramine. Do not use if you are taking other medications that contain promethazine. Do not use if you are taking other medications that contain cetirizine. Do not use if you are taking other medications that contain loratadine. Do not use if you are taking other medications that contain fexofenadine. Do not use if you are taking other medications that contain levocetirizine. Do not use if you are taking other medications that contain desloratadine. Do not use if you are taking other medications that contain rupatadine. Do not use if you are taking other medications that contain ebastine.	Do not use if you are pregnant or breastfeeding. Do not use if you are taking other medications that contain acetaminophen, dextromethorphan, or phenylephrine. Do not use if you are taking other medications that contain doxylamine. Do not use if you are taking other medications that contain pseudoephedrine. Do not use if you are taking other medications that contain chlorpheniramine. Do not use if you are taking other medications that contain diphenhydramine. Do not use if you are taking other medications that contain promethazine. Do not use if you are taking other medications that contain cetirizine. Do not use if you are taking other medications that contain loratadine. Do not use if you are taking other medications that contain fexofenadine. Do not use if you are taking other medications that contain levocetirizine. Do not use if you are taking other medications that contain desloratadine. Do not use if you are taking other medications that contain rupatadine. Do not use if you are taking other medications that contain ebastine.																			

**DAYTIME AND NIGHTTIME COLD AND FLU**  
 acetaminophen, dextromethorphan hbr, phenylephrine hcl, acetaminophen, dextromethorphan hbr, doxylamine succinate kit

Product Information			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:85828-773
<b>Packaging</b>			

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:85828-773-02	1 in 1 BOTTLE; Type 0: Not a Combination Product	03/02/2026	

### Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE, PLASTIC	354 mL
Part 2	1 BOTTLE, PLASTIC	354 mL

### Part 1 of 2

### DAYTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, phenylephrine hcl liquid

### Product Information

Item Code (Source)	NDC:85828-772
Route of Administration	ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

### Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

## Product Characteristics

Color	orange	Score	
Shape		Size	
Flavor	APRICOT	Imprint Code	
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/02/2026	

## Part 2 of 2

### NIGHTTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate liquid

## Product Information

Item Code (Source)	NDC:85828-750
Route of Administration	ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	

<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>SORBITOL SOLUTION</b> (UNII: 8KW3E207O2)	

### Product Characteristics

<b>Color</b>	red	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	CHERRY	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/02/2026	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/02/2026	

**Labeler** - Grocery Outlet, Inc. (029161585)

**Registrant** - TIME CAP LABORATORIES, INC. (037052099)

### Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		677604129	manufacture(85828-773)