

DAIRYLAND BRAND PRE-POST- iodine liquid
Stearns Packaging Corporation

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Pre-Post 1/2%

IODOPHORS are considered relatively non-toxic, but should always be used in accordance with label directions. Consult your veterinarian in case of teat irritation.

TEAT DIPPING IN COLD WEATHER:

When the temperature falls below 10 degrees F, or when wind chill is significant, special precautions should be taken to avoid chapped and frozen teats.

1. Teats should be dry before turning cows out into cold weather.
2. When teats are dipped after milking, allow 30 seconds contact, and blot off any excess teat dip with a single service towel.
3. Warm the teat dip to reduce drying time.
4. Provide windbreaks in outside areas for cows.
5. Monitor fresh cows with swollen udders and teats since they are more susceptible to chapped and frozen teats.

PRODUCT INFORMATION: This teat dip is an IODOPHOR (a combination of iodine and complexing agents), and when properly used, is effective as an aid in reducing the spread of mastitis causing organisms.

DO NOT DILUTE.

USE AT FULL STRENGTH.

PROTECT FROM FREEZING

Important Use Instructions

For Pre-Milking

1. Wash each individual teat with a sanitizing udder wash.
NOTE: For best results, keep hair around teat based clipped.
2. Hand strip teats to initial let down.
3. Pre-dip entire teat with PRE-POST 1/2% on all four quarters. Repeat process on next 4 to 6 cows or until first cow prepped has at least 1 minute contact time.
4. Wipe each individual teat with a clean paper towel. Utilize a "pulling down" motion which removes iodine solution, germs, and sediment from teat skin. Very important!
5. Place machine on cow and repeat wiping process on remaining cows that have been predipped.

For Post Milking

DIRECTIONS FOR TEAT DIPPING. Dip entire teat with undiluted dip solution. Allow to air dry and remain on teats for prolonged contact time. Clean, sanitize and thoroughly dry each teat prior to next milking. Replace dip solution when visibly cloudy or dirty.

FIRST-AID:

EYES: If contact with eyes occurs, flush with plenty of cool water for 15 minutes. Consult a physician.

INTERNAL: May be harmful if swallowed. If ingested, drink large amounts of water. DO NOT induce vomiting. Get medical attention immediately.

24 HOUR EMERGENCY CONTACT
1-800-255-3924

DAIRYLAND BRAND

PRE-POST 1/2%

A SANITIZING PRE AND POST DIP

NPE-Free Aid in Reducing the Spread of Organisms Which May Cause Mastitis

Active Ingredient.....0.5% Available Iodine

Emollient System Contains.....5% Glycerin Minimum pH.....4.0

KEEP OUT OF THE REACH OF CHILDREN

See side panel for Precautionary Statements STEARNS PACKAGING CORPORATION

4200 Sycamore Ave. (53714), PO Box 3216, Madison, WI 53704

(800) 655-5008 www.stearnspkg.com

NET CONTENTS: 1 U.S. FL. GALLON (3.78 LITERS)

DO NOT DILUTE. USE AT FULL STRENGTH. PROTECT FROM FREEZING. Important Use Instructions

FOR PRE-MILKING

1. Wash each individual teat with a sanitizing udder wash.
- NOTE: For best results, keep teat around teat base clipped.
2. Hand strip teats to initiate let down.
3. Pre-dip entire teat with PRE-POST 1/2% on all four quarters. Repeat process on next 4 teats or until first cow prepreg has at least 1 minute contact time.
4. Wipe each individual teat with a clean paper towel. Utilize a "pulling down" motion which removes iodine solution, grease, and sediment from teat sheath. Very important!
5. Place machine on cow and repeat milking process on remaining cows that have been pre-dipped.

FOR POST MILKING

DIRECTIONS FOR TEAT DIPPING: Dip entire teat with undiluted dip solution. Allow to air dry and remain on teats for prolonged contact time. Clean, sanitize and thoroughly dry each teat prior to next milking. Replace dip solution when visibly cloudy or dirty.

PRODUCT INFORMATION: This teat dip is a 100%NPE (a combination of iodine and complexing agents) and when properly used, is effective as an aid in reducing the spread of mastitis causing organisms. 100%NPEs are considered relatively non-toxic, but should always be used in accordance with label directions. Consult your veterinarian in case of teat irritation.

TEAT DIPPING IN COLD WEATHER: When the temperature falls below 10°F, or when wind chill is significant, special precautions should be taken to avoid chapped and frozen teats.

1. Teats should be dry before being covered in cold weather.
2. When teats are dipped after milking, allow 30 seconds contact, and blot off any excess teat dip with a single service towel.
3. Warm the teat dip to reduce drying time.
4. Provide windbreaks in outside areas for cows.
5. Monitor fresh cows with swollen udders and teats since they are more susceptible to chapped and frozen teats.

DAIRYLAND BRAND

PRE-POST 1/2%
A SANITIZING PRE & POST DIP

NPE-Free Aid in Reducing the Spread of Organisms Which May Cause Mastitis

Active Ingredient0.5% Available Iodine
Emollient System Contains5% Glycerin
Minimum pH4.0

KEEP OUT OF THE REACH OF CHILDREN
See side panel for Precautionary Statements.

STEARNS PACKAGING CORPORATION
4200 Sycamore Ave. (53714), PO Box 3216, Madison, WI 53704
(800) 655-5008 • www.stearnspkg.com

NET CONTENTS: 1 U.S. GALLON (3.78 L)

FIRST-AID:
EYES: If contact with eyes occurs, flush with plenty of cool water for 15 minutes. Consult a physician.
INTERNAL: May be harmful if swallowed. If ingested, drink large amounts of water. DO NOT induce vomiting. Get medical attention immediately.

PRIMEROS AUXILIOS:
OJOS: Si ocurre contacto con los ojos, lavar con abundante agua fría durante 15 minutos. Consulte a un médico.
INTERNO: Puede ser nocivo en caso de ingestión. Beba grandes cantidades de agua si se ingiere este producto. NO induzca el vómito. Obtenga atención médica inmediatamente.

24 HOUR EMERGENCY CONTACT
24 horas servicio telefónico de repuesta a emergencias:
1-800-255-3924

#1244010
REORDER NUMBER

LOT NO:

EXP. DATE:

USE ESTE PRODUCTO CON TODA SU FUERZA. NO DEJE CONGELAR. Instrucciones de uso importantes.

PRE-SELLADO

1. Lave cada pezón individual con un lavado de la ubre de desinfección.
- NOTA: Para obtener los mejores resultados mantenga el pelo alrededor de la base de cada pezón bien recortado.
2. Utilice una solución de empuje con la mano para iniciar la liberación de la leche.
3. Repita el tratamiento del pre-velado por todo el pezón con Pre-Post 1/2% en los cuatro cuartos de la ubre. Repita el proceso en las siguientes 4 ubres hasta que la primera vaca ha tenido por lo menos 1 minuto de tiempo de contacto.
4. Limpie cada pezón individual con una toalla de papel limpia. Utilice una "tirando abajo" de movimiento, lo que eliminará las soluciones de yodo, los pigmentos y los sedimentos de la piel del pezón. Esto es muy importante!
5. Coloque la máquina de ordeño en la ubre y repita el proceso de limpieza de las vacas restantes que requieren el tratamiento pre-celada.

EL SELADO POST-ORDEÑO

NOTIFICACIONES POR EL SELADO DE POST-ORDEÑO: Semejante toda la hembra con la inmersión del pezón sin diluir. Deje que el sellador se seque al aire y permanecer en el pezón por una cantidad prolongada de tiempo. Limpie, desinfecte y seca el fondo cada pezón antes de la siguiente ordeño. Vuelva a colocar el sellador de inmersión del pezón cuando se vuelva volublemente tibia a vaca.

INFORMACION DEL PRODUCTO: Este producto sellador es un yodoforo (una combinación de yodo y formadores de complejo) y cuando se utiliza correctamente es eficaz como una ayuda en la reducción de la propagación de organismos causantes de mastitis. Todavía se consideran relativamente no tóxicos, pero siempre se deben utilizar de acuerdo con instrucciones de la etiqueta. Consulte a su veterinario en caso de irritación de los pezones.

BANO DE PEZONES EN TIEMPO FRÍO: Cuando la temperatura cae por debajo de 10°F (-12°C), o cuando el factor de enfriamiento del viento es importante tome precauciones especiales para evitar los pezones agrietados y congelados.

1. Los pezones deben estar secos antes de aplicar las vacas en el frío.
2. Cuando los pezones se sumergen después del ordeño, espere 30 segundos de contacto, y posteriormente sacarlos el exceso de baño de pezones con una toalla desechable.
3. Caliente el baño de pezones para reducir el tiempo de secado.
4. Proporcione protección contra el viento en las áreas fuera para las vacas.
5. Vigile las vacas frescas con ubres hinchadas y pezones ya que son más susceptibles a los pezones agrietados y congelados.

#627610 Rev. 2016

DAIRYLAND BRAND PRE-POST

iodine liquid

Product Information				
Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:60282-2106	
Route of Administration	TOPICAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
IODINE (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4)		IODINE	5 g in 1 L	
Inactive Ingredients				
Ingredient Name		Strength		
GLYCERIN (UNII: PDC6A3C0OX)		50 g in 1 L		
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:60282-2106-1	3.78 L in 1 BOTTLE, PLASTIC		
2	NDC:60282-2106-2	18.93 L in 1 PAIL		
3	NDC:60282-2106-3	56.78 L in 1 DRUM		

4	NDC:60282-2106-4	208.2 L in 1 DRUM		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
unapproved drug other		06/21/2004		

Labeler - Stearns Packaging Corporation (006069256)

Registrant - Stearns Packaging Corporation (006069256)

Establishment			
Name	Address	ID/FEI	Business Operations
Stearns Packaging Corporation		006069256	manufacture, api manufacture

Revised: 12/2017

Stearns Packaging Corporation