

DAIRYLAND BRAND PRE-POST- iodine liquid
Stearns Packaging Corporation

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Pre-Post 1/2%

IODOPHORS are considered relatively non-toxic, but should always be used in accordance with label directions. Consult your veterinarian in case of teat irritation.

TEAT DIPPING IN COLD WEATHER:

When the temperature falls below 10 degrees F, or when wind chill is significant, special precautions should be taken to avoid chapped and frozen teats.

1. Teats should be dry before turning cows out into cold weather.
2. When teats are dipped after milking, allow 30 seconds contact, and blot off any excess
teat dip with a single service towel.
3. Warm the teat dip to reduce drying time.
4. Provide windbreaks in outside areas for cows.
5. Monitor fresh cows with swollen udders and teats since they are more susceptible to chapped and frozen teats.

PRODUCT INFORMATION: This teat dip is an IODOPHOR (a combination of iodine and complexing agents), and when properly used, is effective as an aid in reducing the spread of mastitis causing organisms.

DO NOT DILUTE.
USE AT FULL STRENGTH.
PROTECT FROM FREEZING
Important Use Instructions

For Pre-Milking

1. Wash each individual teat with a sanitizing udder wash.
NOTE: For best results, keep hair around teat based clipped.
2. Hand strip teats to initial let down.
3. Pre-dip entire teat with PRE-POST 1/2% on all four quarters. Repeat process on next 4 to 6 cows or until first cow prepped has at least 1 minute contact time.
4. Wipe each individual teat with a clean paper towel. Utilize a "pulling down" motion which removes iodine solution, germs, and sediment from teat skin. Very important!
5. Place machine on cow and repeat wiping process on remaining cows that have been predipped.

For Post Milking

DIRECTIONS FOR TEAT DIPPING. Dip entire teat with undiluted dip solution. Allow to air dry and remain on teats for prolonged contact time. Clean, sanitize and thoroughly dry each teat prior to next milking. Replace dip solution when visibly cloudy or dirty.

FIRST-AID:

EYES: If contact with eyes occurs, flush with plenty of cool water for 15 minutes. Consult a physician.

INTERNAL: May be harmful if swallowed. If ingested, drink large amounts of water. DO NOT induce vomiting. Get medical attention immediately.

24 HOUR EMERGENCY CONTACT

1-800-255-3924

DAIRYLAND BRAND

PRE-POST 1/2%

A SANITIZING PRE AND POST DIP

NPE-Free Aid in Reducing the Spread of Organisms Which May Cause Mastitis

Active Ingredient.....0.5% Available Iodine

Emollient System Contains.....5% Glycerin Minimum pH.....4.0

KEEP OUT OF THE REACH OF CHILDREN

See side panel for Precautionary Statements STEARNS PACKAGING CORPORATION

4200 Sycamore Ave. (53714), PO Box 3216, Madison, WI 53704

(800) 655-5008 www.stearnspkg.com

NET CONTENTS: 1 U.S. FL. GALLON (3.78 LITERS)

DO NOT DILUTE. USE AT FULL STRENGTH. PROTECT FROM FREEZING. Important Use Instructions

FOR PRE-MILKING

1. Wash each individual teat with a cambizing udder wash.
2. **NOTE:** For best results, keep hair around teat base clipped.
3. Repeat process on next 4 to 6 cows or until first cow prepped but at least 1 minute contact time.
4. Wipe each individual teat with a clean paper towel. Utilize a "pulling down" motion which removes iodine solution, germs, and sediment from teat skin. Very important!
5. Place machine on cow and repeat wiping process on remaining cows that have been prepped.

FOR POST MILKING

DIRECTIONS FOR TEAT DIPPING: Dip entire teat with undiluted dip solution. Allow to air dry and remain on teats for prolonged contact time. Clean, sanitize and thoroughly dry each teat prior to next milking. Replace dip solution when visibly cloudy or dirty.

PRODUCT INFORMATION: This test dip is an IODOPHOR (a combination of iodine and complexing agents), and when properly used, is effective as an aid in reducing the spread of mastitis causing organisms. IODOPHORS are considered relatively non-toxic, but should always be used in accordance with label directions. Consult your veterinarian in case of teat irritation.

TEAT DIPPING IN COLD WEATHER: When the temperature falls below 10°F, or when wind chill is significant, special precautions should be taken to avoid chapped and frozen teats.

1. Teats should be dry before dipping into cold weather.
2. When teats are dipped after milking, allow 30 seconds contact, and blot off any excess teat dip with a single service towel.
3. Warm the teat dip to reduce drying time.
4. Provide windbreaks in outside areas for cows.
5. Monitor fresh cows with swollen udders and teats since they are more susceptible to chapped and frozen teats.

DAIRYLAND BRAND

PRE-POST 1/2%
A SANITIZING PRE & POST DIP

NPE-Free Aid in Reducing the Spread of Organisms Which May Cause Mastitis

Active Ingredient0.5% Available Iodine
Emollient System Contains5% Glycerin
Minimum pH4.0

KEEP OUT OF THE REACH OF CHILDREN
See side panel for Precautionary Statements.

STEARNS PACKAGING CORPORATION
4200 Sycamore Ave. (53714), PO Box 3216, Madison, WI 53704
(800) 655-5008 • www.stearnspkg.com

NET CONTENTS: 1 U.S. GALLON (3.78 L)

FIRST-AID:
EYES: If contact with eyes occurs, flush with plenty of cool water for 15 minutes. Consult a physician.
INTERNAL: May be harmful if swallowed. If ingested, drink large amounts of water. DO NOT induce vomiting. Get medical attention immediately.

PRIMEROS AUXILIOS:
QUJOS: Si ocurre contacto con los ojos, lavar con abundante agua fría durante 15 minutos. Consulte a un médico.
INTERNO: Puede ser nocivo en caso de ingestión. Beba grandes cantidades de agua si se ingiere este producto. NO induzca el vómito. Obtenga atención médica inmediatamente.

24 HOUR EMERGENCY CONTACT
24 horas servicio telefónico de repuesta a emergencias:
1-800-255-3924

#1244010
REGORDER NUMBER

LOT NO:

EXP. DATE:

USE ESTE PRODUCTO CON TODA SU FUERZA. NO SE DILUYA. Instrucciones de uso importantes.

PRE-SELLADO

1. Lave cada pezon individual con un lavado de la ubre de desinfección.
2. Utilice una toalla de papel para limpiar el área alrededor de la base de cada pezon bien mojado.
3. NOTA: Para obtener los mejores resultados mantenga el pelo alrededor de la base de cada pezon bien mojado.
4. Repita el tratamiento de pre-sellado por todos los pezones con Pre-Post 1/2% en los cuatro cuartos de la ubre. Repita el proceso en las pezoneras 4-6 veces hasta que la primera vaca ha finalizado por lo menos 1 minuto de tiempo de contacto.
5. Limpie cada pezon individual con una toalla de papel limpia. Utilice una "tracción abajo" de movimiento, lo que eliminará la solución de yodo, los gérmenes y los sedimentos de la piel del pezon. Esto es muy importante!
6. Coloque la máquina de ordeño en la ubre y repita el proceso de limpieza de las vacas restantes que recibieron el tratamiento pre-sellado.

EL SELADO POST-ORDEÑO

INSTRUCCIONES POR EL SELADO DE POST-ORDEÑO: Sumerja toda la tetina con la inmersión del pezon en el líquido. Deje que el líquido se seque al aire y permanecer en el pezon por una cantidad prolongada de tiempo. Limpie, desinfecte y seca a fondo cada pezon antes de la siguiente ordeño. Vuelva a colocar el solución de inmersión del pezon cuando se vacie totalmente la teta o la vaca.

INFORMACIÓN DEL PRODUCTO: Este producto sellador es un yodoforo (una combinación de yodo y formadores de complejos) y cuando se utiliza correctamente es eficaz como una ayuda en la reducción de la propagación de organismos causantes de mastitis. Iodoforos se consideran relativamente no tóxicos, pero siempre se deben utilizar de acuerdo con instrucciones de la etiqueta. Consulte a su veterinario en caso de irritación de los pezones.

BANJO DE PEZONES EN TIEMPO FRÍO: Cuando la temperatura cae por debajo de 10°F (-12°C), o cuando el factor de enfriamiento del viento es importante tome precauciones especiales para evitar los pezones agrietados y congelados.

1. Los pezones deben estar secos antes de colocar las vacas en el frío.
2. Cuando los pezones se sumergen después del ordeño, espere 30 segundos de contacto, y posteriormente secarlos el exceso de baño de pezones con una toalla desechable.
3. Cubierta el baño de pezones para reducir el tiempo de secado.
4. Proporcione protección contra el viento en las áreas fuera para las vacas.
5. Vigile las vacas frescas con ubres hinchadas y pezones ya que son más susceptibles a los pezones agrietados y congelados.

TAKE TIME
OBSERVE LABEL DIRECTIONS

DAIRYLAND BRAND PRE-POST iodine liquid

Product Information			
Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:60282-2106
Route of Administration	TOPICAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
IODINE (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4)	IODINE	5 g in 1 L	

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	50 g in 1 L

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:60282-2106-1	3.78 L in 1 BOTTLE, PLASTIC		
2	NDC:60282-2106-2	18.93 L in 1 PAIL		
3	NDC:60282-2106-3	56.78 L in 1 DRUM		
4	NDC:60282-2106-4	208.2 L in 1 DRUM		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		06/21/2004	

Labeler - Stearns Packaging Corporation (006069256)

Registrant - Stearns Packaging Corporation (006069256)

Establishment

Name	Address	ID/FEI	Business Operations
Stearns Packaging Corporation		006069256	manufacture

Establishment

Name	Address	ID/FEI	Business Operations
Lab Express International		800875106	api manufacture

Revised: 11/2025

Stearns Packaging Corporation