

**OMEPRAZOLE- omeprazole tablet, delayed release**  
**P & L Development, LLC**

-----  
**OMEPRAZOLE MAGNESIUM DELAYED-RELEASE TABLETS**

***Active ingredient (in each tablet)***

Omeprazole delayed-release tablet 20 mg  
(equivalent to 20.6 mg omeprazole magnesium)

***Purpose***

Acid reducer

***Use***

- treats frequent heartburn (occurs ***2 or more*** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

***Warnings***

**Allergy alert:**

- Do not use if you are allergic to omeprazole
- Omeprazole may cause severe skin reactions. Symptoms may include:
  - skin reddening
  - blisters
  - rash

if an allergic reaction occurs, stop use and seek medical help right away.

**Do not use if you have:**

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**

These may be signs of a serious condition. See your doctor.

**Ask a doctor before use if you have:**

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting

- stomach pain

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug.

Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if:**

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

***Directions***

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

**14-Day Course of Treatment**

- swallow 1 tablet with a glass of water before eating in the morning
- take everyday for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- **swallow whole. Do not chew or crush tablets.**

**Repeated 14-Day Courses (if needed)**

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
- children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

***Other information***

- read the directions and warnings before use
- Keep the carton. It contains important information.
- store at 20° to 25°C (68° to 77°F) and protect from moisture

### ***Inactive ingredients***

acetyl tributyl citrate, colloidal silicon dioxide, corn starch, croscarmellose sodium, hydroxypropyl cellulose, hypromellose 2910, magnesium stearate, methacrylic acid copolymer type C, microcrystalline cellulose, polyethylene glycol 400, polyethylene glycol 3350, polyethylene glycol 8000, polyvinyl alcohol, red iron oxide, sucrose, talc, titanium dioxide, triethyl citrate

### ***Tips for Managing Heartburn***

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

### **Principal display panel**

Compare to the active ingredient in Prilosec® OTC†

24-hour

### **Omeprazole**

Delayed-release tablets 20 mg

Acid reducer

Treats FREQUENT heartburn !

May take 1 to 4 days for full effect

Tablets

One 14-day course of treatment

†This product is not affiliated with, manufactured by or produced by the makers or owners of Prilosec OTC®.

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

Distributed by:

**PL Developments**  
 200 Hicks Street  
 Westbury, NY 11590

**Package label**

Distributed by:  
**PL Developments**  
 200 Hicks Street  
 Westbury, NY 11590

EXCELLENCE  
QUALITY

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

This product is not affiliated with, manufactured by or produced by the makers or owners of Prilosec OTC®.

<p><b>Drug Facts (continued)</b></p> <p>Acid reducers may interact with certain prescription drugs.</p> <p><b>Stop use and ask a doctor if:</b></p> <ul style="list-style-type: none"> <li>■ your heartburn continues or worsens</li> <li>■ you need to take this product for more than 14 days</li> <li>■ you need to take more than 1 course of treatment every 4 months</li> <li>■ you get diarrhea</li> <li>■ you develop a rash or joint pain</li> </ul> <p><b>If pregnant or breast-feeding,</b> ask a health professional before use.</p> <p><b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.</p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>■ for adults 18 years of age and older</li> <li>■ this product is to be used once a day (every 24 hours), every day for 14 days</li> <li>■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours</li> </ul> <p><b>14-Day Course of Treatment</b></p> <ul style="list-style-type: none"> <li>■ swallow 1 tablet with a glass of water before eating in the morning</li> <li>■ take every day for 14 days</li> <li>■ do not take more than 1 tablet a day</li> <li>■ do not use for more than 14 days unless directed by your doctor</li> <li>■ swallow whole. Do not crush or chew tablets.</li> </ul> <p><b>Repeated 14-Day Courses (if needed)</b></p> <ul style="list-style-type: none"> <li>■ you may repeat a 14-day course every 4 months</li> <li>■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor</li> </ul>	<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.</li> </ul> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>■ read the directions and warnings before use</li> <li>■ keep the carton. It contains important information.</li> <li>■ store at 20° to 25°C (68° to 77°F) and protect from moisture</li> </ul> <p><b>Inactive ingredients</b> acetyl tributyl citrate, colloidal silicon dioxide, corn starch, croscarmellose sodium, hydroxypropyl cellulose, hypromellose 2910, magnesium stearate, methacrylic acid copolymer type C, microcrystalline cellulose, polyethylene glycol 400, polyethylene glycol 3350, polyethylene glycol 8000, polyvinyl alcohol, red iron oxide, sucrose, talc, titanium dioxide, triethyl citrate</p> <p><b>Questions or comments?</b>        Call 1-877-753-3935 Monday-Friday 9AM-5PM EST</p> <p><b>Tips for Managing Heartburn</b></p> <ul style="list-style-type: none"> <li>■ Do not lie flat or bend over after eating</li> <li>■ Do not wear tight-fitting clothing around the stomach</li> <li>■ Do not eat before bedtime</li> <li>■ Raise the head of your bed</li> <li>■ Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables</li> <li>■ Eat slowly and avoid big meals</li> <li>■ If overweight, lose weight</li> <li>■ Quit smoking</li> </ul>	<div style="text-align: center;">  <p style="font-size: x-small;">Compare to the active ingredient in Prilosec OTC®† NDC 59726-298-14</p> </div> <div style="text-align: center; border: 1px solid black; border-radius: 50%; padding: 20px; margin: 10px auto; width: 80%;"> <p style="font-size: x-small;">see current drug facts</p> <p style="font-size: x-small;">24-hour</p> <h1 style="margin: 0;">Omeprazole</h1> <p style="font-size: x-small;">delayed-release tablets, 20 mg acid reducer</p> <p style="font-size: x-small;">treats FREQUENT heartburn! may take 1 to 4 days for full effect</p> </div> <div style="text-align: center; margin-top: 10px;"> <p style="font-size: x-small;">14 tablets one 14-day course of treatment</p>  <p style="font-size: x-small;">actual size</p> </div>	<p><b>Drug Facts</b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="font-size: x-small;">Active ingredient (in each tablet)</th> <th style="font-size: x-small;">Purpose</th> </tr> </thead> <tbody> <tr> <td style="font-size: x-small;">Omeprazole delayed-release tablet 20 mg (equivalent to 20.6 mg omeprazole magnesium).....</td> <td style="font-size: x-small;">Acid reducer</td> </tr> </tbody> </table> <p><b>Use</b> ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p> <p><b>Warnings</b></p> <p><b>Allergy alert:</b> ■ Do not use if you are allergic to omeprazole. ■ Omeprazole may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If an allergic reaction occurs, stop use and seek medical help right away.</p> <p><b>Do not use if you have:</b> ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain. These may be signs of a serious condition. See your doctor.</p> <p><b>Ask a doctor before use if you have:</b> ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain</p> <p><b>Ask a doctor or pharmacist before use if you are taking a prescription drug.</b></p>	Active ingredient (in each tablet)	Purpose	Omeprazole delayed-release tablet 20 mg (equivalent to 20.6 mg omeprazole magnesium).....	Acid reducer
Active ingredient (in each tablet)	Purpose						
Omeprazole delayed-release tablet 20 mg (equivalent to 20.6 mg omeprazole magnesium).....	Acid reducer						



3 59726 05114 9

PLD-B849A  
FC009469

Lot No:  
Exp. Date:

**Omeprazole delayed release tablets 20 mg**

**OMEPRAZOLE**  
 omeprazole tablet, delayed release

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:59726-298
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>OMEPRAZOLE MAGNESIUM</b> (UNII: 426QFE7XLK) (OMEPRAZOLE - UNII:KG60484QX9)	OMEPRAZOLE	20.6 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>ACETYLTRIBUTYL CITRATE</b> (UNII: 0ZBX0N59RZ)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>HYDROXYPROPYL CELLULOSE, UNSPECIFIED</b> (UNII: 9XZ8H6N6OH)	
<b>HYPROMELLOSE 2910 (15 MPA.S)</b> (UNII: 36SFW2JZ0W)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER</b> (UNII: NX76LV5T8J)	
<b>POLYETHYLENE GLYCOL 400</b> (UNII: B697894SGQ)	
<b>POLYETHYLENE GLYCOL 3350</b> (UNII: G2M7P15E5P)	
<b>POLYETHYLENE GLYCOL 8000</b> (UNII: Q662QK8M3B)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>TRIETHYL CITRATE</b> (UNII: 8Z96QXD6UM)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	

## Product Characteristics

<b>Color</b>	red (reddish brown)	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	13mm
<b>Flavor</b>		<b>Imprint Code</b>	O
<b>Contains</b>			

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:59726-298-14	1 in 1 BOX	01/31/2025	
1		14 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206582	01/31/2025	

**Labeler** - P & L Development, LLC (800014821)

Revised: 6/2025

P & L Development, LLC