

ANTIMONIUM METALLICUM- antimonium metallicum pellet
Hahnemann Laboratories, Inc.

Disclaimer: This homeopathic product has not been evaluated by the Food and Drug Administration for safety or efficacy. FDA is not aware of scientific evidence to support homeopathy as effective.

Single label image contains content for sections 1-17.

ANTIMONIUM METALLICUM

30C

EST 63545

CONSTIPATION

Contents: 5g

HOMEOPATHIC MEDICINE.

Dose: Adult and child take 2-3 pellets once

or as directed by your prescriber. Warning:

If pregnant, nursing, or if symptoms persist

consult your practitioner.

USE according to standard Homeopathic indications for self-limiting conditions, such as listed above.


DIRECTIONS: Dissolve one dose under tongue. Repeat as needed or as directed. Stop taking if symptoms worsen.

Keep out of reach of children.

ACTIVE INGREDIENT listed above.

QUESTIONS: Toll free 888-427-5422

INACTIVE INGREDIENT: sucrose.



0000

Hahnemann Laboratories, Inc. 1949 Fourth St. San Rafael, CA 94901
TWIST TO OPEN. NO OTHER SEALING

antimonium metallicum pellet						
Product Information						
Product Type		HUMAN OTC DRUG	Item Code (Source)	NDC:63545-651		
Route of Administration		BUCCAL, SUBLINGUAL, ORAL				
Active Ingredient/Active Moiety						
Ingredient Name			Basis of Strength	Strength		
ANTIMONY (UNII: 9IT35J3UV3) (ANTIMONY - UNII:9IT35J3UV3)			ANTIMONY	30 [hp_C] in 1 [hp_C]		
Inactive Ingredients						
Ingredient Name			Strength			
SUCROSE (UNII: C151H8M554)						
Product Characteristics						
Color	white	Score	no score			
Shape	ROUND	Size	3mm			
Flavor		Imprint Code				
Contains						
Packaging						
#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:63545-651-01	240 [hp_C] in 1 VIAL, GLASS; Type 0: Not a Combination Product	12/01/2016			

ANTIMONIUM METALLICUM

30C EST 63545

CONSTIPATION Contents: 5g

HOMEOPATHIC MEDICINE.

Dose: Adult and child take 2-3 pellets once

or as directed by your prescriber. Warning:

If pregnant, nursing, or if symptoms persist

consult your practitioner.

USE according to standard Homeopathic indications for

self-limiting conditions, such as listed above.

DIRECTIONS: Dissolve one dose under tongue. Repeat as

needed or as directed. Stop taking if symptoms worsen.

Keep out of reach of children.

ACTIVE INGREDIENT listed above.

QUESTIONS: Toll free 888-427-6422

INACTIVE INGREDIENT: sucrose.

0000

Hahnemann Laboratories, Inc. 1949 Fourth St. San Rafael, CA 94901
TWIST TOP OPEN AND USE AS DIRECTED

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

unapproved homeopathic	12/01/2016
------------------------	------------

Labeler - Hahnemann Laboratories, Inc. (147098081)

Establishment

Name	Address	ID/FEI	Business Operations
Hahnemann Laboratories, Inc.		147098081	manufacture(63545-651)

Revised: 12/2016

Hahnemann Laboratories, Inc.