

**TOP VALUE EFFERVESCENT PAIN RELIEF- aspirin, citric acid, sodium bicarbonate tablet, effervescent**  
**Tower Laboratories Ltd**

-----  
**Effervescent Pain Relief Tablets 36 ct.**

**Active Ingredients**

(in each effervescent tablet)

Aspirin 325 mg (NSAID)\*

Citric acid 1000 mg

Sodium bicarbonate (heat-treated) 1916 mg

\*Nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever

Antacid

Antacid

**Uses**

for the temporary relief of ■ heartburn, acid indigestion and sour stomach when accompanied by headache or body aches and pains ■ upset stomach with headache from overindulgence in food or drink

■ pain alone (headaches or body aches and pains)

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

## **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- if you are allergic to aspirin ■ if overly full from food or drink. Ask a doctor if severe stomach pain occurs after taking this product.

## **Ask a doctor before use if**

- stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have a sodium-restricted diet

## **Ask a doctor or pharmacist before use if you are**

- presently taking a prescription drug. Antacids may interact with certain prescription drugs.
- taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis

## **When using this product**

do not take more than 8 tablets (adults) or 4 tablets (60 years and over) in a 24-hour period or use the maximum dosage for more than 10 days

## **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away.
- pain lasts for more than 10 days or gets worse
- new symptoms occur
- redness or swelling is present
- ringing in the ears or a loss of hearing occurs

## **If pregnant or breast-feeding,**

ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by**

**a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

### **Directions**

■ do not exceed recommended dosage

■ to avoid serious injury, completely dissolve tablets in 4 oz. of water before taking adults and children 12 years and over (up to 60 years of age) 2 tablets every 4 hours

do not exceed 8 tablets in 24 hours

60 years and over 2 tablets every 4 hours do not exceed 4 tablets in 24 hours

### **Other information**

■ **each tablet contains:** sodium 524 mg

■ store at room temperature (59°-86°F)

### **Inactive Ingredients**

■ none

**Serious side effects associated with the use of this product can be reported to the following address:**

Tower Laboratories LTD

P.O. Box 306

8 Industrial Park Road

Centerbrook, CT 06409

**PACKAGE LABEL. PRINCIPAL DISPLAY PANEL**



TOP VALUE EFFERVESCENT PAIN RELIEF			
aspirin, citric acid, sodium bicarbonate tablet, effervescent			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:50201-4213
<b>Route of Administration</b>	ORAL		
<b>Active Ingredient/Active Moiety</b>			
Ingredient Name		Basis of Strength	Strength
<b>SODIUM BICARBONATE</b> (UNII: 8MDF5V39QO) (BICARBONATE ION - UNII:HN1ZRA3Q20)		SODIUM BICARBONATE	1916 mg
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)		ASPIRIN	325 mg
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL) (ANHYDROUS CITRIC ACID - UNII:XF417D3PSL)		ANHYDROUS CITRIC ACID	1000 mg

## Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	25mm
<b>Flavor</b>		<b>Imprint Code</b>	L421
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50201-4213-6	18 in 1 CARTON	12/01/2006	
1		2 in 1 POUCH; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	12/01/2006	

**Labeler** - Tower Laboratories Ltd (001587203)

**Registrant** - Tower Laboratories Ltd (869024500)

Revised: 12/2024

Tower Laboratories Ltd