

**MEDI-FIRST BURN- lidocaine hydrochloride gel**  
**Unifirst First Aid Corporation**

-----  
**Medi-First Burn Gel Drug Facts**

***Drug Facts***

***Active ingredient (in each gram)***

Lidocaine HCl 2.0%

***Purpose***

Topical analgesic

***Uses***

For the temporary relief of pain associated with minor burns.

***Warnings***

**For external use only.**

**Do not use**

- in large quantities, particularly over raw or blistered areas

**When using this product**

- avoid contact with the eyes

**Stop use ask a doctor if**

- condition worsens
- symptoms persist for more than 7 days or clears up and occurs again within a few days

**Keep out of the reach of children.**

If swallowed, get medical help or contact a Poison Control Center right away.

***Directions***

- **adults & children 2 years and older:** apply to affected area not more than 3 to 4 times daily
- **children under 2:** consult a doctor

***Inactive ingredients***

aloe vera, carbomer, germaben II, propylene glycol, purified water, menthol,

triethanolamine, vitamin E acetate

## Questions?

Call 1-800-634-7680

Mfg. for Medique Products, 17080 Alico Commerce Ct., Ste 1, Fort Myers, FL 33967

## Medi-First Burn Gel Label

Medi-First ®

Burn Gel

Lidocaine HCl / Topical Analgesic

Product # 47769

NET CONTENTS: 6 Units of 1/8 oz

|  |   |   |                                     |
|--|---|---|-------------------------------------|
| <b>Drug Facts</b>  | <b>Burn Gel</b><br>Product # 47769<br>6 per Box | <b>MEDI-FIRST®</b><br><b>Burn Gel</b><br>Lidocaine HCl / Topical Analgesic<br>Product # 47769<br>NET CONTENTS: 6 Units of 1/8oz |                                     |
| <b>Active ingredient (in each gram)</b><br>Lidocaine HCl 2.0%.....   |   |   | <b>Purpose</b><br>Topical analgesic |
| <b>Uses</b> For the temporary relief of pain associated with minor burns.  |   |   |                                     |
| <b>Warnings</b><br>For external use only.  |   |   |                                     |
| Do not use ■ in large quantities, particularly over raw surfaces or blistered areas  |   |   |                                     |
| When using this product ■ avoid contact with the eyes  |   |   |                                     |
| Stop use ask a doctor if ■ condition worsens ■ symptoms persist for more than 7 days or clear up and occur again within a few days                       |   |   |                                     |
| <b>Directions</b> ■ adults & children 2 years and older: apply to affected area not more than 3 to 4 times daily<br>■ children under 2: consult a doctor |   |   |                                     |
| <b>Inactive ingredients</b> aloe vera, carbomer, germben II, propylene glycol, purified water, menthol, triethanolamine, vitamin E acetate               |   |   |                                     |
| <b>Questions?</b> Call 1-800-634-7680  |   | Retain carton for complete product information.   |                                     |

## MEDI-FIRST BURN

lidocaine hydrochloride gel

### Product Information

|                                |                |                           |               |
|--------------------------------|----------------|---------------------------|---------------|
| <b>Product Type</b>            | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:47682-477 |
| <b>Route of Administration</b> | TOPICAL        |                           |               |

### Active Ingredient/Active Moiety

| Ingredient Name   | Basis of Strength                 | Strength     |
|---|-----------------------------------|--------------|
| <b>LIDOCAINE HYDROCHLORIDE</b> (UNII: V13007Z41A) (LIDOCAINE - UNII:98PI200987) | LIDOCAINE HYDROCHLORIDE ANHYDROUS | 20 mg in 1 g |

### Inactive Ingredients

| Ingredient Name                             | Strength |
|---|----------|
| <b>DIAZOLIDINYL UREA</b> (UNII: H5RIZ3MPW4) |          |
| <b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)  |          |
| <b>WATER</b> (UNII: 059QF0KO0R)             |          |
| <b>ALOE VERA LEAF</b> (UNII: ZY81Z83H0X)    |          |

|  |  |
|--|--|
| <b>MENTHOL</b> (UNII: L7T10EIP3A)  |  |
| <b>TROLAMINE</b> (UNII: 9O3K93S3TK)  |  |
| <b>.ALPHA.-TOCOPHEROL ACETATE</b> (UNII: 9E8X80D2L0)                               |  |
| <b>CARBOMER INTERPOLYMER TYPE A (ALLYL SUCROSE CROSSLINKED)</b> (UNII: 59TL3WG5CO) |  |
| <b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)  |  |
| <b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)  |  |

### Packaging

| # | Item Code        | Package Description                                  | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:47682-477-69 | 6 in 1 BOX, UNIT-DOSE                                | 07/30/2014           |                    |
| 1 |                  | 3.3 g in 1 PACKET; Type 0: Not a Combination Product |                      |                    |

### Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M015                                     | 07/30/2014           |                    |

**Labeler** - Unifirst First Aid Corporation (832947092)

Revised: 9/2025

Unifirst First Aid Corporation