

MAGNESIA ACETICA- magnesia acetica pellet
Hahnemann Laboratories, Inc.

Disclaimer: This homeopathic product has not been evaluated by the Food and Drug Administration for safety or efficacy. FDA is not aware of scientific evidence to support homeopathy as effective.

Single label image contains content for sections 1-17.

MAGNESIA ACETICA

30C

EST 63545

DIARRHEA

Contents: 5g

HOMEOPATHIC MEDICINE.

Dose: Adult and child take 2-3 pellets once or as directed by your prescriber. **Warning: If pregnant, nursing, or if symptoms persist consult your practitioner.**

USE according to standard Homeopathic indications for self-limiting conditions, such as listed above.
DIRECTIONS: Dissolve one dose under tongue. Repeat as needed or as directed. Stop taking if symptoms worsen.

Keep out of reach of children.

ACTIVE INGREDIENT listed above.

QUESTIONS: Toll free 888-427-6422

INACTIVE INGREDIENT: sucrose.

|||||

0000

Hahnemann Laboratories, Inc. 1941 Fourth St. San Rafael, CA 94901
TAPE TO OPEN (SEE ONLY IF SEALED)

magnesia acetica pellet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63545-662
Route of Administration	BUCCAL, ORAL, SUBLINGUAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MAGNESIUM ACETATE TETRAHYDRATE (UNII: I01G0EJC3B) (MAGNESIUM CATION - UNII:T6V3LHY838)	MAGNESIUM ACETATE TETRAHYDRATE	30 [hp_C] in 1 [hp_C]

Inactive Ingredients

Ingredient Name	Strength
SUCROSE (UNII: C151H8M554)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	3mm
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63545-662-01	240 [hp_C] in 1 VIAL, GLASS; Type 0: Not a Combination Product	12/01/2016	

MAGNESIA ACETICA

30C

EST. 63545

DIARRHEA

Contents: 5g

HOMEOPATHIC MEDICINE.

Dose: Adult and child take 2-3 pellets once or as directed by your prescriber. Warning: If pregnant, nursing, or if symptoms persist consult your practitioner.

USE according to standard Homeopathic indications for self-limiting conditions, such as listed above.
DIRECTIONS: Dissolve one dose under tongue. Repeat as needed or as directed. Stop taking if symptoms worsen.
Keep out of reach of children.
ACTIVE INGREDIENT listed above.
QUESTIONS: Toll free 888-427-6422
INACTIVE INGREDIENT: sucrose.

|||||
0000

Hahnemann Laboratories, Inc. 190 Fourth St. San Rafael, CA 94901
TWIST TO OPEN. ONLY IF SEALED

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

unapproved homeopathic

12/01/2016

Labeler - Hahnemann Laboratories, Inc. (147098081)

Establishment

Name	Address	ID/FEI	Business Operations
Hahnemann Laboratories, Inc.		147098081	manufacture(63545-662)

Revised: 12/2016

Hahnemann Laboratories, Inc.