

VETERINARY LACTATED RINGERS- sodium chloride, sodium lactate, potassium chloride, calcium chloride injection, solution
Baxter Healthcare Corporation

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Veterinary Lactated Ringer's Injection, USP
For Animal Use Only

Lactated Ringer's Injection, USP is a sterile, nonpyrogenic solution for fluid and electrolyte replenishment in single dose containers for intravenous administration. It contains no antimicrobial agents. Discard unused portion. Composition, osmolarity, pH, ionic concentration and caloric content are shown in Table 1.

Table 1

| | Size (mL) | Composition (g/L) | | | | Osmolarity (mOsmol/L) (calc) | pH | Ionic Concentration (mEq/L) | | | | | Caloric Content (kcal/L) |
|---|-----------|------------------------------|---|--------------------------------|--|------------------------------|---------------------|-----------------------------|-----------|---------|----------|---------|--------------------------|
| | | Sodium Chloride, USP, (NaCl) | Sodium Lactate, (C ₃ H ₅ NaO ₃) | Potassium Chloride, USP, (KCl) | Calcium Chloride, USP (CaCl ₂ ·2H ₂ O) | | | Sodium | Potassium | Calcium | Chloride | Lactate | |
| Veterinary Lactated Ringer's Injection, USP | 5000 | 6 | 3.1 | 0.3 | 0.2 | 273 | 6.5 (6.0 to 7.5) | 130 | 4 | 2.7 | 109 | 28 | 9 |

The plastic container is fabricated from a specially formulated polyvinyl chloride. The amount of water that can permeate from inside the container into the overwrap is insufficient to affect the solution significantly. Solutions in contact with the plastic container can leach out certain of its chemical components in very small amounts within the expiration period, e.g., di-2-ethylhexyl phthalate (DEHP), up to 5 parts per million. However, the safety of the plastic has been confirmed in tests in animals according to USP biological tests for plastic containers as well as by tissue culture toxicity studies.

Lactated Ringer's Injection, USP has value as a source of water and electrolytes. It is capable of inducing diuresis depending on the clinical condition of the

patient.

Lactated Ringer's Injection, USP produces a metabolic alkalinizing effect. Lactate ions are metabolized ultimately to carbon dioxide and water, which requires the consumption of hydrogen cations.

Lactated Ringer's Injection, USP is indicated as a source of water and electrolytes or as an alkalinizing agent.

None known

Lactated Ringer's Injection, USP should be used with great care, if at all, in patients with congestive heart failure, severe renal insufficiency, and in clinical states in which there exists edema with sodium retention.

Lactated Ringer's Injection, USP should be used with great care, if at all, in patients with hyperkalemia, severe renal failure, and in conditions in which potassium retention is present.

Lactated Ringer's Injection, USP should be used with great care in patients with metabolic or respiratory alkalosis. The administration of lactate ions should be done with great care in those conditions in which there is an increased level or an impaired utilization of these ions, such as severe hepatic insufficiency.

Lactated Ringer's Injection, USP should not be administered simultaneously with blood through the same administration set because of the likelihood of coagulation.

The intravenous administration of Lactated Ringer's Injection, USP can cause fluid and/or solute overloading resulting in dilution of serum electrolyte concentrations, overhydration, congested states, or pulmonary edema. The risk of dilutional states is inversely proportional to the electrolyte concentrations of the injections. The risk of solute overload causing congested states with peripheral and pulmonary edema is directly proportional to the electrolyte concentrations of the injections.

In patients with diminished renal function, administration of Lactated Ringer's Injection, USP may result in sodium or potassium retention.

Lactated Ringer's Injection, USP is not for use in the treatment of lactic acidosis.

Reactions which may occur because of the solution or the technique of administration include febrile response, infection at the site of injection, venous thrombosis or phlebitis extending from the site of injection, extravasation, and hypervolemia. If an adverse reaction does occur, discontinue the infusion, evaluate the patient, institute appropriate therapeutic countermeasures, and save the remainder of the fluid for examination if deemed necessary.

Clinical evaluation and periodic laboratory determinations are necessary to monitor changes in fluid balance, electrolyte concentrations, and acid base balance during prolonged parenteral therapy or whenever the condition of the patient warrants such evaluation.

Lactated Ringer's Injection, USP must be used with caution. Excess administration may result in metabolic alkalosis.

Caution must be exercised in the administration of Lactated Ringer's Injection, USP to patients receiving corticosteroids or corticotropin.

Do not administer unless solution is clear and seal is intact.

Reactions which may occur because of the solution or the technique of administration include febrile response, infection at the site of injection, venous thrombosis or phlebitis extending from the site of injection, extravasation, and hypervolemia.

If an adverse reaction does occur, discontinue the infusion, evaluate the patient, institute appropriate therapeutic countermeasures, and save the remainder of the fluid for examination if deemed necessary.

As directed by a veterinarian. Dosage is dependent upon the age, weight and clinical condition of the patient as well as laboratory determinations.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration whenever solution and container permit.

All injections in plastic containers are intended for intravenous administration using sterile equipment.

Additives may be incompatible. Complete information is not available. Those additives known to be incompatible should not be used. Consult with pharmacist, if available. If, in the informed judgment of the veterinarian, it is deemed advisable to introduce additives, use aseptic technique. Mix thoroughly when additives have been introduced. Do not store solutions containing additives.

In an event of overhydration or solute overload, re-evaluate the patient and institute appropriate corrective measures. See Warnings, Precautions and Adverse Events.

Lactated Ringer's Injection, USP in plastic container is available as follows:

| Size (mL) | Code | NDC |
|-----------|--------|------------------|
| 5000 | 2B8219 | NDC 0338-0092-02 |

Exposure of pharmaceutical products to heat should be minimized. Avoid excessive heat. It is recommended the product be stored at room temperature (25°C); brief exposure up to 40°C does not adversely affect the product.

Directions for use of plastic container

To Open

Tear overwrap down side at slit and remove solution container. Visually inspect the container. If the outlet port protector is damaged, detached, or not present, discard container as solution path sterility may be impaired. Some opacity of the plastic due to moisture absorption during the sterilization process may be observed. This is normal and does not affect the solution quality or safety. The opacity will diminish gradually. Check for minute leaks by squeezing inner bag firmly. If leaks are found, discard solution as sterility may be impaired. If supplemental medication is desired, follow directions below.

Preparation for Administration

1. Suspend container from eyelet support.
2. Remove plastic protector from outlet port at bottom of container.
3. Attach administration set. Refer to complete directions accompanying set.

To Add Medication

WARNING: Additives may be incompatible.

To add medication before solution administration

1. Prepare medication site.
2. Using syringe with 19 to 22 gauge needle, puncture medication port and inject.
3. Mix solution and medication thoroughly. For high density medication such as potassium chloride, squeeze ports while ports are upright and mix thoroughly.

To add medication during solution administration

1. Close clamp on the set.
2. Prepare medication site.
3. Using syringe with 19 to 22 gauge needle, puncture resealable medication port and inject.
4. Remove container from IV pole and/or turn to an upright position.
5. Evacuate both ports by squeezing them while container is in the upright position.
6. Mix solution and medication thoroughly.
7. Return container to in use position and continue administration.

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

Baxter Healthcare Corporation

Deerfield, IL 60015 USA

For customer service call (800) 933-0303

Printed in USA

Baxter is a registered trademark of Baxter International, Inc.

07-19-00-0639

Rev. January 2019

4500

Veterinary Lactated Ringer's Injection USP

4 |

5000 mL

3500

EACH 100 mL CONTAINS 600 mg SODIUM CHLORIDE USP
310 mg SODIUM LACTATE 30 mg POTASSIUM CHLORIDE USP
20 mg CALCIUM CHLORIDE USP pH 6.5 (6.0 to 7.5) mEq/L
SODIUM 130 POTASSIUM 4 CALCIUM 2.7 CHLORIDE 109
LACTATE 28 OSMOLARITY 273 mOsmol/L (CALC)
STERILE NONPYROGENIC SINGLE DOSE CONTAINER CONTAINS
NO ANTIMICROBIAL AGENTS USE SOLUTION PROMPTLY
FOLLOWING INITIAL ENTRY **NOT FOR USE IN THE TREATMENT
OF LACTIC ACIDOSIS** ADDITIVES MAY BE INCOMPATIBLE
CONSULT WITH PHARMACIST IF AVAILABLE WHEN INTRODUCING
ADDITIVES USE ASEPTIC TECHNIQUE MIX THOROUGHLY DISCARD
UNUSED PORTION DOSAGE INTRAVENOUSLY AS DIRECTED BY
A VETERINARIAN SEE PACKAGE INSERT CAUTIONS SQUEEZE
AND INSPECT INNER BAG WHICH MAINTAINS PRODUCT
STERILITY DISCARD IF LEAKS ARE FOUND DO NOT ADMINISTER
SIMULTANEOUSLY WITH BLOOD DO NOT USE UNLESS SOLUTION
IS CLEAR AND SEAL IS INTACT STORE UNIT IN MOISTURE
BARRIER OVERWRAP AT ROOM TEMPERATURE (25°C/77°F)
UNTIL READY TO USE AVOID EXCESSIVE HEAT

3000

2 |

2000

FOR ANIMAL USE ONLY

1500

CAUTION FEDERAL (USA) LAW RESTRICTS THIS DRUG TO USE BY
OR ON THE ORDER OF A LICENSED VETERINARIAN

BAXTER IS A REGISTERED TRADEMARK OF
BAXTER INTERNATIONAL, INC.

FOR CUSTOMER SERVICE CALL
800 933 0303

NDC 0338-0092-02
2B8219
07-25-00-0419

1000

BAXTER HEALTHCARE CORPORATION

DEERFIELD, IL 60015 USA
MADE IN USA

Baxter

500

Container Label

**Veterinary Lactated
Ringer's Injection USP**

5000 mL

EACH 100 ML CONTAINS 600 MG SODIUM CHLORIDE USP
310 MG SODIUM LACTATE 30 MG POTASSIUM CHLORIDE USP
20 MG CALCIUM CHLORIDE USP PH 6.5 (6.0 TO 7.5) MEQ/L
SODIUM 130 POTASSIUM 4 CALCIUM 2.7 CHLORIDE 109
LACTATE 28 OSMOLARITY 273 MOSMOL/L (CALC)
STERILE NONPYROGENIC SINGLE DOSE CONTAINER CONTAINS
NO ANTIMICROBIALAGENTS USE SOLUTION PROMPTLY
FOLLOWING INITIAL ENTRY **NOT FOR USE IN THE TREATMENT
OF LACTIC ACIDOSIS** ADDITIVES MAY BE INCOMPATIBLE
CONSULT WITH PHARMACIST IF AVAILABLE WHEN INTRODUCING
ADDITIVES USE ASEPTIC TECHNIQUE MIX THOROUGHLY DISCARD
UNUSED PORTION DOSAGE INTRAVENOUSLY AS DIRECTED BY
A VETERINARIAN SEE PACKAGE INSERT CAUTIONS SQUEEZE
AND INSPECT INNER BAG WHICH MAINTAINS PRODUCT
STERILITY DISCARD IF LEAKS ARE FOUND DO NOT ADMINISTER
SIMULTANEOUSLY WITH BLOOD DO NOT USE UNLESS SOLUTION
IS CLEAR AND SEAL IS INTACT STORE UNIT IN MOISTURE
BARRIER OVERWRAP AT ROOM TEMPERATURE (25°C/77°F)
UNTIL READY TO USE AVOID EXCESSIVE HEAT

FOR ANIMAL USE ONLY

CAUTION FEDERAL (USA) LAW RESTRICTS THIS DRUG TO USE BY
OR ON THE ORDER OF A LICENSED VETERINARIAN
BAXTER IS A REGISTERED TRADEMARK OF BAXTER INTERNATIONAL, INC.

FOR CUSTOMER SERVICE CALL
800 933 0303

BAXTER HEALTHCARE CORPORATION
DEERFIELD, IL 60015 USA

MADE IN USA

NDC 0338-0092-02
2B8219
07-25-00-0419

Baxter Logo

4500

4000

3500

3000

2500

2000

1500

1000

500

VETERINARY LACTATED RINGERS

sodium chloride, sodium lactate, potassium chloride, calcium chloride injection, solution

Product Information

| | | | |
|--------------------------------|--------------------------|---------------------------|---------------|
| Product Type | PRESCRIPTION ANIMAL DRUG | Item Code (Source) | NDC:0338-0092 |
| Route of Administration | INTRAVENOUS | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|--------------------|------------------|
| SODIUM CHLORIDE (UNII: 451W47IQ8X) (CHLORIDE ION - UNII:Q32ZN48698, SODIUM CATION - UNII:LYR4MONH37) | SODIUM CHLORIDE | 600 mg in 100 mL |
| SODIUM LACTATE (UNII: TU7HW0W0QT) (LACTIC ACID - UNII:33X04XA5AT, SODIUM CATION - UNII:LYR4MONH37) | SODIUM LACTATE | 310 mg in 100 mL |
| POTASSIUM CHLORIDE (UNII: 660YQ98I10) (POTASSIUM CATION - UNII:295O53K152, CHLORIDE ION - UNII:Q32ZN48698) | POTASSIUM CHLORIDE | 30 mg in 100 mL |
| CALCIUM CHLORIDE (UNII: M4I0D6VV5M) (CALCIUM CATION - UNII:2M83C4R6ZB, CHLORIDE ION - UNII:Q32ZN48698) | CALCIUM CHLORIDE | 20 mg in 100 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|---------------------------------|----------|
| WATER (UNII: 059QF0KO0R) | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---------------------|----------------------|--------------------|
| 1 | NDC:0338-0092-02 | 2 in 1 CARTON | | |
| 1 | | 5000 mL in 1 BAG | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------------|--|----------------------|--------------------|
| unapproved drug other | | 01/01/2019 | |

Labeler - Baxter Healthcare Corporation (005083209)

Establishment

| Name | Address | ID/FEI | Business Operations |
|-------------------------------|---------|-----------|--|
| Baxter Healthcare Corporation | | 059140764 | ANALYSIS, LABEL, MANUFACTURE, PACK, STERILIZE, API MANUFACTURE |

Establishment

| Name | Address | ID/FEI | Business Operations |
|-------------------------------|---------|-----------|---------------------|
| Baxter Healthcare Corporation | | 194684502 | ANALYSIS |

Revised: 1/2019

Baxter Healthcare Corporation