CHILDRENS PAIN AND FEVER- acetaminophen suspension Meijer Distribution Inc

Meijer Distribution, Inc. Children's Pain & Fever Drug Facts

Active ingredient (in each 5 mL)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to:
 - · the common cold
 - flu
 - headache
 - sore throat
 - toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has

liver disease

Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin

When using this product

do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see overdose warning)
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child's dose into the dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL

60-71	9-10 years	12.5 mL
72-95	11 years	15 mL

^{*}or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- each 5 mL contains: sodium 3 mg
- store at 20-25°C (68-77°F)
- do not use if printed neckband is broken or missing

Inactive ingredients

anhydrous citric acid, calcium sulfate, carrageenan, flavor, glycerin, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, tribasic sodium phosphate

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

meijer®

CHILDREN'S | age 2-11 years

pain & fever

acetaminophen 160 mg per 5 mL | pain reliever fever reducer

COMPARE TO CHILDREN'S TYLENOL® ACTIVE INGREDIENT

DYE FREE


FAST, EFFECTIVE RELIEF OF CHILDREN'S PAIN & FEVER

CHERRY FLAVOR

Oral Suspension

Free of: Dyes, Alcohol, Aspirin, Ibuprofen, Parabens, High Fructose Corn Syrup

8 FL OZ (236 mL)





meijer

CHILDREN'S | age 2-11 years

pain & fever

acetaminophen 160 mg per 5 mL pain reliever | fever reduc

FAST, EFFECTIVE RELIEF OF CHILDREN'S PAIN & FEVER

CHERRY FLAVOR

NDC 41250-959-34

Drug Facts

Active ingredient (in each 5 mL)

Purpose

- Uses temporarily:

 = reduces fever
 = relivese minor aches and pains due to:
 = the common cold
 = the use the common to the temporarily the temp

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes
■ more than 5 doese in 24 hours, which is the maximum delly

I mote users of some and an anomal moteral manufacture and with other drugs containing acetaminophen

Altergy alert: Acetaminophen may cause severe skin reactions.

Symptoms may include:

If a skin reaction is a skin reaction is a skin reaction in a skin reaction is a skin reaction in a skin reaction is a skin reaction occurs, skop use and seek medical help right away, for a score threat warning! If soer throat is severe, persists for more than 2 days, is accompanied or followed by ferest, headache, rash, nausea, or vernifing, consult a doctor promptly.

- Do not use

 with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a ductor or pharmacist.

 If your child has even had an allergic reaction to this product or

Ask a doctor before use if your child has liver disea

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

When using this product do not exceed recommended dose se warning)

- Stop use and ask a doctor if

 Bein gets worse or lasts more than 5 days

 fever gets worse or lasts more than 3 days

 fever gets worse or lasts more than 3 days

 mew symptoms occur

 redness or swelling is present

 These could be signs of a serious condition.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quck medical attention is critical even if you do not notice any

- Directions

 It his product does not contain directions or complete warnings for adult use do not give more than directed (see overdose warning)

 I shake well before using

 I'ml = milliture

 I'ml right does on chart. If possible, use weight to does otherwise, use greekers cap and squeeze your child's dose into the doing cup possessive cap and squeeze your child's dose into the doing cup possessive cap and squeeze your child's dose into the doing cup possessive cap and squeeze your child's dose into the doing cup possessive cap and squeeze your child's dose into the doing cup possessive cap and squeeze your child's dose into the doing cup possessive cap and squeeze your child's dose into the doing cup possessive cap and squeeze your child's dose into the doing cup possessive cap and squeeze your child's dose into the doing cup possessive cap and squeeze your child's dose into the doing cup possessive cap and squeeze your child's dose into the doing cup possessive cap and squeeze your child's dose into the doing cup possessive cap and squeeze your child's dose into the doing cup possessive cap and squeeze your child's dose into the doing cup possessive cap and squeeze your child's dose into the doing cup possessive cap and squeeze your child's dose into the doing cup possessive cap and squeeze your child's dose into the doing cup possessive cap and squeeze your child's dose into the doing cup possessive cap and squeeze your child's dose into the doing cup possessive cap and squeeze your child's dose into the doing cup possessive cap and squeeze your child's dose into the doing cup possessive cap and squeeze your child's dose into the doing cup possessive cap and squeeze your child's dose into the doing cup possessive cap and squeeze your child's dose into the doing cup possessive cap and squeeze your child's dose into the doing cup possessive cap and squeeze your child's dose into the doing cup possessive cap and squeeze your child's dose into the doing cup possessive cap and cap and cap and cap and cap and cap an

Drug Facts (continued)

lo not give more than 5 times in 24 hours			
Weight (lb)	Age (yr)	Dose (mL)*	
under 24	under 2 years	ask a doctor	
24-35	2-3 years	5 mL	
36-47	4-5 years	7.5 mL	
48-59	6-8 years	10 mL	
00.71	0.10 mars	19 E ml	

11 years 15 mL

"or as directed by a doctor
Ittention: use only enclosed dosing cup specifically designed for
use with this product. Do not use any other dosing device.

- Other information
 each 5 mL contains: sodium 3 mg
 store at 20-25°C (68-77°F)
 do not use if printed neckband is broken or missing

Inactive ingredients anydrous citric acid, calcium sulfate, carrageman, flavor, glycerin, hydroxyethyl cellulose, microcystelline cellulose and carboxymethylcellulose sodium propylene glycol, purified water, sodium benzoate, sorbitol s sucralose, tribasic sodium phosphate

Questions or comments? 1-800-719-9260

**This product is not manufactured or distributed by Kerwue Inc., owner of the registered trademark Tylenol*.

DIST. BY MEIJER DISTRIBUTION, INC. Grand Rapids, MI 49544 www.meijer.com



meijer

CHILDREN'S | age 2-11 years

pain & fever

acetaminophen 160 mg per 5 mL | pain reliever fever reducer

COMPARE TO CHILDREN'S TYLENOL®

FAST, EFFECTIVE RELIEF OF CHILDREN'S PAIN & FEVER

CHERRY FLAVOR

Oral Suspension Free of: Dyes, Alcohol, Aspirin, Ibuprofen, Parabens, High Fructose Corn Syrup





95934 6E C1







STEP 2. Remove child protective cap



STEP 3. Squeeze your child's dose into the dosing cup



IMPORTANT: Please see dosing chart or ask a doctor for your child's correct dose





acetaminophen suspension

Product Information Product Type HUMAN OTC DRUG Item Code (Source) NDC:41250-959 **Route of Administration ORAL**

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	160 mg in 5 mL	

Inactive Ingredients			
Ingredient Name	Strength		
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)			
CALCIUM SULFATE, UNSPECIFIED FORM (UNII: WATODDB505)			
CARRAGEENAN (UNII: 5C69YCD2YJ)			
GLYCERIN (UNII: PDC6A3C0OX)			
HYDROXYETHYL CELLULOSE, UNSPECIFIED (UNII: T4V6TWG28D)			
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)			
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K6790BS311)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
WATER (UNII: 059QF0KO0R)			
SODIUM BENZOATE (UNII: OJ245FE5EU)			
SORBITOL SOLUTION (UNII: 8KW3E207O2)			
SUCRALOSE (UNII: 96K6UQ3ZD4)			
SODIUM PHOSPHATE, TRIBASIC (UNII: A752Q30A6X)			

Product Characteristics			
Color	WHITE (opaque, viscous)	Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-959- 26	1 in 1 CARTON	03/28/2012	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:41250-959- 34	1 in 1 CARTON	06/30/2025	
2		236 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	03/28/2012	

Labeler - Meijer Distribution Inc (006959555)

Revised: 7/2025 Meijer Distribution Inc